



Observation of *s*-Channel Production of Single Top Quarks at the Tevatron

- T. Aaltonen,^{†,21} V. M. Abazov,^{‡,13} B. Abbott,^{‡,116} B. S. Acharya,^{‡,80} M. Adams,^{‡,98} T. Adams,^{‡,97} J. P. Agnew,^{‡,94}
G. D. Alexeev,^{‡,13} G. Alkhazov,^{‡,88} A. Alton,^{‡,31,ii} S. Amerio,^{‡,39a,39b} D. Amidei,^{‡,31} A. Anastassov,^{‡,15,v} A. Annovi,^{‡,17}
J. Antos,^{‡,12} G. Apolinari,^{‡,15} J. A. Appel,^{‡,15} T. Arisawa,^{‡,52} A. Artikov,^{‡,13} J. Asaadi,^{‡,47} W. Ashmanskas,^{‡,15}
A. Askew,^{‡,97} S. Atkins,^{‡,106} B. Auerbach,^{‡,2} K. Augsten,^{‡,62} A. Aurisano,^{‡,47} C. Avila,^{‡,60} F. Azfar,^{‡,38} F. Badaud,^{‡,65}
W. Badgett,^{‡,15} T. Bae,^{‡,25} L. Bagby,^{‡,15} B. Baldin,^{‡,15} D. V. Bandurin,^{‡,51} S. Banerjee,^{‡,80} A. Barbaro-Galtieri,^{‡,26}
E. Barberis,^{‡,107} P. Baringer,^{‡,105} V. E. Barnes,^{‡,43} B. A. Barnett,^{‡,23} P. Barria,^{‡,41a,41c} J. F. Bartlett,^{‡,15} P. Bartos,^{‡,12}
U. Bassler,^{‡,70} M. Bauche,^{‡,39a,39b} V. Bazterra,^{‡,98} A. Bean,^{‡,105} F. Bedeschi,^{‡,41} M. Begalli,^{‡,57} S. Behari,^{‡,15}
L. Bellantoni,^{‡,15} G. Bellettini,^{‡,41a,41b} J. Bellinger,^{‡,54} D. Benjamin,^{‡,14} A. Beretvas,^{‡,15} S. B. Beri,^{‡,78} G. Bernardi,^{‡,69}
R. Bernhard,^{‡,74} I. Bertram,^{‡,92} M. Besançon,^{‡,70} R. Beuselinck,^{‡,93} P. C. Bhat,^{‡,15} S. Bhatia,^{‡,108} V. Bhatnagar,^{‡,78}
A. Bhatti,^{‡,45} K. R. Bland,^{‡,5} G. Blazey,^{‡,99} S. Blessing,^{‡,97} K. Bloom,^{‡,109} B. Blumenfeld,^{‡,23} A. Bocci,^{‡,14} A. Bodek,^{‡,44}
A. Boehnlein,^{‡,15} D. Boline,^{‡,113} E. E. Boos,^{‡,86} G. Borissov,^{‡,92} D. Bortoletto,^{‡,43} M. Borysova,^{‡,91,tt} J. Boudreau,^{‡,42}
A. Boveia,^{‡,11} A. Brandt,^{‡,119} O. Brandt,^{‡,75} L. Brigliadori,^{‡,6a,6b} R. Brock,^{‡,32} C. Bromberg,^{‡,32} A. Bross,^{‡,15}
D. Brown,^{‡,69} E. Brucken,^{‡,21} X. B. Bu,^{‡,15} J. Budagov,^{‡,13} H. S. Budd,^{‡,44} M. Buehler,^{‡,15} V. Buescher,^{‡,76}
V. Bunichev,^{‡,86} S. Burdin,^{‡,92,ij} K. Burkett,^{‡,15} G. Busetto,^{‡,39a,39b} P. Bussey,^{‡,19} C. P. Buszello,^{‡,90} P. Butti,^{‡,41a,41b}
A. Buzatu,^{‡,19} A. Calamba,^{‡,10} E. Camacho-Pérez,^{‡,83} S. Camarda,^{‡,4} M. Campanelli,^{‡,28} F. Canelli,^{‡,11,cc} B. Carls,^{‡,22}
D. Carlsmith,^{‡,54} R. Carosi,^{‡,41} S. Carrillo,^{‡,16,l} B. Casal,^{‡,9,j} M. Casarsa,^{‡,48} B. C. K. Casey,^{‡,15} H. Castilla-Valdez,^{‡,83}
A. Castro,^{‡,6a,6b} P. Catastini,^{‡,20} S. Caughron,^{‡,32} D. Cauz,^{‡,48a,48b,48c} V. Cavaliere,^{‡,22} M. Cavalli-Sforza,^{‡,4} A. Cerri,^{‡,26,e}
L. Cerrito,^{‡,28,q} S. Chakrabarti,^{‡,113} K. M. Chan,^{‡,103} A. Chandra,^{‡,121} E. Chapon,^{‡,70} G. Chen,^{‡,105} Y. C. Chen,^{‡,1}
M. Chertok,^{‡,7} G. Chiarelli,^{‡,41} G. Chlachidze,^{‡,15} K. Cho,^{‡,25} S. W. Cho,^{‡,82} S. Choi,^{‡,82} D. Chokheli,^{‡,13}
B. Choudhary,^{‡,79} S. Cihangir,^{‡,15} D. Claes,^{‡,109} A. Clark,^{‡,18} C. Clarke,^{‡,53} J. Clutter,^{‡,105} M. E. Convery,^{‡,15} J. Conway,^{‡,7}
M. Cooke,^{‡,15,ss} W. E. Cooper,^{‡,15,y} M. Corbo,^{‡,15,y} M. Corcoran,^{‡,121} M. Cordelli,^{‡,17} F. Couderc,^{‡,70} M.-C. Cousinou,^{‡,67}
C. A. Cox,^{‡,7} D. J. Cox,^{‡,7} M. Cremonesi,^{‡,41} D. Cruz,^{‡,47} J. Cuevas,^{‡,9,x} R. Culbertson,^{‡,15} D. Cutts,^{‡,118} A. Das,^{‡,95}
N. d'Ascenzo,^{‡,15,u} M. Datta,^{‡,15,ff} G. Davies,^{‡,93} P. de Barbaro,^{‡,44} S. J. de Jong,^{‡,84,85} E. De La Cruz-Burelo,^{‡,83}
F. Déliot,^{‡,70} R. Demina,^{‡,44} L. Demortier,^{‡,45} M. Deninno,^{‡,6} D. Denisov,^{‡,15} S. P. Denisov,^{‡,87} M. D'Errico,^{‡,39a,39b}
S. Desai,^{‡,15} C. Deterre,^{‡,75,kk} K. DeVaughan,^{‡,109} F. Devoto,^{‡,21} A. Di Canto,^{‡,41a,41b} B. Di Ruzza,^{‡,15,p} H. T. Diehl,^{‡,15}
M. Diesburg,^{‡,15} P. F. Ding,^{‡,94} J. R. Dittmann,^{‡,5} A. Dominguez,^{‡,109} S. Donati,^{‡,41a,41b} M. D'Onofrio,^{‡,27}
M. Dorigo,^{‡,48a,48d} A. Driutti,^{‡,48a,48b,48c} A. Dubey,^{‡,79} L. V. Dudko,^{‡,86} A. Duperrin,^{‡,67} S. Dutt,^{‡,78} M. Eads,^{‡,99}
K. Ebina,^{‡,52} R. Edgar,^{‡,31} D. Edmunds,^{‡,32} A. Elagin,^{‡,47} J. Ellison,^{‡,96} V. D. Elvira,^{‡,15} Y. Enari,^{‡,69} R. Erbacher,^{‡,7}
S. Errede,^{‡,22} B. Esham,^{‡,22} H. Evans,^{‡,101} V. N. Evdokimov,^{‡,87} S. Farrington,^{‡,38} L. Feng,^{‡,99} T. Ferbel,^{‡,44}
J. P. Fernández Ramos,^{‡,29} F. Fiedler,^{‡,76} R. Field,^{‡,16} F. Filthaut,^{‡,84,85} W. Fisher,^{‡,32} H. E. Fisk,^{‡,15} G. Flanagan,^{‡,15,s}
R. Forrest,^{‡,7} M. Fortner,^{‡,99} H. Fox,^{‡,92} M. Franklin,^{‡,20} J. C. Freeman,^{‡,15} H. Frisch,^{‡,11} S. Fuess,^{‡,15} Y. Funakoshi,^{‡,52}
C. Galloni,^{‡,41a,41b} P. H. Garbincius,^{‡,15} A. Garcia-Bellido,^{‡,44} J. A. García-González,^{‡,83} A. F. Garfinkel,^{‡,43}
P. Garosi,^{‡,41a,41c} V. Gavrilov,^{‡,33} W. Geng,^{‡,67,32} C. E. Gerber,^{‡,98} H. Gerberich,^{‡,22} E. Gerchtein,^{‡,15} Y. Gershtein,^{‡,110}
S. Giagu,^{‡,46} V. Giakoumopoulou,^{‡,3} K. Gibson,^{‡,42} C. M. Ginsburg,^{‡,15} G. Ginther,^{‡,15,44} N. Giokaris,^{‡,3} P. Giromini,^{‡,17}
G. Giurgiu,^{‡,23} V. Glagolev,^{‡,13} D. Glenzinski,^{‡,15} M. Gold,^{‡,34} D. Goldin,^{‡,47} A. Golosanov,^{‡,15} G. Golovanov,^{‡,13}
G. Gomez,^{‡,9} G. Gomez-Ceballos,^{‡,30} M. Goncharov,^{‡,30} O. González López,^{‡,29} I. Gorelov,^{‡,34} A. T. Goshaw,^{‡,14}
K. Goulianos,^{‡,45} E. Gramellini,^{‡,6} P. D. Grannis,^{‡,113} S. Greder,^{‡,71} H. Greenlee,^{‡,15} G. Grenier,^{‡,72} S. Grinstein,^{‡,4}
Ph. Gris,^{‡,65} J.-F. Grivaz,^{‡,68} A. Grohsjean,^{‡,70,kk} C. Grossi-Pilcher,^{‡,11} R. C. Group,^{‡,51,15} S. Grünendahl,^{‡,15}
M. W. Grünewald,^{‡,81} T. Guillemin,^{‡,68} J. Guimaraes da Costa,^{‡,20} G. Gutierrez,^{‡,15} P. Gutierrez,^{‡,116} S. R. Hahn,^{‡,15}
J. Haley,^{‡,117} J. Y. Han,^{‡,44} L. Han,^{‡,59} F. Happacher,^{‡,17} K. Hara,^{‡,49} K. Harder,^{‡,94} M. Hare,^{‡,50} A. Harel,^{‡,44}
R. F. Harr,^{‡,53} T. Harrington-Taber,^{‡,15,m} K. Hatakeyama,^{‡,5} J. M. Hauptman,^{‡,104} C. Hays,^{‡,38} J. Hays,^{‡,93} T. Head,^{‡,94}
T. Hebbeker,^{‡,73} D. Hedin,^{‡,99} H. Hegab,^{‡,117} J. Heinrich,^{‡,40} A. P. Heinson,^{‡,96} U. Heintz,^{‡,118} C. Hensel,^{‡,56}
I. Heredia-De La Cruz,^{‡,83,ii} M. Herndon,^{‡,54} K. Herner,^{‡,15} G. Hesketh,^{‡,94,nn} M. D. Hildreth,^{‡,103} R. Hirosky,^{‡,51}
T. Hoang,^{‡,97} J. D. Hobbs,^{‡,113} A. Hocker,^{‡,15} B. Hoeneisen,^{‡,64} J. Hogan,^{‡,121} M. Hohlfeld,^{‡,76} J. L. Holzbauer,^{‡,108}
Z. Hong,^{‡,47} W. Hopkins,^{‡,15,f} S. Hou,^{‡,1} I. Howley,^{‡,119} Z. Hubacek,^{‡,62,70} R. E. Hughes,^{‡,35} U. Husemann,^{‡,55}
M. Hussein,^{‡,32,aa} J. Huston,^{‡,32} V. Hynek,^{‡,62} I. Iashvili,^{‡,112} Y. Ilchenko,^{‡,120} R. Illingworth,^{‡,15} G. Introzzi,^{‡,41a,41e,41f}
M. Iori,^{‡,46a,46b} A. S. Ito,^{‡,15} A. Ivanov,^{‡,7,0} S. Jabeen,^{‡,118} M. Jaffré,^{‡,68} E. James,^{‡,15} D. Jang,^{‡,10} A. Jayasinghe,^{‡,116}
B. Jayatilaka,^{‡,15} E. J. Jeon,^{‡,25} M. S. Jeong,^{‡,82} R. Jesik,^{‡,93} P. Jiang,^{‡,59} S. Jindariani,^{‡,15} K. Johns,^{‡,95} E. Johnson,^{‡,32}

- M. Johnson,^{‡,15} A. Jonckheere,^{‡,15} M. Jones,^{‡,43} P. Jonsson,^{‡,93} K. K. Joo,^{‡,25} J. Joshi,^{‡,96} S. Y. Jun,^{‡,10} A. W. Jung,^{‡,15}
 T. R. Junk,^{‡,15} A. Juste,^{‡,89} E. Kajfasz,^{‡,67} M. Kambeitz,^{‡,24} T. Kamon,^{‡,25,47} P. E. Karchin,^{‡,53} D. Karmanov,^{‡,86}
 A. Kasmi,^{‡,5} Y. Kato,^{‡,37,n} I. Katsanos,^{‡,109} R. Kehoe,^{‡,120} S. Kermiche,^{‡,67} W. Ketchum,^{‡,11,gg} J. Keung,^{‡,40}
 N. Khalatyan,^{‡,15} A. Khanov,^{‡,117} A. Kharchilava,^{‡,112} Y. N. Kharzheev,^{‡,13} B. Kilminster,^{‡,15,cc} D. H. Kim,^{‡,25}
 H. S. Kim,^{‡,25} J. E. Kim,^{‡,25} M. J. Kim,^{‡,17} S. H. Kim,^{‡,49} S. B. Kim,^{‡,25} Y. J. Kim,^{‡,25} Y. K. Kim,^{‡,11} N. Kimura,^{‡,52}
 M. Kirby,^{‡,15} I. Kiselevich,^{‡,33} K. Knoepfel,^{‡,15} J. M. Kohli,^{‡,78} K. Kondo,^{‡,52,*} D. J. Kong,^{‡,25} J. Konigsberg,^{‡,16}
 A. V. Kotwal,^{‡,14} A. V. Kozelov,^{‡,87} J. Kraus,^{‡,108} M. Kreps,^{‡,24} J. Kroll,^{‡,40} M. Kruse,^{‡,14} T. Kuhr,^{‡,24} A. Kumar,^{‡,112}
 A. Kupco,^{‡,63} M. Kurata,^{‡,49} T. Kurča,^{‡,72} V. A. Kuzmin,^{‡,86} A. T. Laasanen,^{‡,43} S. Lammel,^{‡,15} S. Lammers,^{‡,101}
 M. Lancaster,^{‡,28} K. Lannon,^{‡,35,w} G. Latino,^{‡,41a,41c} P. Lebrun,^{‡,72} H. S. Lee,^{‡,82} H. S. Lee,^{‡,25} J. S. Lee,^{‡,25} S. W. Lee,^{‡,104}
 W. M. Lee,^{‡,15} X. Lei,^{‡,95} J. Lellouch,^{‡,69} S. Leo,^{‡,41} S. Leone,^{‡,41} J. D. Lewis,^{‡,15} D. Li,^{‡,69} H. Li,^{‡,51} L. Li,^{‡,96}
 Q. Z. Li,^{‡,15} J. K. Lim,^{‡,82} A. Limosani,^{‡,14,r} D. Lincoln,^{‡,15} J. Linnemann,^{‡,32} V. V. Lipaev,^{‡,87} E. Lipeles,^{‡,40}
 R. Lipton,^{‡,15} A. Lister,^{‡,18,a} H. Liu,^{‡,51} H. Liu,^{‡,120} Q. Liu,^{‡,43} T. Liu,^{‡,15} Y. Liu,^{‡,59} A. Lobodenko,^{‡,88} S. Lockwitz,^{‡,55}
 A. Loginov,^{‡,55} M. Lokajicek,^{‡,63} R. Lopes de Sa,^{‡,113} D. Lucchesi,^{‡,39a,39b} A. Lucă,^{‡,17} J. Lueck,^{‡,24} P. Lujan,^{‡,26}
 P. Lukens,^{‡,15} R. Luna-Garcia,^{‡,83,oo} G. Lungu,^{‡,45} A. L. Lyon,^{‡,15} J. Lys,^{‡,26} R. Lysak,^{‡,12,d} A. K. A. Maciel,^{‡,56}
 R. Madar,^{‡,74} R. Madrak,^{‡,15} P. Maestro,^{‡,41a,41c} R. Magaña-Villalba,^{‡,83} S. Malik,^{‡,45} S. Malik,^{‡,109} V. L. Malyshev,^{‡,13}
 G. Manca,^{‡,27,b} A. Manousakis-Katsikakis,^{‡,3} J. Mansour,^{‡,75} L. Marchese,^{‡,6,hh} F. Margaroli,^{‡,46} P. Marino,^{‡,41a,41d}
 J. Martínez-Ortega,^{‡,83} M. Martínez,^{‡,4} K. Matera,^{‡,22} M. E. Mattson,^{‡,53} A. Mazzacane,^{‡,15} P. Mazzanti,^{‡,6}
 R. McCarthy,^{‡,113} C. L. McGivern,^{‡,94} R. McNulty,^{‡,27,i} A. Mehta,^{‡,27} P. Mehtala,^{‡,21} M. M. Meijer,^{‡,84,85}
 A. Melnitchouk,^{‡,15} D. Menezes,^{‡,99} P. G. Mercadante,^{‡,58} M. Merkin,^{‡,86} C. Mesropian,^{‡,45} A. Meyer,^{‡,73} J. Meyer,^{‡,75,qq}
 T. Miao,^{‡,15} F. Miconi,^{‡,71} D. Mietlicki,^{‡,31} A. Mitra,^{‡,1} H. Miyake,^{‡,49} S. Moed,^{‡,15} N. Moggi,^{‡,6} N. K. Mondal,^{‡,80}
 C. S. Moon,^{‡,15,y} R. Moore,^{‡,15,dd,ee} M. J. Morello,^{‡,41a,41d} A. Mukherjee,^{‡,15} M. Mulhearn,^{‡,51} Th. Muller,^{‡,24} P. Murat,^{‡,15}
 M. Mussini,^{‡,6a,6b} J. Nachtman,^{‡,15,m} Y. Nagai,^{‡,49} J. Naganoma,^{‡,52} E. Nagy,^{‡,67} I. Nakano,^{‡,36} A. Napier,^{‡,50}
 M. Narain,^{‡,118} R. Nayyar,^{‡,95} H. A. Neal,^{‡,31} J. P. Negret,^{‡,60} J. Nett,^{‡,47} C. Neu,^{‡,51} P. Neustroev,^{‡,88} H. T. Nguyen,^{‡,51}
 T. Nigmanov,^{‡,42} L. Nodulman,^{‡,2} S. Y. Noh,^{‡,25} O. Norniella,^{‡,22} T. Nunnemann,^{‡,77} L. Oakes,^{‡,38} S. H. Oh,^{‡,14}
 Y. D. Oh,^{‡,25} I. Oksuzian,^{‡,51} T. Okusawa,^{‡,37} R. Orava,^{‡,21} J. Orduna,^{‡,121} L. Ortolan,^{‡,4} N. Osman,^{‡,67} J. Osta,^{‡,103}
 C. Pagliarone,^{‡,48} A. Pal,^{‡,119} E. Palencia,^{‡,9,e} P. Palni,^{‡,34} V. Papadimitriou,^{‡,15} N. Parashar,^{‡,102} V. Parihar,^{‡,118}
 S. K. Park,^{‡,82} W. Parker,^{‡,54} R. Partridge,^{‡,118,mm} N. Parua,^{‡,101} A. Patwa,^{‡,114,rr} G. Paulett,^{‡,48a,48b,48c} M. Paulini,^{‡,10}
 C. Paus,^{‡,30} B. Penning,^{‡,15} M. Perfilov,^{‡,86} Y. Peters,^{‡,94} K. Petridis,^{‡,44} G. Petrillo,^{‡,68} P. Pétroff,^{‡,14} T. J. Phillips,^{‡,14}
 G. Piacentino,^{‡,41} E. Pianori,^{‡,40} J. Pilot,^{‡,7} K. Pitts,^{‡,22} C. Plager,^{‡,8} M.-A. Pleier,^{‡,114} V. M. Podstavkov,^{‡,15}
 L. Pondrom,^{‡,54} A. V. Popov,^{‡,87} S. Poprocki,^{‡,15,f} K. Potamianos,^{‡,26} A. Pranko,^{‡,26} M. Prewitt,^{‡,121} D. Price,^{‡,94}
 N. Prokopenko,^{‡,87} F. Prokoshin,^{‡,13,z} F. Ptohos,^{‡,17,g} G. Punzi,^{‡,41a,41b} J. Qian,^{‡,31} A. Quadt,^{‡,75} B. Quinn,^{‡,108}
 N. Ranjan,^{‡,43} P. N. Ratoff,^{‡,92} I. Razumov,^{‡,87} I. Redondo Fernández,^{‡,29} P. Renton,^{‡,38} M. Rescigno,^{‡,46} F. Rimondi,^{‡,6,*}
 I. Ripp-Baudot,^{‡,71} L. Ristori,^{‡,41,15} F. Rizatdinova,^{‡,117} A. Robson,^{‡,19} T. Rodriguez,^{‡,40} S. Rolli,^{‡,50,h} M. Rominsky,^{‡,15}
 M. Ronzani,^{‡,41a,41b} R. Roser,^{‡,15} J. L. Rosner,^{‡,11} A. Ross,^{‡,92} C. Royon,^{‡,70} P. Rubinov,^{‡,15} R. Ruchti,^{‡,103}
 F. Ruffini,^{‡,41a,41c} A. Ruiz,^{‡,9} J. Russ,^{‡,10} V. Rusu,^{‡,15} G. Sajot,^{‡,66} W. K. Sakamoto,^{‡,44} Y. Sakurai,^{‡,52}
 A. Sánchez-Hernández,^{‡,83} M. P. Sanders,^{‡,77} L. Santi,^{‡,48a,48b,48c} A. S. Santos,^{‡,56,pp} K. Sato,^{‡,49} G. Savage,^{‡,15}
 V. Saveliev,^{‡,15,u} A. Savoy-Navarro,^{‡,15,y} L. Sawyer,^{‡,106} T. Scanlon,^{‡,93} R. D. Schamberger,^{‡,113} Y. Scheglov,^{‡,88}
 H. Schellman,^{‡,100} P. Schlabach,^{‡,15} E. E. Schmidt,^{‡,15} C. Schwanenberger,^{‡,94} T. Schwarz,^{‡,31} R. Schwienhorst,^{‡,32}
 L. Scodellaro,^{‡,9} F. Scuri,^{‡,41} S. Seidel,^{‡,34} Y. Seiya,^{‡,37} J. Sekaric,^{‡,105} A. Semenov,^{‡,13} H. Severini,^{‡,116} F. Sforza,^{‡,41a,41b}
 E. Shabalina,^{‡,75} S. Z. Shalhout,^{‡,7} V. Shary,^{‡,70} S. Shaw,^{‡,32} A. A. Shchukin,^{‡,87} T. Shears,^{‡,27} P. F. Shepard,^{‡,42}
 M. Shimojima,^{‡,49,t} M. Shochet,^{‡,11} I. Shreyber-Tecker,^{‡,33} V. Simak,^{‡,62} A. Simonenko,^{‡,13} P. Skubic,^{‡,116} P. Slattery,^{‡,44}
 K. Sliwa,^{‡,50} D. Smirnov,^{‡,103} J. R. Smith,^{‡,7} F. D. Snider,^{‡,15} G. R. Snow,^{‡,109} J. Snow,^{‡,115} S. Snyder,^{‡,114}
 S. Söldner-Rembold,^{‡,94} H. Song,^{‡,42} L. Sonnenschein,^{‡,73} V. Sorin,^{‡,4} K. Soustruznik,^{‡,61} R. St. Denis,^{‡,19,*}
 M. Stancari,^{‡,15} J. Stark,^{‡,66} D. Stentz,^{‡,15,v} D. A. Stoyanova,^{‡,87} M. Strauss,^{‡,116} J. Strologas,^{‡,34} Y. Sudo,^{‡,49}
 A. Sukhanov,^{‡,15} I. Suslov,^{‡,13} L. Suter,^{‡,94} P. Svoisky,^{‡,116} K. Takemasa,^{‡,49} Y. Takeuchi,^{‡,49} J. Tang,^{‡,11} M. Tecchio,^{‡,31}
 P. K. Teng,^{‡,1} J. Thom,^{‡,15,f} E. Thomson,^{‡,40} V. Thukral,^{‡,47} M. Titov,^{‡,70} D. Toback,^{‡,47} S. Tokar,^{‡,12} V. V. Tokmenin,^{‡,13}
 K. Tollefson,^{‡,32} T. Tomura,^{‡,49} D. Tonelli,^{‡,15,e} S. Torre,^{‡,17} D. Torretta,^{‡,15} P. Totaro,^{‡,41a,41d} M. Trovato,^{‡,44} Y.-T. Tsai,^{‡,44}
 D. Tsbychev,^{‡,113} B. Tuchming,^{‡,70} C. Tully,^{‡,111} F. Ukegawa,^{‡,49} S. Uozumi,^{‡,25} L. Uvarov,^{‡,88} S. Uvarov,^{‡,88}
 S. Uzunyan,^{‡,99} R. Van Kooten,^{‡,101} W. M. van Leeuwen,^{‡,84} N. Varelas,^{‡,98} E. W. Varnes,^{‡,95} I. A. Vasilyev,^{‡,87}
 F. Vázquez,^{‡,16,l} G. Velev,^{‡,15} C. Vellidis,^{‡,15} A. Y. Verkhhee,^{‡,13} C. Vernieri,^{‡,41a,41d} L. S. Vertogradov,^{‡,13}
 M. Verzocchi,^{‡,15} M. Vesterinen,^{‡,94} M. Vidal,^{‡,43} D. Vilanova,^{‡,70} R. Vilar,^{‡,9} J. Vizán,^{‡,9,bb} M. Vogel,^{‡,34} P. Vokac,^{‡,62}

G. Volpi,^{†,17} P. Wagner,^{†,40} H. D. Wahl,^{‡,97} R. Wallny,^{†,15,j} M. H. L. S. Wang,^{‡,15} S. M. Wang,^{†,1} J. Warchol,^{‡,103}
D. Waters,^{†,28} G. Watts,^{‡,122} M. Wayne,^{‡,103} J. Weichert,^{‡,76} L. Welty-Rieger,^{‡,100} W. C. Wester III,^{†,15} D. Whiteson,^{‡,40,c}
A. B. Wicklund,^{‡,2} S. Wilbur,^{†,7} H. H. Williams,^{†,40} M. R. J. Williams,^{‡,101} G. W. Wilson,^{‡,105} J. S. Wilson,^{‡,31}
P. Wilson,^{‡,15} B. L. Winer,^{†,35} P. Wittich,^{†,15,f} M. Wobisch,^{‡,106} S. Wolbers,^{‡,15} H. Wolfe,^{‡,35} D. R. Wood,^{‡,107}
T. Wright,^{†,31} X. Wu,^{†,18} Z. Wu,^{†,5} T. R. Wyatt,^{‡,94} Y. Xie,^{‡,15} R. Yamada,^{‡,15} K. Yamamoto,^{‡,37} D. Yamato,^{‡,37}
S. Yang,^{‡,59} T. Yang,^{†,15} U. K. Yang,^{†,25} Y. C. Yang,^{†,25} W.-M. Yao,^{‡,26} T. Yasuda,^{‡,15} Y. A. Yatsunenko,^{‡,13}
W. Ye,^{‡,113} Z. Ye,^{‡,15} G. P. Yeh,^{†,15,m} K. Yi,^{†,15,m} H. Yin,^{‡,15} K. Yip,^{‡,114} J. Yoh,^{†,15} K. Yorita,^{†,52} T. Yoshida,^{‡,37,k}
S. W. Youn,^{‡,15} G. B. Yu,^{†,14} I. Yu,^{†,25} J. M. Yu,^{‡,31} A. M. Zanetti,^{†,48} Y. Zeng,^{†,14} J. Zennamo,^{‡,112}
T. G. Zhao,^{‡,94} B. Zhou,^{‡,31} C. Zhou,^{†,14} J. Zhu,^{‡,31} M. Zielinski,^{‡,44} D. Ziemska,^{‡,101}
L. Zivkovic,^{‡,69} and S. Zucchelli^{‡,6a,6b}
(CDF Collaboration)[†]

(D0 Collaboration)[‡]¹Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China²Argonne National Laboratory, Argonne, Illinois 60439, USA³University of Athens, 157 71 Athens, Greece⁴Institut de Fisica d'Altes Energies, ICREA, Universitat Autonoma de Barcelona, E-08193 Bellaterra (Barcelona), Spain⁵Baylor University, Waco, Texas 76798, USA^{6a}Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy^{6b}University of Bologna, I-40127 Bologna, Italy⁷University of California, Davis, Davis, California 95616, USA⁸University of California, Los Angeles, Los Angeles, California 90024, USA⁹Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain¹⁰Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA¹¹Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA¹²Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia¹³Joint Institute for Nuclear Research, RU-141980 Dubna, Russia¹⁴Duke University, Durham, North Carolina 27708, USA¹⁵Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA¹⁶University of Florida, Gainesville, Florida 32611, USA¹⁷Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy¹⁸University of Geneva, CH-1211 Geneva 4, Switzerland¹⁹Glasgow University, Glasgow G12 8QQ, United Kingdom²⁰Harvard University, Cambridge, Massachusetts 02138, USA²¹Division of High Energy Physics, Department of Physics, University of Helsinki, FIN-00014, Helsinki, Finland;

Helsinki Institute of Physics, FIN-00014 Helsinki, Finland

²²University of Illinois, Urbana, Illinois 61801, USA²³The Johns Hopkins University, Baltimore, Maryland 21218, USA²⁴Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany²⁵Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea; Ewha Womans University, Seoul 120-750, Korea²⁶Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA²⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom²⁸University College London, London WC1E 6BT, United Kingdom²⁹Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain³⁰Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA³¹University of Michigan, Ann Arbor, Michigan 48109, USA³²Michigan State University, East Lansing, Michigan 48824, USA³³Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia³⁴University of New Mexico, Albuquerque, New Mexico 87131, USA³⁵The Ohio State University, Columbus, Ohio 43210, USA³⁶Okayama University, Okayama 700-8530, Japan³⁷Osaka City University, Osaka 558-8585, Japan³⁸University of Oxford, Oxford OX1 3RH, United Kingdom^{39a}Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy

- ^{39b}*University of Padova, I-35131 Padova, Italy*
- ⁴⁰*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ^{41a}*Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy*
- ^{41b}*University of Pisa, I-56127 Pisa, Italy*
- ^{41c}*University of Siena, I-56127 Pisa, Italy*
- ^{41d}*Scuola Normale Superiore, I-56127 Pisa, Italy*
- ^{41e}*INFN Pavia, I-27100 Pavia, Italy*
- ^{41f}*University of Pavia, I-27100 Pavia, Italy*
- ⁴²*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- ⁴³*Purdue University, West Lafayette, Indiana 47907, USA*
- ⁴⁴*University of Rochester, Rochester, New York 14627, USA*
- ⁴⁵*The Rockefeller University, New York, New York 10065, USA*
- ^{46a}*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, I-00185 Roma, Italy*
- ^{46b}*Sapienza Università di Roma, I-00185 Roma, Italy*
- ⁴⁷*Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA*
- ^{48a}*Istituto Nazionale di Fisica Nucleare Trieste, I-33100 Udine, Italy*
- ^{48b}*Gruppo Collegato di Udine, I-33100 Udine, Italy*
- ^{48c}*University of Udine, I-33100 Udine, Italy*
- ^{48d}*University of Trieste, I-34127 Trieste, Italy*
- ⁴⁹*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁵⁰*Tufts University, Medford, Massachusetts 02155, USA*
- ⁵¹*University of Virginia, Charlottesville, Virginia 22906, USA*
- ⁵²*Waseda University, Tokyo 169, Japan*
- ⁵³*Wayne State University, Detroit, Michigan 48201, USA*
- ⁵⁴*University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁵⁵*Yale University, New Haven, Connecticut 06520, USA*
- ⁵⁶*LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*
- ⁵⁷*Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil*
- ⁵⁸*Universidade Federal do ABC, Santo André, Brazil*
- ⁵⁹*University of Science and Technology of China, Hefei, People's Republic of China*
- ⁶⁰*Universidad de los Andes, Bogotá, Colombia*
- ⁶¹*Charles University, Faculty of Mathematics and Physics, Center for Particle Physics, Prague, Czech Republic*
- ⁶²*Czech Technical University in Prague, Prague, Czech Republic*
- ⁶³*Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic*
- ⁶⁴*Universidad San Francisco de Quito, Quito, Ecuador*
- ⁶⁵*LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France*
- ⁶⁶*LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France*
- ⁶⁷*CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France*
- ⁶⁸*LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France*
- ⁶⁹*LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France*
- ⁷⁰*CEA, Irfu, SPP, Saclay, France*
- ⁷¹*IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France*
- ⁷²*IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France*
- ⁷³*III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany*
- ⁷⁴*Physikalisches Institut, Universität Freiburg, Freiburg, Germany*
- ⁷⁵*II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany*
- ⁷⁶*Institut für Physik, Universität Mainz, Mainz, Germany*
- ⁷⁷*Ludwig-Maximilians-Universität München, München, Germany*
- ⁷⁸*Panjab University, Chandigarh, India*
- ⁷⁹*Delhi University, Delhi, India*
- ⁸⁰*Tata Institute of Fundamental Research, Mumbai, India*
- ⁸¹*University College Dublin, Dublin, Ireland*
- ⁸²*Korea Detector Laboratory, Korea University, Seoul, Korea*
- ⁸³*CINVESTAV, Mexico City, Mexico*
- ⁸⁴*Nikhef, Science Park, Amsterdam, The Netherlands*
- ⁸⁵*Radboud University Nijmegen, Nijmegen, The Netherlands*
- ⁸⁶*Moscow State University, Moscow, Russia*
- ⁸⁷*Institute for High Energy Physics, Protvino, Russia*
- ⁸⁸*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*
- ⁸⁹*Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Física d'Altes Energies (IFAE), Barcelona, Spain*

- ⁹⁰*Uppsala University, Uppsala, Sweden*
⁹¹*Taras Shevchenko National University of Kyiv, Kiev, Ukraine*
⁹²*Lancaster University, Lancaster LA1 4YB, United Kingdom*
⁹³*Imperial College London, London SW7 2AZ, United Kingdom*
⁹⁴*The University of Manchester, Manchester M13 9PL, United Kingdom*
⁹⁵*University of Arizona, Tucson, Arizona 85721, USA*
⁹⁶*University of California Riverside, Riverside, California 92521, USA*
⁹⁷*Florida State University, Tallahassee, Florida 32306, USA*
⁹⁸*University of Illinois at Chicago, Chicago, Illinois 60607, USA*
⁹⁹*Northern Illinois University, DeKalb, Illinois 60115, USA*
¹⁰⁰*Northwestern University, Evanston, Illinois 60208, USA*
¹⁰¹*Indiana University, Bloomington, Indiana 47405, USA*
¹⁰²*Purdue University Calumet, Hammond, Indiana 46323, USA*
¹⁰³*University of Notre Dame, Notre Dame, Indiana 46556, USA*
¹⁰⁴*Iowa State University, Ames, Iowa 50011, USA*
¹⁰⁵*University of Kansas, Lawrence, Kansas 66045, USA*
¹⁰⁶*Louisiana Tech University, Ruston, Louisiana 71272, USA*
¹⁰⁷*Northeastern University, Boston, Massachusetts 02115, USA*
¹⁰⁸*University of Mississippi, University, Mississippi 38677, USA*
¹⁰⁹*University of Nebraska, Lincoln, Nebraska 68588, USA*
¹¹⁰*Rutgers University, Piscataway, New Jersey 08855, USA*
¹¹¹*Princeton University, Princeton, New Jersey 08544, USA*
¹¹²*State University of New York, Buffalo, New York 14260, USA*
¹¹³*State University of New York, Stony Brook, New York 11794, USA*
¹¹⁴*Brookhaven National Laboratory, Upton, New York 11973, USA*
¹¹⁵*Langston University, Langston, Oklahoma 73050, USA*
¹¹⁶*University of Oklahoma, Norman, Oklahoma 73019, USA*
¹¹⁷*Oklahoma State University, Stillwater, Oklahoma 74078, USA*
¹¹⁸*Brown University, Providence, Rhode Island 02912, USA*
¹¹⁹*University of Texas, Arlington, Texas 76019, USA*
¹²⁰*Southern Methodist University, Dallas, Texas 75275, USA*
¹²¹*Rice University, Houston, Texas 77005, USA*
¹²²*University of Washington, Seattle, Washington 98195, USA*

(Received 21 February 2014; published 9 June 2014)

We report the first observation of single-top-quark production in the s channel through the combination of the CDF and D0 measurements of the cross section in proton-antiproton collisions at a center-of-mass energy of 1.96 TeV. The data correspond to total integrated luminosities of up to 9.7 fb^{-1} per experiment. The measured cross section is $\sigma_s = 1.29^{+0.26}_{-0.24} \text{ pb}$. The probability of observing a statistical fluctuation of the background to a cross section of the observed size or larger is 1.8×10^{-10} , corresponding to a significance of 6.3 standard deviations for the presence of an s -channel contribution to the production of single-top quarks.

DOI: 10.1103/PhysRevLett.112.231803

PACS numbers: 14.65.Ha, 12.15.Hh, 12.15.Ji, 13.85.Qk

The top quark, with a mass of $m_t = 173.2 \pm 0.9 \text{ GeV}$ [1], is the most massive and one of the most puzzling elementary particles of the standard model (SM). Detailed studies of top-quark production and decay provide powerful tests of strong and electroweak interactions, as well as sensitivity to physics beyond the standard model (BSM) [2]. At the Tevatron, where protons (p) and antiprotons (\bar{p}) collide at a center-of-mass energy of $\sqrt{s} = 1.96 \text{ TeV}$, top quarks are produced predominantly in pairs ($t\bar{t}$) via the strong interaction [3]. Top quarks are also produced singly in $p\bar{p}$ collisions via the electroweak interaction. The single-top-quark production cross section is expected to be proportional to the square of the magnitude of

the quark-mixing Cabibbo-Kobayashi-Maskawa matrix [4] element V_{tb} , and consequently sensitive to potential contributions from a fourth generation of quarks [5,6], as well as flavor-changing neutral currents [7–10], anomalous top-quark couplings [11–13], heavy W' bosons [14–17], charged Higgs bosons [18,19], or other new phenomena [20,21].

At the Tevatron, there are two important processes in which a single top quark is produced in association with other quarks. The dominant channel proceeds through the exchange of a spacelike virtual W boson between a light quark and a bottom quark (b quark) in the t channel [22–24]. A second mode occurs through the exchange of a




FIG. 1 (color online). Dominant Feynman diagrams for (a) s -channel and (b) t -channel single-top-quark production at the Tevatron.

timelike virtual W boson in the s channel, which produces a top quark and a b quark [25]. Figure 1 shows the leading Feynman diagrams for the s - and t -channel production modes. Independent measurements of s -channel and t -channel production are important, since BSM contributions could have different effects on the two modes [20].

Single-top-quark production, independent of channel, was reported by the CDF and D0 Collaborations in Refs. [26,27] and [28,29], respectively. The D0 Collaboration subsequently measured with larger data sets the production cross section for the combined s and t channels [30], and obtained $\sigma_{s+t} = 4.11^{+0.60}_{-0.55}$ pb using a data set of 9.7 fb^{-1} in agreement with the SM prediction of 3.15 ± 0.19 pb ($m_t = 172.5 \text{ GeV}$) [24,31].

After establishing the $s + t$ process, the cross sections of the individual production modes were measured independently. Several differences in the properties of s - and t -channel events can be used to distinguish them from one another. Events originating from t -channel production typically contain one light-flavor jet in the forward detector region (at large pseudorapidity), which is useful for distinguishing them from events associated with s -channel production and other SM background processes. Moreover, events from the s -channel process are more likely to contain two jets originating from b quarks (b jets) within the central region of the detector where they can be identified. Hence, single-top-like events with two identified b jets are more likely to have originated from s -channel production. Exploiting these differences, the D0 Collaboration observed the t -channel process [32], and measured its cross section to be $\sigma_t = 3.07^{+0.54}_{-0.55}$ pb using a data set of 9.7 fb^{-1} [30]. This compares to the SM prediction of 2.10 ± 0.13 pb ($m_t = 172.5 \text{ GeV}$) [24]. At the CERN LHC proton-proton (pp) collider, t -channel production was also observed by the ATLAS and CMS Collaborations [33,34].

Observing the s -channel process is more difficult, since the expected cross section is smaller than that of the t channel and its kinematic features are less distinct from the background. However, the Tevatron has an advantage over the LHC in this mode, since valence quarks ($q\bar{q}'$ from $p\bar{p}$) generally initiate s -channel single-top-quark production, leading to a larger signal-to-background ratio at the Tevatron than at the LHC. Due to this advantage, the

CDF and D0 Collaborations have reported evidence for s -channel production independently of each other [30,35], while the LHC experiments have to date reported only unpublished upper limits on the cross section in pp collisions.

In this Letter, we report a combination of s -channel cross section analyses performed by the CDF [35,36] and D0 [30] Collaborations. The CDF and D0 detectors are central magnetic spectrometers surrounded by electromagnetic and hadronic calorimeters and muon detectors [37–39]. The combined measurement utilizes the full Tevatron Run II data sets corresponding to up to 9.7 fb^{-1} of integrated luminosity per experiment.

The data are selected using a logical OR of many online selection requirements, which preserve high signal efficiency for offline analysis. Since the magnitude of the W -top-bottom quark coupling is much larger than the W -top-down and W -top-strange quark couplings [40], each top quark decays almost exclusively to a W boson and a b quark. The selection is split into two distinct final-state topologies, both designed to select single-top-quark events in which the W boson decays leptonically.

One final-state topology ($\ell + \text{jets}$, $\ell = e$ or μ), analyzed by both collaborations, contains single-top-quark events in which the W boson decays leptonically producing an electron or a muon. We select events that (i) contain only one isolated lepton ℓ with large transverse momentum p_T , (ii) have large missing transverse energy E_T [39], (iii) have either two jets (CDF analysis) or two or three jets (D0 analysis) with large p_T , and (iv) have one or two b jets. To identify b jets, multivariate techniques are used that discriminate b jets from jets originating from light quarks and gluons [41,42]. Additional selection criteria are applied to exclude kinematic regions that are difficult to model, and to minimize the quantum chromodynamics (QCD) multijet background where one jet is misreconstructed as a lepton and spurious E_T arises from jet energy mismeasurements.

The other final-state topology, analyzed by the CDF Collaboration, involves E_T and jets, but no reconstructed isolated charged leptons ($E_T + \text{jets}$). The CDF analysis avoids overlap with the $\ell + \text{jets}$ sample by explicitly vetoing events with identified leptons [36]. Large missing transverse energy is required and events with two or three reconstructed jets are accepted. This additional sample increases the acceptance for s -channel signal events by encompassing those in which the W -boson decay produces a muon or electron that is either not reconstructed or not isolated, or a hadronically decaying tau lepton that is reconstructed as a third jet. After the basic event selection, QCD multijet events dominate the $E_T + \text{jets}$ event sample. To reduce this multijet background, a neural-network event selection is optimized to preferentially select signallike events.

Events passing the $\ell + \text{jets}$ and $E_T + \text{jets}$ selections are further separated into independent analysis channels based on the number of reconstructed jets as well as the number

and quality of b -tagged jets. Each of the analyzed channels has a different background composition and signal (s) to background (b) ratio. Analyzing them separately enhances the sensitivity to single-top-quark production [30,35,36].

Both collaborations use Monte Carlo (MC) generators to simulate the kinematic properties of signal and background events, except in the case of multijet production, for which the model is derived from data. The CDF analysis models single-top-quark signal events at next-to-leading-order (NLO) accuracy in the strong coupling constant α_s using the POWHEG [43] generator. The D0 analysis uses the SINGLETOP [44] event generator, based on NLO COMPHEP calculations that match the event kinematic features predicted by NLO calculations [45,46]. Spin information in the decays of the top quark and the W boson is preserved for both POWHEG and SINGLETOP.

Kinematic properties of background events associated with the $W + \text{jets}$ and $Z + \text{jets}$ processes are simulated using the ALPGEN leading-order MC generator [47], and those of diboson processes (WW , WZ , and ZZ) are modeled using PYTHIA [48]. The $t\bar{t}$ process is modeled using PYTHIA in the CDF analysis and by ALPGEN in the D0 analysis. Higgs-boson processes are modeled using simulated events generated with PYTHIA for a Higgs-boson mass of $m_H = 125$ GeV. In all cases PYTHIA is used to model proton remnants and simulate the hadronization of all generated partons. The mass of the top quark in simulated events is set to $m_t = 172.5$ GeV, which is consistent with the current Tevatron average value [1]. All MC events are processed through GEANT-based detector simulations [49] and reconstructed by the same software packages used for the collider data.

Predictions for the normalization of simulated background-process contributions are estimated using both simulation and data. Data are used to normalize the W plus light-flavor and heavy-flavor jet contributions using enriched $W + \text{jets}$ data samples that have negligible signal content [27,30,36]. All other simulated background samples are normalized to the theoretical cross sections at NLO combined with next-to-next-to-leading log (NNLL) resummation [24] for t -channel single-top-quark production, at next-to-NLO [50] for $t\bar{t}$, at NLO [51] for $Z + \text{jets}$ and diboson production, and including all relevant higher-order QCD and electroweak corrections for Higgs-boson production [52]. Differences observed between simulated events and data in lepton and jet reconstruction efficiencies, resolutions, jet-energy scale (JES), and b -tagging efficiencies are adjusted in the simulation to match the data, through correction functions obtained from measurements in independent data samples.

We form multivariate discriminants, optimized for separating the s -channel single-top-quark signal events in each of the analysis samples from the larger background contributions, to extract the cross section measurements [53]. The combined cross section measurement is obtained using a Bayesian statistical analysis of each bin of the observed

TABLE I. Systematic uncertainties associated with the CDF and the D0 single-top-quark s -channel cross section measurements in $\ell + \text{jets}$ final states. The values shown for each category indicate the range of uncertainties applied to the predicted normalizations for signal and background contributions over the full set of analysis samples from each experiment. The check marks indicate which categories contribute uncertainties on the shape of the final multivariate discriminant output variable. It is also noted if categories are treated as fully correlated between the two experiments.

Systematic uncertainty	CDF Norm(%)	D0 Norm(%)	Corre- lated Dist		
Lumi from detector	4.5	4.5	No		
Lumi from cross section	4.0	4.0	Yes		
Signal modeling	2–10	✓	3–8	Yes	
Background (simulation)	2–12	✓	2–11	✓	Yes
Background (data)	15–40	✓	19–50	✓	No
Detector modeling	2–10	✓	1–5	✓	No
b -jet-tagging	10–30		5–40	✓	No
JES	0–20	✓	0–40	✓	No

discriminant distribution from each sample, comparing data to the modeled distributions for each of the contributing signal and background processes [54].

A complete list of systematic uncertainties for the $\ell + \text{jets}$ analyses is given in Table I. These can arise from uncertainties on differential distributions (Dist) and their normalizations (Norm). The CDF $E_T + \text{jets}$ analysis has a similar set of systematic uncertainties that are fully correlated with the CDF $\ell + \text{jets}$ analysis except for the uncertainty related to the data-based background. Sources of systematic uncertainty common to measurements of both collaborations are assumed to be 100% correlated, while other uncertainties are assumed to be uncorrelated. The dependence of the results on the correlation assumptions has been found to be negligible. The categories of uncertainty correspond generally to those in Refs. [1,3], and can be summarized as follows.

Detector-specific luminosity uncertainty: The component of the uncertainty on integrated luminosity that comes from the uncertainty on the acceptance and efficiency of the luminosity detector is taken as uncorrelated between the CDF [55] and D0 [56] measurements.

Luminosity from cross section: The portion of the uncertainty in integrated luminosity that comes from uncertainties on the inelastic and diffractive cross sections is fully correlated between the CDF and D0 measurements.

Signal modeling: The systematic uncertainty associated with uncertainties in the modeling of the single-top-quark signal, including uncertainties from the choice of the description of initial- and final-state QCD radiation, and proton and antiproton parton density functions, also covering uncertainties in the applied hadronization models, is taken as fully correlated between the CDF and D0 measurements.




FIG. 2 (color online). Distribution of the discriminant histograms, summed for bins with similar signal-to-background ratio (s/b). The expected sum of the backgrounds is shown by the unfilled histogram, and the uncertainty of the background is represented by the gray shaded band. The expected s -channel signal contribution is shown by a filled blue histogram.

Background from simulation: The systematic uncertainty associated with uncertainties in the modeling of various background contributions is taken as fully correlated between the CDF and D0 measurements. This includes uncertainties in $t\bar{t}$ and diboson process normalizations originating from theoretical calculations.

Background based on data: The systematic uncertainty associated with the modeling of various background sources obtained using data-driven methods is uncorrelated




FIG. 3 (color online). The background-subtracted distribution of the discriminant histograms, summed for bins with similar signal-to-background ratio (s/b). The background and s -channel signal (blue filled histogram) have been normalized to the most likely values returned from the likelihood fit and the uncertainty on the background uncertainty has also been constrained by the data (gray shaded band).

between the CDF and D0 measurements. This includes uncertainties on the normalization of $W + \text{jets}$, $Wb\bar{b}$, and $Wc\bar{c}$ events as well as uncertainties on the modeling of the contributions and discriminant-variable shapes for the $W + \text{jets}$ and QCD multijet production processes.

Detector modeling: The systematic uncertainty on efficiencies for identifying reconstructed objects and to cover observed mismodeling of the data from the simulations is uncorrelated between the CDF and D0 measurements.

b-jet tagging: The systematic uncertainty associated with the modeling of b -jet tagging efficiencies and associated mistag rates is uncorrelated between the CDF [41] and D0 [42] measurements.

Jet energy scale: This systematic uncertainty originates from using calibration-data samples to establish the JES. For the CDF analyses, this corresponds to uncertainties associated with the η -dependent JES corrections, which are estimated using dijet events in data. For the D0 analysis, this includes uncertainties in calorimeter response for light jets, uncertainties from η - and p_T -dependent JES corrections, and other small contributions. This uncertainty is assumed to be uncorrelated between the CDF [57] and D0 [58] measurements.

The Bayesian posterior probability density as a function of the s -channel signal cross section σ_s is given by

$$p(\sigma_s) = \int L(\sigma_s, \{\theta\} | \text{data}) \pi(\sigma_s) \Pi(\{\theta\}) d\{\theta\}, \quad (1)$$

where L is the joint binned likelihood function for all channels

$$L = \prod_{i=\text{bins,channels}} \frac{(s_i + b_i)^{n_i} e^{-(s_i + b_i)}}{n_i!}. \quad (2)$$




FIG. 4 (color online). The posterior probability distribution for the combination of the CDF and D0 analysis channels compared with the NLO + NNLL theoretical prediction [31].




FIG. 5 (color online). Measured single-top-quark s -channel production cross sections from each of the individual analyses and various combinations of these analyses compared with the NLO + NNLL theoretical prediction [31].

The number of observed events in bin i is n_i . $\{\theta\}$ is the set of nuisance parameters representing the systematic uncertainties (assuming priors following a Gaussian probability density function), and $\Pi(\{\theta\})$ is the product of the prior probability densities encoding the systematic uncertainties on $\{\theta\}$. The predictions for the number of signal events s_i and background events b_i depend on the values of the nuisance parameters that are integrated over in Eq. (1). The prior density for the signal cross section, $\pi(\sigma_s)$, is taken to be a uniform prior for non-negative cross sections. We quote the measured cross section as the value that maximizes its posterior likelihood, and the uncertainty as the smallest interval that contains 68% of the integrated area of the posterior density.

Figure 2 shows the signal and background expectations and the data as a function of $\log_{10}(s/b)$ of the collected bins, for the combined CDF and D0 analyses. The respective background-subtracted $\log_{10}(s/b)$ discriminant distribution using the most likely values for the signal and background yields derived from the likelihood fit is shown in Fig. 3. The extracted posterior probability distribution for σ_s is presented in Fig. 4, and Fig. 5 gives a graphical presentation of the individual and combined measurements. All measurements agree within their uncertainties with the SM prediction $\sigma_s^{\text{SM}} = 1.05 \pm 0.06 \text{ pb}$ ($m_t = 172.5 \text{ GeV}$) [31]. The most probable value for the combined cross section is $\sigma_s = 1.29^{+0.26}_{-0.24} \text{ pb}$ for a top-quark mass of 172.5 GeV. The total expected uncertainty is 20%, and the expected uncertainty without considering systematic uncertainties is 14%. The dependence of the measured value on the assumed value of the top-quark mass is negligible compared to the uncertainty on the measurement [27,30].




FIG. 6 (color online). Log-likelihood ratios using an asymptotic approximation for the background-only (solid green line) and SM-signal-plus-background (dashed blue) hypotheses from the combined measurement.

The statistical significance of this result is quantified through a calculated p value based on an asymptotic log-likelihood ratio (LLR) approach [59], including systematic uncertainties. The p value quantifies the probability that the measured value of the cross section or a larger value could result from a background fluctuation in the absence of signal. The distributions of the LLR resulting from fits of simulated samples that include background-only, or signal-plus-background, contributions are presented in Fig. 6. The probability to measure an s -channel cross section of at least the observed value in the absence of signal is 1.8×10^{-10} , corresponding to a significance of 6.3 standard deviations (s.d.), with a sensitivity expected, assuming the central value of the SM s -channel cross-section prediction, of 5.1 s.d.

In summary, we report the first observation of s -channel single-top-quark production with a significance of 6.3 s.d. by combining the CDF and D0 measurements. The combined value of the s -channel single-top-quark production cross section is $\sigma_s = 1.29^{+0.26}_{-0.24} \text{ pb}$, in agreement with the SM expectation.

We thank the Fermilab staff and technical staffs of the participating institutions for their vital contributions. We acknowledge support from the U.S. DOE and NSF (U.S.), ARC (Australia), CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil), NSERC (Canada), NSC, CAS, and CNSF (China), Colciencias (Colombia), MSMT and GACR (Czech Republic), the Academy of Finland, CEA and CNRS/IN2P3 (France), BMBF and DFG (Germany), DAE and DST (India), SFI (Ireland), INFN (Italy), MEXT (Japan), the Korean World Class University Program and NRF (Korea), CONACyT (Mexico), FOM (Netherlands), MON, NRC KI, and RFBR (Russia), the Slovak R&D Agency, the Ministerio de Ciencia e Innovación, and

Programa Consolider-Ingenio 2010 (Spain), The Swedish Research Council (Sweden), SNSF (Switzerland), STFC and the Royal Society (United Kingdom), the A. P. Sloan Foundation (U.S.), and the EU community Marie Curie Fellowship Contract No. 302103.

^{*}Deceased.

^aVisitor from University of British Columbia, Vancouver, BC V6T 1Z1, Canada.

^bVisitor from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy.

^cVisitor from University of California Irvine, Irvine, California 92697, USA.

^dVisitor from Institute of Physics, Academy of Sciences of the Czech Republic, 182 21, Czech Republic.

^eVisitor from CERN, CH-1211 Geneva, Switzerland.

^fVisitor from Cornell University, Ithaca, New York 14853, USA.

^gVisitor from University of Cyprus, Nicosia CY-1678, Cyprus.

^hVisitor from Office of Science, U.S. Department of Energy, Washington, DC 20585, USA.

ⁱVisitor from University College Dublin, Dublin 4, Ireland.

^jVisitor from ETH, 8092 Zürich, Switzerland.

^kVisitor from University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017.

^lVisitor from Universidad Iberoamericana, Lomas de Santa Fe, México, C.P. 01219, Distrito Federal.

^mVisitor from University of Iowa, Iowa City, Iowa 52242, USA.

ⁿVisitor from Kinki University, Higashi-Osaka City, Japan 577-8502.

^oVisitor from Kansas State University, Manhattan, Kansas 66506, USA.

^pVisitor from Brookhaven National Laboratory, Upton, New York 11973, USA.

^qVisitor from Queen Mary, University of London, London, E1 4NS, United Kingdom.

^rVisitor from University of Melbourne, Victoria 3010, Australia.

^sVisitor from Muons, Inc., Batavia, Illinois 60510, USA.

^tVisitor from Nagasaki Institute of Applied Science, Nagasaki 851-0193, Japan.

^uVisitor from National Research Nuclear University, Moscow 115409, Russia.

^vVisitor from Northwestern University, Evanston, Illinois 60208, USA.

^wVisitor from University of Notre Dame, Notre Dame, Indiana 46556, USA.

^xVisitor from Universidad de Oviedo, E-33007 Oviedo, Spain.

^yVisitor from CNRS-IN2P3, Paris, F-75205 France.

^zVisitor from Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile.

^{aa}Visitor from The University of Jordan, Amman 11942, Jordan.

^{bb}Visitor from Universite catholique de Louvain, 1348 Louvain-La-Neuve, Belgium.

^{cc}Visitor from University of Zürich, 8006 Zürich, Switzerland.

^{dd}Visitor from Massachusetts General Hospital, Boston, Massachusetts 02114, USA.

^{ee}Visitor from Harvard Medical School, Boston, Massachusetts 02114, USA.

^{ff}Visitor from Hampton University, Hampton, Virginia 23668, USA.

^{gg}Visitor from Los Alamos National Laboratory, Los Alamos, New Mexico 87544, USA.

^{hh}Visitor from Università degli Studi di Napoli Federico I, I-80138 Napoli, Italy.

ⁱⁱVisitor from Augustana College, Sioux Falls, South Dakota, USA.

^{jj}Visitor from The University of Liverpool, Liverpool, United Kingdom.

^{kk}Visitor from DESY, Hamburg, Germany.

^{ll}Visitor from Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Mexico.

^{mm}Visitor from SLAC, Menlo Park, California, USA.

ⁿⁿVisitor from University College London, London, United Kingdom.

^{oo}Visitor from Centro de Investigacion en Computacion - IPN, Mexico City, Mexico.

^{pp}Visitor from Universidade Estadual Paulista, São Paulo, Brazil.

^{qq}Visitor from Karlsruher Institut für Technologie (KIT) - Steinbuch Centre for Computing (SCC).

^{rr}Visitor from Office of Science, U.S. Department of Energy, Washington, D.C. 20585, USA.

^{ss}Visitor from American Association for the Advancement of Science, Washington, D.C. 20005, USA.

^{tt}Visitor from National Academy of Science of Ukraine (NASU) - Kiev Institute for Nuclear Research (KINR).

[1] T. Aaltonen *et al.* (CDF and D0 Collaborations), *Phys. Rev. D* **86**, 092003 (2012); arXiv:1305.3929.

[2] C. T. Hill and S. J. Parke, *Phys. Rev. D* **49**, 4454 (1994).

[3] T. Aaltonen *et al.* (CDF and D0 Collaborations), *Phys. Rev. D* **89**, 072001 (2014).

[4] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).

[5] M. S. Chanowitz, *Phys. Rev. D* **79**, 113008 (2009).

[6] J. Alwall, R. Frederix, J.-M. Gerard, A. Giannanco, M. Herquet, S. Kalinin, E. Kou, V. Lemaitre, and F. Maltoni, *Eur. Phys. J. C* **49**, 791 (2007).

[7] T. M. P. Tait and C. P. Yuan, *Phys. Rev. D* **55**, 7300 (1997).

[8] M. E. Luke and M. J. Savage, *Phys. Lett. B* **307**, 387 (1993).

[9] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **102**, 151801 (2009).

[10] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **693**, 81 (2010).

[11] A. P. Heinson, A. S. Belyaev, and E. E. Boos, *Phys. Rev. D* **56**, 3114 (1997).

[12] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **708**, 21 (2012).

[13] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **713**, 165 (2012).

[14] E. H. Simmons, *Phys. Rev. D* **55**, 5494 (1997).

[15] A. Datta, P. J. O'Donnell, Z. H. Lin, X. Zhang, and T. Huang, *Phys. Lett. B* **483**, 203 (2000).

[16] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **103**, 041801 (2009).

- [17] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **699**, 145 (2011).
- [18] C. S. Li, R. J. Oakes, and J. M. Yang, *Phys. Rev. D* **55**, 1672 (1997).
- [19] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **102**, 191802 (2009).
- [20] T. M. P. Tait and C. P. Yuan, *Phys. Rev. D* **63**, 014018 (2000).
- [21] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **108**, 201802 (2012).
- [22] S. S. D. Willenbrock and D. A. Dicus, *Phys. Rev. D* **34**, 155 (1986).
- [23] C.-P. Yuan, *Phys. Rev. D* **41**, 42 (1990).
- [24] N. Kidonakis, *Phys. Rev. D* **83**, 091503 (2011). The cross section for the t -channel single-top-quark process is $2.10 \pm 0.13 \text{ pb}$ ($m_t = 172.5 \text{ GeV}$).
- [25] S. Cortese and R. Petronzio, *Phys. Lett. B* **253**, 494 (1991).
- [26] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **103**, 092002 (2009).
- [27] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **82**, 112005 (2010).
- [28] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **103**, 092001 (2009).
- [29] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **84**, 112001 (2011).
- [30] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **726**, 656 (2013).
- [31] N. Kidonakis, *Phys. Rev. D* **81**, 054028 (2010). The s -channel single-top-quark cross section is $1.05 \pm 0.06 \text{ pb}$ ($m_t = 172.5 \text{ GeV}$).
- [32] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **705**, 313 (2011).
- [33] G. Aad *et al.* (ATLAS Collaboration), *Phys. Lett. B* **717**, 330 (2012).
- [34] S. Chatrchyan *et al.* (CMS Collaboration), *J. High Energy Phys.* **12** (2012) 035.
- [35] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **112**, 231804 (2014).
- [36] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **112**, 231805 (2014).
- [37] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005).
- [38] V. M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006).
- [39] The CDF and D0 experiments use cylindrical coordinate systems with origins in the centers of the detectors, where θ and ϕ are the polar and azimuthal angles, respectively, and pseudorapidity is $\eta = -\ln[\tan(\theta/2)]$. The transverse energy, as measured by the calorimeter, is defined to be $E_T = E \sin \theta$. The missing E_T (\vec{E}_T) is defined by $\vec{E}_T = -\sum_i E_T^i \hat{n}_i$, i = calorimeter tower number, where \hat{n}_i is a unit vector perpendicular to the beam axis and pointing at the i th calorimeter tower. \vec{E}_T is corrected for high-energy muons and for mismeasurements of jet energies. We define $E_T = |\vec{E}_T|$. The transverse momentum p_T is defined to be $p \sin \theta$.
- [40] J. Beringer *et al.* [Particle Data Group Collaboration], *Phys. Rev. D* **86**, 010001 (2012).
- [41] J. Freeman, T. Junk, M. Kirby, Y. Oksuzian, T. Phillips, F. Snider, M. Trovato, J. Vizan, and W. Yao, *Nucl. Instrum. Methods Phys. Res., Sect. A* **697**, 64 (2013).
- [42] V. M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **620**, 490 (2010); V. M. Abazov *et al.* (D0 Collaboration), arXiv:1312.7623.
- [43] S. Alioli, P. Nason, C. Oleari, and E. Re, *J. High Energy Phys.* **09** (2009) 111.
- [44] E. E. Boos, V. E. Bunichev, L. V. Dudko, V. I. Savrin, and V. V. Sherstnev, *Phys. At. Nucl.* **69**, 1317 (2006). We use SINGLETOP version 4.2p1.
- [45] Z. Sullivan, *Phys. Rev. D* **70**, 114012 (2004).
- [46] J. M. Campbell, R. Frederix, F. Maltoni, and F. Tramontano, *Phys. Rev. Lett.* **102**, 182003 (2009).
- [47] M. L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A. D. Polosa, *J. High Energy Phys.* **07** (2003) 001. We use ALPGEN version 2.11.
- [48] T. Sjöstrand, S. Mrenna, and P. Skands, *J. High Energy Phys.* **05** (2006) 026. We use PYTHIA version 6.409.
- [49] R. Brun and F. Carminati, CERN Program Library Long Writeup, Report No. W5013, 1993.
- [50] S. Moch and P. Uwer, *Phys. Rev. D* **78**, 034003 (2008). The top-quark-pair cross section is $7.46 \pm 0.75 \text{ pb}$ ($m_t = 172.5 \text{ GeV}$).
- [51] R. K. Ellis, *Nucl. Phys. B, Proc. Suppl.* **160**, 170 (2006). We use MCFM version 5.1.
- [52] J. Baglio and A. Djouadi, *J. High Energy Phys.* **10** (2010) 064.
- [53] For the D0 Collaboration this is different from Ref. [30], where a combined s - and t -channel discriminant was used to measure both single top quark channels separately, without assuming their SM cross sections. In this Letter, we use a discriminant where the s channel is the only signal and the t channel is considered as background normalized to its SM cross section.
- [54] I. Bertram *et al.*, Report No. FERMILAB-TM-2104 (2000).
- [55] S. Klimenko, J. Konigsberg, and T. M. Liss, Report No. FERMILAB-FN-0741.
- [56] T. Andeen *et al.*, Report No. FERMILAB-TM-2365 (2007).
- [57] A. Bhatti *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **566**, 375 (2006).
- [58] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **85**, 052006 (2012).
- [59] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, *Eur. Phys. J. C* **71**, 1554 (2011).