

Journal Pre-proof

Recent progress on wastewater-based epidemiology for COVID-19 surveillance: A systematic review of analytical procedures and epidemiological modeling

Stefano Ciannella, Cristina González-Fernández, Jenifer Gomez-Pastora



PII: S0048-9697(23)01569-3

DOI: <https://doi.org/10.1016/j.scitotenv.2023.162953>

Reference: STOTEN 162953

To appear in: *Science of the Total Environment*

Received date: 1 February 2023

Revised date: 13 March 2023

Accepted date: 15 March 2023

Please cite this article as: S. Ciannella, C. González-Fernández and J. Gomez-Pastora, Recent progress on wastewater-based epidemiology for COVID-19 surveillance: A systematic review of analytical procedures and epidemiological modeling, *Science of the Total Environment* (2023), <https://doi.org/10.1016/j.scitotenv.2023.162953>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Recent progress on wastewater-based epidemiology for COVID-19 surveillance: a systematic review of analytical procedures and epidemiological modeling

Stefano Ciannella¹

sciannel@ttu.edu

Cristina González-Fernández^{1,2}

gon72807@ttu.edu

Jenifer Gomez-Pastora^{1,*}

jenifer.gomez@ttu.edu

¹ Department of Chemical Engineering, Texas Tech University, Lubbock TX, USA 79409

² Departamento de Ingenierías Química y Biomolecular, Universidad de Cantabria, Avda. Los Castros, s/n, 39005 Santander, Spain

* **Corresponding author:** Email: jenifer.gomez@ttu.edu; Phone: +1 806 834 3553

Abstract

On March 11, 2020, the World Health Organization declared the coronavirus disease 2019 (COVID-19), whose causative agent is the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), a pandemic. This virus is predominantly transmitted via respiratory droplets and shed via sputum, saliva, urine, and stool. Wastewater-based epidemiology (WBE) has been able to monitor the circulation of viral pathogens in the population. This tool demands both in-lab and computational work to be meaningful for, among other purposes, the prediction of outbreaks. In this context, we present a systematic review that organizes and discusses laboratory procedures for SARS-CoV-2 RNA quantification from a wastewater matrix, along with modeling techniques applied to the development of WBE for COVID-19 surveillance. The goal of this review is to present the current panorama of WBE operational aspects as well as to identify current challenges related to it. Our review was conducted in a reproducible manner by following the PRISMA guidelines for systematic reviews. We identified a lack of standardization in wastewater analytical procedures. Regardless, the real-time quantitative polymerase chain reaction (RT-qPCR) approach was the most reported technique employed to detect and quantify viral RNA in wastewater samples. As a more convenient sample matrix, we suggest the solid portion of wastewater to be considered in future investigations due to its higher viral load compared to the liquid fraction. Regarding the epidemiological modeling, the data-driven approach was consistently used for the prediction of variables associated with outbreaks. Future efforts should also be directed toward the development of rapid, more economical, portable, and accurate detection devices.

Keywords: COVID-19; wastewater-based epidemiology; SARS-CoV-2 detection; systematic review; wastewater; epidemiological modeling.

1 INTRODUCTION

The Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has caused a major pandemic where millions of people have been infected globally. It belongs to the *Coronaviridae* family and comprises spiked glycoproteins (S) on the surface of a spherical virion that varies from 60 to 140 nm in diameter and is surrounded by a lipid envelope (Zhu et al., 2020). Particularly, SARS-CoV-2 is the causative agent of the coronavirus disease 2019 (COVID-19), which is a life-threatening disease that represents a major threat to public health (Bar-Or et al., 2021; Flood et al., 2021; Li et al., 2020). SARS-CoV-2 is predominantly transmitted via respiratory droplets, which are generated during sneezing, breathing or coughing and direct or indirect contact through different secretions (Tanhaei et al., 2021; van Doremalen et al., 2020). In this regard, this virus has not only been detected in sputum and saliva but its RNA has been also found in stools and urine, as well as in anal/rectal swabs (Cheung et al., 2020; Mesoraca et al., 2020; Peng et al., 2020). Rather than testing individuals, wastewater-based epidemiology (WBE) has been applied to detect viral pathogens in sewage shed from stool and urine, thus representing a viable alternative to estimate the infection prevalence in the community. WBE was theorized in 2001 (Jones-Lepp, 2001) with the original purpose of monitoring the use of illicit drugs at the community level (Claro et al., 2021). Recently, it has been successfully applied for the detection and monitoring of several viral pathogens in the population (e.g., poliovirus, enterovirus, norovirus, and hepatitis) (Barbosa et al., 2022; Hellmer et al., 2014; Medema et al., 2020; Nasseri et al., 2021; Robotto et al., 2022).

It has been proven that SARS-CoV-2 can be shed in feces after its replication in human intestine enterocytes (Ding and Liang, 2020; Haramoto et al., 2020; Lamers et al., 2020; Lescure et al., 2020), even when the patient had no gastrointestinal symptoms (Xiao et al., 2020a; Zuo et al., 2021). The shedding of this virus from stools can occur after becoming undetectable in the respiratory tract (Wu et al., 2020). Thereby, the SARS-CoV-2 shedding period was found to be longer in fecal than in upper respiratory samples, but its RNA is generally detected earlier in the latter (Zhang et al., 2021a). Additionally, it has been reported that SARS-CoV-2 RNA could be shed through respiratory and fecal routes before the infected individual exhibits symptoms (Buscarini et al., 2020; He et al., 2020; Zhang et al., 2021b). According to the above-mentioned insights, shedding in feces, sputum and saliva contribute to the SARS-CoV-2 load in wastewater (Markt et al., 2021). Interestingly, however, the analysis of wastewater performed through cell culture indicated that the SARS-CoV-2 particles were found non-infectious (Tiwari et al., 2022), bringing evidence to previous observations suggesting that SARS-CoV-2 is not potentially associated with a waterborne transmission risk in community wastewater influents (Rimoldi et al., 2020; Westhaus et al., 2021). The detection of SARS-CoV-2 in wastewater (or, interchangeably, sewage), even at low COVID-19 prevalence, makes sewage surveillance a sensitive tool to monitor its circulation in the population (Prakash, 2021). Quantifying a specific genome of an enteric virus in wastewater is an indirect, noninvasive form of assessing the current health status of the local population (Prevost et al., 2015). Moreover, wastewater surveillance enables both providing early notice of the SARS-CoV-2 (re)emergence in a population when applied routinely (Karthikeyan et

al., 2021; Zhao et al., 2022), and supplementing clinical testing by assessing temporal and spatial trends, evaluating asymptomatic and symptomatic individuals, and observing the efficiency of public preventive strategies (Castiglioni et al., 2022; Gupta et al., 2020; Tomasino et al., 2021).

The development of analytical methods for WBE purposes, starting from sampling and viral detection to RNA quantification, emerges as an important research theme that has been approached by a considerable number of studies in the last two years (Carducci et al., 2020; Kabdasli and Tunay, 2021; Kitajima et al., 2020). In this context, a dearth of standardization in the sample analysis methodology was identified (Ahmed et al., 2021; Calderon-Franco et al., 2022; McMinn et al., 2021; Peinado et al., 2022), which has been characterized by the use of a myriad of methods to concentrate, extract, detect, and quantify SARS-CoV-2 RNA (de Sousa et al., 2022; Pillay et al., 2021; Xie et al., 2022). Furthermore, the normalization of quantitative information has not been addressed: for example, standard units to express viral loads in wastewater have not been established so far (Shah et al., 2022). Regarding the output of the laboratory analysis, the accurate estimation of viral genomic concentration in wastewater is an issue that must be addressed in future COVID-19 surveillance research since this variable has been used to estimate the number of COVID-19 cases when confronted with clinical testing data (Ahmed et al., 2020a; de Sousa et al., 2022; Pillay et al., 2021). Analytical accuracy is imperative for building the path towards understanding the infection dynamics through WBE by designing trustful correlations and mathematical models relating sewershed viral concentration and epidemiological clinical data. To address this issue, we elevated the need to systematize the available

knowledge on the technology for wastewater analysis as well as the scientific effort to unravel and model COVID-19 infection dynamics through existing or developed WBE mathematical models.

The purpose of this systematic review is twofold: identify the reported methodology of techniques/procedures to quantify SARS-CoV-2 viral RNA in domestic wastewater, and the mathematical methods and models by which viral loads have been associated with epidemiological data. It should be noted that this integrated approach has not been considered by any previously published review on the field. Thus, this study may serve as a reference for upcoming research that requires detailed information on these subjects, thus readers interested in one or both operational aspects of WBE for COVID-19 can find this study relevant given the exposition of methods and findings from a total of 158 studies. This review is structured as follows. After this introduction, Section 2 brings a comprehensive description of the systematic search method and selection process for evidence-based publications discussing analytical methods and mathematical modeling for COVID-19 surveillance up to August 2022. Next, in Section 3, we report our findings from the selected literature regarding in-lab and computational works that have been performed. Section 4 critically discusses our findings in terms of current issues, gaps to fill, and promising alternatives to treat wastewater towards the refinement of WBE for COVID-19 surveillance. Finally, we finish with the main conclusions drawn from this analysis and directions for further research.

2 METHODS

2.1 Search strategy

This systematic literature review (SLR) was conducted by following the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) method (Page et al., 2021). The SLR is a reliable way to acquire a consistent overview of a specific research theme in an organized and replicable manner (Denyer and Tranfield, 2009; Tranfield et al., 2003). Before engaging in the systematic collection of studies, we conducted a non-structured search to identify regularly associated keywords and concepts about the subject. Keywords such as “COVID-19/SARS-CoV-2”, “wastewater”, “surveillance”, “methods”, “analysis”, “modeling”, and “correlation” were commonly used to identify records of peer-reviewed articles in the multidisciplinary literature. Next, we chose the following databases: ISI Web of Science (www.webofscience.com) and Scopus (www.scopus.com) given their relevance in the academic literature (Wang and Maltman, 2016), along with Engineering Village (Elsevier's Compendex) (www.engineeringvillage.com) due to its importance in the interdisciplinary engineering field (Cusker, 2013), and PubMed (MEDLINE) (www.pubmed.ncbi.nlm.nih.gov) for its reliability as a medical database for evidence-based studies such as systematic reviews (Falagas et al., 2008; Gusenbauer and Haddaway, 2020). As presented above, to address the current panorama of WBE for COVID-19 surveillance to a fuller extent, we considered a conjoint exploration of two pillars: laboratory procedures to quantify viral RNA in wastewater, and modeling (computational) methods to, among other goals, predict outbreaks in the community and city levels. The latter pillar is fed with input generated by the former. To fulfill this

purpose, we performed two independent collections of records from the databases (one for viral RNA analysis and quantification and another for epidemiological modeling), with each one adopting suitable search strings as shown in Table 1. We decided to approach the search in this manner due to the extensive number of studies returned upon conducting a single search for both subjects. Then, we were able to significantly reduce the initial number of records found in the databases and work with a reasonable sample of articles. Table 1 summarizes the keywords used as search strings and how they were combined to compose each search.

2.2 Selection of studies and filtering

The academic coverage and analysis of the COVID-19 pandemic and its consequences demand trustable data to mitigate the risk of misconceptions in matters of public health information and public policies (Davenport et al., 2020; Tagliabue et al., 2020). Accordingly, we favored the side of selecting records from trustworthy sources when pondering the trade-off that exists between considering a high-quality level of discussion and broadening the information basis, with the latter often associated with doubtful reliability (Tranfield et al., 2003). Thus, we decided to include only peer-reviewed, original articles, therefore excluding other types of studies and publication formats, such as reviews, short communications, technical reports, letters, notes, abstracts, and surveys. Any available but unpublished work was excluded as well. Studies published between January 2020 and August 2022 were included in the sample. The deduplication, screening, filtering, and application of inclusion criteria were performed in EndNote 20 to reduce the original sample of studies to a trustful and representative collection of knowledge in the field. The initial search using the terms

expressed in Table 1 returned a total of 2,400 articles, most of them unrelated to our subjects. Next, we engaged in the screening phase as described: collected records were primarily screened for their title only, and subsequently, for their abstracts and content in full. Throughout this phase, we considered ineligible any publication that addressed topics outside our focus, such as other types of viral pathogens in wastewater, COVID-19 diagnosis and treatment, other matrices such as soil, leachate and air, elimination of various pathogens in water, water quality, wastewater from aircraft and ships, drug detection, and biosensors, to name a few. About their content, we included publications that presented (1) clear and concise descriptions and/or comparisons of analytical methods, protocols, and technologies currently used for pre-treatment, concentration, extraction, and quantification of SARS-CoV-2 nucleic acid in a wastewater matrix, and/or (2) precise information on the characteristics of studied wastewater, study location, time range, application of statistical tests for correlating WBE variables, as well as any used or developed mathematical model toward exploring COVID-19 infection dynamics.

Regarding the filtering procedures, we first used EndNote 20 to detect and exclude duplicated records independently for each search, then we applied time range and language filters, followed by the last filter regarding the type of publication from an initial total of 1,106 identified literature records. After this screening step, we assessed the remaining studies through the lens of the established inclusion and exclusion criteria, thus excluding 558 and 84 records by title-only and abstract, respectively; these studies were considered out of the scope of this review, and thus deemed ineligible. Besides, 21 studies were unclear about their methodological procedures or had not

presented any type of wanted information, therefore excluded. Last, we combined the two groups into a single pool and ran a second deduplication, thus excluding another seven studies. Finally, we finished with a list of 96 studies from the first group and 62 studies from the second group as presented in Figure 1, amounting to a final pool of 158 works.

2.3 Data extraction

To properly organize the data extraction process, we used a MS Excel spreadsheet with designated columns to include the following information reported by the selected studies: study location, sample collection period, wastewater characteristics (wastewater treatment plant (WWTP) influent, sewage, or treated), sample pre-treatment, concentration/extraction methods, gene targets, quantification method/technology, initial sample volume processed, lowest and highest viral concentrations recovered in both solid and liquid phases, the estimated time offset (lag) between sample analysis and epidemiological reporting, statistical test to correlate wastewater viral load and clinical data and its result, and mathematical modeling strategy. Not all the studies were thorough in reporting this set of systematized categories, nevertheless, we reasoned that these categories were potentially discussed at some level in our sample of studies, and thus every publication on the selected portfolio should be able to contribute within the scope of this review.

3 RESULTS

3.1 Scientific contribution of the selected studies to SARS-CoV-2 WBE

A meaningful result from the analysis of the reviewed publications was the identification of six main types of contributions in the WBE for COVID-19 surveillance field from 2020 to 2022, which are listed next and depicted in Figure 2: (1) quantitative comparison of concentration, extraction or quantification methods through parametric studies, (2) local reporting of SARS-CoV-2 detection in wastewater and the respective methodology, (3) development, adaptation and/or optimization of analysis protocols, (4) building correlations between viral concentration levels and clinical testing data, (5) mathematical modeling, simulation or parameter estimations for SARS-CoV-2 WBE, and (6) phylogenetics, genotyping and/or identification/quantification of variants of concern (VOC).

3.2 Aspects of wastewater analysis for SARS-CoV-2 detection and quantification

SARS-CoV-2 RNA can be found thermically stable in untreated wastewater at temperature values ranging from 4 to 37°C (Ahmed et al., 2020c). This particularly wide range allows laboratory operations to reliably preserve and detect the virus, even having passed through sample collection and processing. The general methodology to generate a quantified viral concentration, in terms of cycle threshold (Ct) units or genomic concentration, from a wastewater sample, follows a sequential procedure of well-defined steps, namely sampling, pre-treatment, concentration, extraction (or isolation), and detection followed by quantification. This framework is depicted in Figure 3. We found that the operationalization of these steps is well diverse, containing different technologies and protocols that use a wide range of reagents (Kaya et al.,

2022). A complete list of the procedures and technology used can be found in Table 2. Next, we describe the general aspects of each step of the current paradigm of wastewater testing for SARS-CoV-2 WBE.

Pre-treatment. Pre-treatment of the wastewater samples has been ignored as a step of the laboratory analysis process in previous review articles, even though several procedures preceding the concentration step were found in roughly 80% of the studies in our pool. Nevertheless, the term “pre-treatment” was employed only in a few studies (Torii et al., 2021; Zhang et al., 2022). Common pre-treatment procedures involve viral inactivation, pre-centrifugation, pH adjustment, and filtration through a single or a sequence of membrane filters. Pre-treatment serves the purpose of removing coarse solid material (Jmii et al., 2021), separating fine solids, and further purifying against bacterial beings (Reynolds et al., 2022). For the inactivation, we found that it can be performed through thermal treatment (Calderon-Franco et al., 2022; McMinn et al., 2021), UV light (Castiglioni et al., 2022; Pellegrinelli et al., 2022), or chemically (Tomasino et al., 2021). Filtration was done at the micrometer level (maximum pore size of 2 μm), and pre-centrifugation was performed at a minimal value of 1,500g but not exceeding 6,000g for a minimum duration of 5 minutes and a maximum of 45 minutes. Adjustment of sample pH was done when required for the following concentration step by using negatively charged membranes or precipitation using polyethylene glycol (PEG) (Farkas et al., 2021; Hasing et al., 2021).

Concentration. Concentration methods should ideally fulfill some features, including but not limited to being sensitive, reproducible, simple from a technical point of view, economical, rapid, and provide high viral recoveries (Prakash, 2021). A single

standardized method for SARS-CoV-2 concentration from sewage has not been reported (Wehrendt et al., 2021). However, several methods have been described in the literature for that purpose (Prakash, 2021). Following the criteria of Birnbaum et al. (2022), these methods can be classified into two categories: (i) size-based techniques, such as ultrafiltration (Dumke et al., 2021; Hasing et al., 2021), ultracentrifugation (Zheng et al., 2022), centrifugal ultrafiltration (Anderson-Coughlin et al., 2021), and adsorption-elution with electronegative membranes (Barril et al., 2021; Jmii et al., 2021), and (ii) entrapment in chemical precipitates, namely, PEG precipitation (Alexander et al., 2020; Farkas et al., 2021), aluminum flocculation (Pino et al., 2021; Salvo et al., 2021), or skimmed milk flocculation (Philo et al., 2021; Pino et al., 2021). Explaining the fundamentals of these concentration methods is beyond the scope of this review, nevertheless the literature is rich in guidelines for the application of these methods. Readers can refer to the studies of Kaya et al. (2022), Dumke et al. (2021), Barril et al. (2021), and Salvo et al. (2021) to understand in detail these concentration methods and how they have been compared quantitatively. Overall, these studies share the conclusion that PEG precipitation, aluminum flocculation, and ultrafiltration methods favor higher viral recovery rates during the concentration step.

Detection & Quantification. The most frequently used method in WBE for SARS-CoV-2 RNA detection is polymerase chain reaction (PCR)-based quantification (Ni et al., 2021). In this regard, real-time reverse transcription–PCR (real-time RT-PCR) has been employed for identifying SARS-CoV-2 genetic targets (Ni et al., 2021; Thongpradit et al., 2022) and is still considered the gold standard method for the detection of SARS-CoV-2 (Ambrosi et al., 2021). Regarding its genomic targets, the nucleocapsid (N) or

the envelope (E) protein genes, as well as the ORF1ab gene are the most often used RT-PCR targets, as presented in Table 2 (Corman et al., 2020; Kitajima et al., 2020). The Centers for Disease Control and Prevention (CDC) indicates the use of probes targeting several loci (N1 and N2) of the nucleocapsid via separate reactions (CDC, 2020). Particularly, N1 is commonly employed as an indicator for detecting SARS-CoV-2 in wastewater (Navarro et al., 2021). Different PCR procedures form a list that encompasses the reverse transcription loop-mediated isothermal amplification (RT-LAMP) (Amoah et al., 2021), the reverse transcription droplet digital PCR (RT-ddPCR) (Flood et al., 2021), the reverse transcription quantitative PCR (RT-qPCR) (Ahmed et al., 2020a) and its variations and improvements (La Posa et al., 2020; Navarro et al., 2021). As can be concluded from Table 2, the RT-qPCR is the most often employed detection/quantification technology and was used in roughly 87% of the studies in our pool.

3.3 Correlating clinical testing data to viral concentrations in wastewater

The correlation between SARS-CoV-2 viral concentration in sewage water and the number of COVID-19 cases is one of the major challenges of applying viral detection in sewage water to track the scale of SARS-CoV-2 spread in a community (Haque et al., 2021). According to Peccia et al. (2020), some studies have reported the successful correlation of viral RNA levels in wastewater and sludge with the number of reported COVID-19 cases. Such correlation is useful to predict the number of active cases in the population (Ahmed et al., 2020a; Hellmer et al., 2014; Li et al., 2021b; Saththasivam et al., 2021). Particularly, with this information, WBE models can translate

viral concentrations in wastewater to the incidence of SARS-CoV-2 shedders within a community (Cao and Francis, 2021).

There are mainly two statistical-based approaches to evaluate these correlations, the estimation of Pearson's correlation coefficient (Forthofer et al., 2007), which is applied to evaluate the level of linear association between two normally distributed variables, and Spearman's rank correlation coefficient for non-normally distributed data prone to contain outliers (Schober et al., 2018). Both coefficients vary within the range from -1 to +1, where -1 indicates a perfect negative relationship between the variables, 0 indicates the inexistence of a linear relationship and +1 points to a strong positive linear association. We found rather high positive coefficients peaking at values of 0.947 (Galani et al., 2022), 0.95 (D'Aoust et al., 2021), and 0.96 (Layton et al., 2022), for instance, but we also collected moderate (Giraud-Billoud et al., 2021; Tandukar et al., 2022; Tomasino et al., 2021) and weak (Ahmed et al., 2020a) correlations when evaluated through these two statistical approaches depending on the nature of their data. Additionally, some studies in the literature have reported negative correlations (Wehrendt et al., 2021). Recently, a meta-analysis study conducted by Li et al. (2023) collected 133 correlation coefficients ranging from -0.38 to 0.99 for Pearson's or Spearman's coefficients; according to the authors, such a wide range of coefficient values is endorsed by several factors, including variations of the environmental conditions, epidemiological conditions, sampling design, air temperature, etc. This observation is consistent with the numbers found throughout our review process, which also showed a spacious range of values.

To maintain concentration levels meaningful and consistent, normalizing the viral concentration is of paramount importance due to the variability of viral levels in wastewater, which is caused by several factors (Li et al., 2023), such as wastewater flow rate, weather conditions, total suspended solids, and daily fecal discharge. This normalization has been reported in the literature to be addressed via various approaches, such as daily mass flux and/or the use of biomarkers (Qiu et al., 2022). One of these biomarkers, the Pepper Mild Mottle Virus (PMMV), found in human fecal excreta (Rosario et al., 2009), has been used to normalize the SARS-CoV-2 signal (LaTurner et al., 2021; Qiu et al., 2022; Robotto et al., 2022), contributing to obtain strong correlations between the viral concentration level in wastewater and COVID-19 clinical cases (D'Aoust et al., 2021). When addressing the correlation between viral concentration in wastewater and COVID-19 cases it should be noted that viral RNA concentrations in wastewater can be considered a lagging indicator since the virus continues to be shed after the infected individuals have been recovered (McMahan et al., 2021). This lag time has been reported in several studies to range between 2 and 28 days (Zhao et al., 2022), but it lacks a well-accepted definition as discussed in the next paragraphs. Such variation in the lag times can be caused by multiple factors, including but not limited to, daily changes in population size, wastewater sampling methods, responses of the society to the pandemic, and variations in the time required for reporting case data (Medema et al., 2020; Peccia et al., 2020). For instance, some authors have reported that the duration of viral shedding in the stools can be extended up to 33 days after obtaining a negative nasopharyngeal swab (Gupta et al., 2020; Jones et al., 2020). From a symptom onset perspective, it has been suggested that

fecal viral shedding can hold up to more than 20 days (Wolfel et al., 2020; Wu et al., 2020), with Miura et al. (2021) having estimated a value of 26 days. Although different lag time values have been proposed in several WBE studies, these works usually lack a definition for this term, which can be a potential source of confusion when comparing SARS-CoV-2 WBE studies. Zhao et al. (2022) consider the lag time as the temporal gap between the measured SARS-CoV-2 concentration peaks and the reported COVID-19 clinical testing cases peaks, while Omori et al. (2021) define this term as “the lag between the detection timing from wastewater and reporting by passive surveillance”; ideally, detection and reporting timing should be concurrent, however in practice that is not the case, especially in low and middle-income countries (Li et al., 2021d). Finally, lag times may also be influenced by SARS-CoV-2 incubation time and shedding duration (Zhao et al., 2022). For instance, Wu et al. (2022) explained that the lag time they reported (4 days) was consistent with the common incubation period from viral infection to symptom inception, which is considered to be between 4 and 5 days. Other studies reported lag periods similar to this value (Lara-Jacobo et al., 2022; Peccia et al., 2020; Xiao et al., 2022).

3.4 Modeling of WBE for COVID-19 surveillance

Modeling techniques for COVID-19 surveillance in wastewater comprise a rather wide spectrum, going from a plethora of regression techniques to the application of conservation principles and more elegant and contemporary data-driven methods. Comparing or ranking the results from each modeling approach is beyond the scope of this review and itself is a complex task, although several studies have presented comparisons between the performance of different predictive models as in Zhao et al.

(2022), Aberi et al. (2021), and Li et al. (2021b). Table 3 brings a complete description of the methodological approaches reported in our pool of studies for modeling WBE for COVID-19 surveillance.

4 DISCUSSION

4.1 Issues and promising alternatives for SARS-CoV-2 analysis in wastewater

Quantifying low viral loads from non-clinical samples represents one of the major challenges of WBE (Calderon-Franco et al., 2022). When viral particles enter the sewage system, dilution occurs by the addition of other types of water (i.e., domestic sewage and stormwater combined or separated in the sewer), while concomitantly being exposed to a diverse range of chemical agents and physical conditions (Haramoto et al., 2020; Krivonakova et al., 2021). In this regard, a concentration step is required due to the low levels at which SARS-CoV-2 RNA is found in wastewater (Boogaerts et al., 2021; Peinado et al., 2022; Zheng et al., 2022). Several works have reported viral RNA detection in untreated wastewater (influent of the WWTP), being its concentration in the range of 10^2 - 10^5 copies per liter and the maximum exceeding 10^6 copies per liter (Kitajima et al., 2020). PEG precipitation represents a simple and low-cost alternative for viral concentration in wastewater (Flood et al., 2021). In PEG precipitation, the solvent is preferentially trapped, and proteins (e.g., virion) are sterically excluded from the solvent phase by PEG. This way, proteins can be concentrated and precipitated once their concentrations surpass the saturated solubility (Torii et al., 2022). Despite being used in many laboratories, this method suffers from losing approximately half of the viral fragments bound to solid matter (Perez-Cataluna et al.,

2021). On the other hand, skimmed milk flocculation can be considered a promising approach for low-resource areas since extensive laboratory resources are not needed. Additionally, this method does not require consumables that are challenging to acquire, thus enabling the performance of uninterrupted surveillance (Philo et al., 2021). It is worth highlighting that some of these methods were developed for detecting non-enveloped enteric viruses (e.g., norovirus, adenovirus, and enterovirus), which have been the focus of most studies that investigate the existence of viruses in municipal wastewater and human excreta (Ahmed et al., 2020c; Flood et al., 2021). Additionally, SARS-CoV-2 concentration methods have been assessed using surrogate viruses to mimic SARS-CoV-2, since personnel with special training and a laboratory that fulfills Biosafety Level 3 are required for the culture of this virus. Examples of these surrogate viruses include Alphacoronavirus HCoV-229E, bovine respiratory syncytial virus (BRSV), bovine coronavirus BCoV, porcine epidemic diarrhea virus (PEDV), murine hepatitis virus, F-specific RNA phages, avian coronavirus of infectious bronchitis virus, mengovirus or Pseudomonas phage Phi6 (Ahmed et al., 2020c; Aquino de Carvalho et al., 2017; Balboa et al., 2020; Flood et al., 2021; Gendron et al., 2010; Hata et al., 2020; Kocamemi et al., 2020; La Rosa et al., 2020; LaTurner et al., 2021; Medema et al., 2020; Randazzo et al., 2020; Torii et al., 2022).

When it comes to the precise detection and viral quantification, multiplex PCR enables multiple target detection and/or quantification with a sensitivity comparable to that of singleplex PCR. Hence, multiplex PCR is a promising technology since it is more cost-effective and time-saving, reduces the required sample volume, and minimizes the variability due to pipetting. Nevertheless, the complexity of this assay requires

optimization to prevent several undesired phenomena, such as primer-probe sets interaction (Navarro et al., 2021; Xiao et al., 2020b). On a different note, the implementation of RT-qPCR to detect SARS-CoV-2 in wastewater has some limitations, as is time-consuming (it could take 24 h), is highly susceptible to the presence of inhibitors, and sample contamination may occur, resulting in false negative results (Ahmed et al., 2022a). Following this problem, it is necessary to explore new, alternative approaches for the detection of SARS-CoV-2 RNA in wastewater. Apart from PCR-based approaches, other methods can also be employed for viral detection (Lara-Jacobo et al., 2022). For instance, metatranscriptomic sequencing also referred to as Next Generation Sequencing is an alternative for SARS-CoV-2 detection and quantification. This technology allows capturing the whole virus genome, which is of paramount importance due to the incidences of mutation events that increase the virulence, thus significantly improving the sensitivity (Boogaerts et al., 2022; Ni et al., 2021). However, the low SARS-CoV-2 RNA concentration in wastewater, along with the existence of nucleic acids from bacteria, other viruses, animal products, and humans, makes conventional metatranscriptomic sequencing an inappropriate technology for WBE applications at this developmental stage (Boogaerts et al., 2022; Ni et al., 2021). To surmount the low viral RNA concentration challenge, the ARTIC Network amplicon library (Nemudryi et al., 2020) and ATOplex (Xiao et al., 2020b) have been developed. Moreover, RT-LAMP has also been used for SARS-CoV-2 detection (Wei et al., 2021). This method is based on the amplification of the nucleic acids under isothermal conditions, thus avoiding the need for thermal cyclers. Different RT-LAMP approaches can be distinguished, such as the colorimetric or visual RT-LAMP or the fluorescent RT-

LAMP. The former enables the visual reading of the results, whereas the latter facilitates the detection of positive amplification by using a fluorescent dye (Amoah et al., 2021; Huang et al., 2020).

Prominent alternatives in the field may be the use of sensors based on electrochemical principles (Chaibun et al., 2021; Kumar et al., 2022; Ramanujam et al., 2021), which have been extensively studied in terms of specificity and selectivity for different types of nanomaterials. Another plausible approach is the use of magnetic devices where magnetic nanoparticles (MNPs) designed for the separation and detection of the viral pathogens in water samples are employed (Gómez-Pastora et al., 2014; Materón et al., 2021; Yue et al., 2020). These devices might be a promising detection technology as they could be simpler, more accurate, economic, rapid, and portable, allowing the measurements to be performed at the WWTPs by plant technicians.

Finally, it has been reported that viral RNA detection can be influenced by several factors, such as the method used for RNA concentration, or the prevalence of COVID-19 infections in the community (Haramoto et al., 2020; Ni et al., 2021). As an instance of the urgent need for analytical accuracy when performing wastewater processing for WBE, we found a decent number of studies that addressed the comparison of the measured viral recovery amongst distinct concentration and extraction procedures; we also found studies exploring the divergences in the detection and quantification of the viral loads using variations of the PCR approach (Ahmed et al., 2022b; Flood et al., 2021). These problematics elevate the magnitude of the issue. Standardization will be a natural consequence of addressing this problem.

4.2. SARS-CoV-2 epidemiological modeling

The common assumption when using WBE for COVID-19 surveillance is that the number of viral copies observed in the wastewater samples and the reported cases from clinical sources result from the real number of infections, which encapsulates symptomatic and asymptomatic cases (Schmitz et al., 2021; Xiao et al., 2022). Following this premise, wastewater-based epidemiological models have demonstrated to be a valuable tool for estimating the number of infected individuals within a population and identifying COVID-19 infection hotspots. WBE has been hampered by the difficulty of properly correlating viral RNA measurements in wastewater to the number of infections. Also, the real number of infected individuals is generally unknown due to the limitations of the current individual testing capacity systems, especially in low-income areas. To overcome these issues, the number of infected individuals has been linked to the viral concentration (gene copies per volume) and the mass rate of viral RNA in wastewater (gene copies per day) (McMahan et al., 2021). The latter is argued to be preferable over the former because of the serial dilutions of wastewater that might occur due to rainfall, for instance, which alters the viral concentration levels along the sewer network. Using the viral mass rates can be a promising approach when developing epidemiological models given that fluctuations in flow rates compensate for the changes in viral concentrations, leaving viral mass rates unaltered. Regarding the modeling techniques, a wide range of tools has been used as demonstrated in Table 3, which can be categorized into (1) regression techniques (Krivonakova et al., 2021; Peccia et al., 2020; Tomasino et al., 2021), (2) conservation principles (McMahan et al., 2021; Saththasivam et al., 2021), and (3) data-driven methods (Aberi et al., 2021; Li et al.,

2021b; Pereira et al., 2020). Different regression approaches have been explored, to name a few: simple univariate and multivariate linear regression (Kuhn et al., 2022; Roka et al., 2021; Zhao et al., 2022), logistic regression (Scott et al., 2021), Autoregressive Integrated Moving Average (ARIMA) (Karthikeyan et al., 2021; Zhao et al., 2022), and the Vector Autoregression (VAR) model (Cao and Francis, 2021; Zhao et al., 2022). WBE can be applied for the back-calculation of infection prevalence. For that purpose, regression techniques are one of the most important tools in WBE modeling. However, these methods may lead to misleading inferences, since they are proposed for independent data with linear correlations, and the WBE data are time series data (Aberi et al., 2021; Cao and Francis, 2021). In the conservation principles category, an example is the application of the susceptible-exposed-infectious-recovered (SEIR) model, which has shown promising results to predict infection prevalence through a set of interconnected ordinary differential equations (Fernandez-Cassi et al., 2021; McMahan et al., 2021; Nourbakhsh et al., 2022; Proverbio et al., 2022). Furthermore, we found that data-driven methods have acquired considerable popularity given the number of studies that employed these approaches to address the complex task of building epidemiological models, with Artificial Neural Networks (ANN) (Galani et al., 2022; Jiang et al., 2022; Li et al., 2021b; Zhu et al., 2022), Adaptive Neuro-Fuzzy Inference System (ANFIS) (Li et al., 2021b), and the Generalized Additive Model (GAM) method (Aberi et al., 2021; Anneser et al., 2022; Vallejo et al., 2022), as examples of a larger group of techniques listed in Table 3. The common interest in using these approaches may have a root in the fact that epidemiological data are generated in large

amounts with a daily frequency, and that data-driven models must be constantly fed and updated with new inputs for better prediction performance.

Ideally, WBE models should account for the changeability and uncertainty in their variables, specifically for the shedding quantities and secretion routes, such as feces, urine, and sputum (Tiwari et al., 2022). We found that the usual way to associate uncertainty appears to be through the Monte Carlo simulation, which was mostly used to associate uncertainty when estimating the infection prevalence (Amereh et al., 2022; de Sousa et al., 2022; Gonzalez-Reyes et al., 2021; Wang et al., 2021) and the shedding rates (de Freitas Bueno et al., 2022; Wu et al., 2022). Going further into modeling uncertainty, it should be noted that relevant variables are potentially able to create a certain degree of uncertainty. These variables are included as model variables or functions in different WBE models. Examples are: the number of active cases influencing viral counts in wastewater (persons) (Gonzalez-Reyes et al., 2021; Rodriguez Rasero et al., 2022) daily stool mass ($\text{g}_{\text{feces}} \cdot \text{person}^{-1}$) (Ahmed et al., 2020a; Amereh et al., 2022; Claro et al., 2021; Pillay et al., 2021), shedding rate of SARS-CoV-2 RNA (gene copies. $\text{g}_{\text{feces}}^{-1}$ or gene copies. $\text{g}_{\text{feces}}^{-1} \cdot \text{day}^{-1}$) (Ahmed et al., 2020a; Claro et al., 2021; Kuhn et al., 2022; Li et al., 2021c; McMahan et al., 2021; Pillay et al., 2021; Schmitz et al., 2021), decay of SARS-CoV-2 RNA due to storage (time^{-1}) (Kaya et al., 2022; Li et al., 2021c; Yanac et al., 2022) and time-dependent RNA degradation (McMahan et al., 2021), the offset between the observed wastewater viral RNA concentration and the estimated patient viral load (Zhu et al., 2022), and RNA temperature-dependent half-life (h) (Ahmed et al., 2020b; McMahan et al., 2021). On the same note, from a clinical perspective, it is not established the influence of the

severeness of the disease on the magnitude of daily shedding (genome copies per gram of stool), and this constitutes another major source of uncertainty in WBE modeling.

Apart from the aforementioned uncertainty sources, Pillay et al. (2021) reported that the variability of the WBE approach may be mainly caused by changes in the environmental conditions (e.g., the viral dilution and stability in water are influenced by rainfall events and temperature) and the unique features of WWTPs. They highlighted the major importance of accurate knowledge of the shedding pattern within the WWTP catchment. Additionally, these authors explained that the weight of stool that is daily produced per person, which is regionally dependent and may be impacted by several factors, influences the accurate estimation of the number of infected individuals. Furthermore, we found several factors that may influence the accuracy of the back-calculation of the infection prevalence, namely population size, bioindicators' stability (PMMoV), excretion rates, sampling method, and sample preparation. Additionally, several parameters, including the temperature, per-capita water, and average travel time in the sewer, represent critical variables that are needed for identifying infection hotspots when the WBE model is applied.

4.3 Current research gaps and future guidelines for SARS-CoV-2 WBE

Further clarifications on SARS-CoV-2 WBE that need to be addressed in the near future include the persistence of the virus in the wastewater, the effect of the shedding dynamics of the virus in feces, whether urban and rural wastewater systems exhibit significative differences in their characteristics, and how the normalization of viral levels in wastewater with regard to population size should be performed (Fitzgerald et

al., 2021). Additionally, one must bear in mind that COVID-19 is unevenly distributed across population types so considering cross-city differences is of paramount importance. Hence, a 'one size fits all' approach should not be applied to disease surveillance (Kuhn et al., 2022). We suggest that public health information should not be predicted by wastewater analysis alone but by a combination of wastewater-derived information and other data sources. This is due to the fact that changes in factors such as local demographics along with the limitations of current clinical testing/reporting systems may affect the potential of domestic wastewater as a source of information for prediction tools (Xiao et al., 2022). Other factors that should be considered in WBE modeling are reported by Kuhn et al. (2022), including the shedding duration (i.e., how long an infected individual may shed viral particles through feces), and the relationship between the infection severity and the number of viral particles that are shed. At this point, it is not completely understood how these two variables may cause changes in the observed wastewater viral concentration.

We also suggest the utilization of the solid portion of the wastewater as an alternative matrix for the analysis. There is evidence that enveloped viruses feature a high inclination to bind to the surface of solids in wastewater in comparison to non-enveloped viruses (Ye et al., 2016). As it was previously mentioned, SARS-CoV-2 possesses a lipid outer envelope (Klein et al., 2020) whose hydrophobicity may promote greater viral binding to solids in the wastewater, thus affecting viral recovery (Ahmed et al., 2020a; Ahmed et al., 2020c; Anderson-Coughlin et al., 2021). It has been pointed out that the chain of wastewater analysis procedures should not only focus on the supernatant fraction but also on the solid portion of the wastewater (Westhaus et al.,

2021; Yanac et al., 2022). Additionally, and from a WBE perspective, concentration levels from the solid portion have been correlated better with COVID-19 incidence numbers when compared to signals obtained from the liquid part of the wastewater (Tanimoto et al., 2022). For modeling purposes, normalization of the concentration levels from the solid portion of the wastewater can be performed through either total suspended solids measurements (Nourbakhsh et al., 2022) or using the concentration of PMMoV; however, comparability between the concentrations obtained from the solid and liquid phases through PMMoV normalization is still restricted (Kim et al., 2022). In this regard, different studies have reported the prevalence of viral particles in the solid phase obtained from domestic wastewater, as well as observations pointing to a significantly higher amount of viral RNA in the solid portion (Kim et al., 2022; Kumblathan et al., 2023). For instance, Kitamura et al. (2021) reported that a higher level of SARS-CoV-2 RNA, compared to PMMoV RNA, was contained in the solid fraction, whereas supernatant fractions comprised lower SARS-CoV-2 RNA levels. They reasoned that the different detection of SARS-CoV-2 RNA and PMMoV RNA in the solid and liquid fractions could result from the fact that PMMoV lacks an envelope, which is present in SARS-CoV-2. Similarly, Li et al. (2021a) found that SARS-CoV-2 RNA was considerably more abundant in the solid than in the liquid fraction. This observation was further endorsed by the studies of Ni et al. (2021) and Tomasino et al. (2021) in terms of viral recovery. The higher viral RNA concentration in the solid phase of wastewater, along with the more time-efficient processing of the solid fraction (Li et al., 2021a; Nourbakhsh et al., 2022), led us to suggest that wastewater solids may represent a

more convenient sample matrix, thus being a promising approach to improve analytical accuracy in WBE for SARS-CoV-2.

As highlighted throughout the present study, WBE represents a valuable tool for predicting COVID-19 cases. To this end, WBE can be implemented via several statistical models with data gathered from wastewater (Ando et al., 2023; Anneser et al., 2022). However, establishing and standardizing protocols are still required so that worldwide conducted studies could be successfully compared (Amereh et al., 2022; Fitzgerald et al., 2021). In further words, the current lack of standardization is revealed by the wide range of sample initial volumes and concentration methods that have been reported by the different studies. Thereby, sample initial volumes range from 2 mL to 1 L; moreover, the extensive variety of concentration methods that have been used include size-based and entrapment in chemical precipitates techniques, such as conventional filtration, ultrafiltration, ultracentrifugation, centrifugation, filtration using negatively charged membranes, precipitation, and direct extraction, as well as their combinations. Furthermore, it is still unclear how different pre-treatment techniques may affect the detection performance through PCR methods. In this sense, the recognition of the pre-treatment step in the wastewater analysis process is crucial to further develop standard protocols for SARS-CoV-2 detection and quantification. Finally, quality controls, variable testing, and the optimization of the methodology are considerably lacking; however, they are required in order to provide analytical accuracy (Calderon-Franco et al., 2022).

5 CONCLUSIONS

SARS-CoV-2 will remain a constant threat to public health given the increasing infectibility of new VOCs. In this study, we reviewed the recent WBE research endeavor to mitigate the hefty burden of COVID-19 on the health systems around the globe. More specifically, this review collects and organizes the recent progress on the analytical methods reported between 2020 and 2022 to detect and quantify SARS-CoV-2 RNA from wastewater samples. We also review the methods by which SARS-CoV-2 wastewater-based epidemiological modeling has been approached to use the output of lab analysis for diverse purposes, such as predicting outbreaks in a community, estimation of active human shedders (or infected individuals), and shedding rates, to name a few. Correlating the amount of genomic material in wastewater with the number of COVID-19 cases within a community is a component of epidemiological modeling that has been tried through a wide range of mathematical methods, with data-driven models considered the most popular approach to address predictions of variables correlated to outbreaks within a certain time horizon, based on genomic viral material measurements in domestic wastewaters. We also highlight the promising opportunities to improve the accuracy and rapidness of viral detection using the solid portion of wastewater as an alternative testing matrix, and the design of novel sensors based on electrochemical or magnetic devices. However, as evidenced throughout this work, recent research has not focused on ways to standardize the analytical procedures for comparability between different locations. Implementing the WBE surveillance as a prediction tool for outbreaks and infection waves, which in turn would result in the mitigation of the COVID-19 burden, remains challenging. To promote the worldwide

applicability of WBE surveillance, this lack of standardization should be managed along with the establishment of a testing framework that accounts for the different analytical sensitivities throughout the different steps of the analysis. Notably, this study contributes to future research as a reference guide for what has been proposed and worked so far to understand the dynamics of viral concentrations in wastewater. Since the effort to mitigate the effects of COVID-19 is a global one, future research must expand the scope of this review and consider the needs of low-income countries, whose health systems are often restricted and the implementation of the WBE surveillance strategy can thus become more arduous.

ACKNOWLEDGEMENTS

Financial support from Texas Tech University is acknowledged. Dr. Cristina González-Fernández thanks the Spanish Ministry of Universities for the Margarita Salas postdoctoral fellowship (complementary call of grants for the qualification of the Spanish university system for 2021-2023, University of Cantabria), funded by the European Union – Next Generation EU.

REFERENCES

- Aberi P, Arabzadeh R, Insam H, Markt R, Mayr M, Kreuzinger N, et al. Quest for Optimal Regression Models in SARS-CoV-2 Wastewater Based Epidemiology. *Int J Environ Res Public Health* 2021; 18.
- Ahmed F, Islam MA, Kumar M, Hossain M, Bhattacharya P, Islam MT, et al. First detection of SARS-CoV-2 genetic material in the vicinity of COVID-19 isolation Centre in Bangladesh: Variation along the sewer network. *Sci Total Environ* 2021; 776: 145724.
- Ahmed W, Angel N, Edson J, Bibby K, Bivins A, O'Brien LW, et al. First confirmed detection of SARS-CoV-2 in untreated wastewater in Australia: A proof of concept for the wastewater surveillance of COVID-19 in the community. *Sci Total Environ* 2020a; 728: 138764.
- Ahmed W, Bertsch PM, Bibby K, Haramoto E, Hewitt J, Huygens F, et al. Decay of SARS-CoV-2 and surrogate murine hepatitis virus RNA in untreated wastewater to inform application in wastewater-based epidemiology. *Environ Res* 2020b; 191: 110092.
- Ahmed W, Bertsch PM, Bivins A, Bibby K, Farkas K, Gathercole A, et al. Comparison of virus concentration methods for the RT-qPCR-based recovery of murine hepatitis virus, a surrogate for SARS-CoV-2 from untreated wastewater. *Sci Total Environ* 2020c; 739: 139960.
- Ahmed W, Simpson SL, Bertsch PM, Bibby K, Bivins A, Blackall LL, et al. Minimizing errors in RT-PCR detection and quantification of SARS-CoV-2 RNA for wastewater surveillance. *Sci Total Environ* 2022a; 805: 149877.
- Ahmed W, Smith WJM, Metcalfe S, Jackson G, Choi PM, Morrison M, et al. Comparison of RT-qPCR and RT-dPCR Platforms for the Trace Detection of SARS-CoV-2 RNA in Wastewater. *ACS ES T Water* 2022b; 2: 1871-1880.
- Alexander MR, Rootes CL, van Vuren PJ, Stewart CR. Concentration of infectious SARS-CoV-2 by polyethylene glycol precipitation. *J Virol Methods* 2020; 286: 113977.

- Ambrosi C, Prezioso C, Checconi P, Scribano D, Sarshar M, Capannari M, et al. SARS-CoV-2: Comparative analysis of different RNA extraction methods. *J Virol Methods* 2021; 287: 114008.
- Amereh F, Jahangiri-Rad M, Mohseni-Bandpei A, Mohebbi SR, Asadzadeh-Aghdaei H, Dabiri H, et al. Association of SARS-CoV-2 presence in sewage with public adherence to precautionary measures and reported COVID-19 prevalence in Tehran. *Sci Total Environ* 2022; 812: 152597.
- Amoah ID, Mthethwa NP, Pillay L, Deepnarain N, Pillay K, Awolusi OO, et al. RT-LAMP: A Cheaper, Simpler and Faster Alternative for the Detection of SARS-CoV-2 in Wastewater. *Food Environ Virol* 2021; 13: 447-456.
- Anderson-Coughlin BL, Shearer AEH, Omar AN, Womack KE, Kniel KE. Recovery of SARS-CoV-2 from Wastewater Using Centrifugal Ultrafiltration. *Methods Protoc* 2021; 4: 9.
- Ando H, Murakami M, Ahmed W, Iwamoto R, Okabe S, Kitajima M. Wastewater-based prediction of COVID-19 cases using a highly sensitive SARS-CoV-2 RNA detection method combined with mathematical modeling. *Environ Int* 2023; 173: 107743.
- Anneser E, Riseberg E, Brooks YM, Corlin L, Stringer C. Modeling the relationship between SARS-CoV-2 RNA in wastewater or sludge and COVID-19 cases in three New England regions. *J Water Health* 2022; 20: 816-828.
- Aquino de Carvalho M, Stachler EN, Cimabue N, Bibby K. Evaluation of Phi6 Persistence and Suitability as an Enveloped Virus Surrogate. *Environ Sci Technol* 2017; 51: 8692-8700.
- Balboa S, Mauricio-Iglesias M, Rodriguez S, Martínez-Lamas L, Vasallo FJ, Regueiro B, et al. The fate of SARS-CoV-2 in WWTPs points out the sludge line as a suitable spot for monitoring. *medRxiv* 2020: 2020.05.25.20112706.
- Bar-Or I, Yaniv K, Shagan M, Ozer E, Weil M, Indenbaum V, et al. Regressing SARS-CoV-2 Sewage Measurements Onto COVID-19 Burden in the Population: A Proof-of-Concept for Quantitative Environmental Surveillance. *Front Public Health* 2021; 9: 561710.

- Barbosa MRF, Garcia SC, Bruni AC, Machado FS, de Oliveira RX, Dropa M, et al. One-year surveillance of SARS-CoV-2 in wastewater from vulnerable urban communities in metropolitan Sao Paulo, Brazil. *J Water Health* 2022; 20: 471-490.
- Barril PA, Pianciola LA, Mazzeo M, Ousset MJ, Jaureguiberry MV, Alessandrello M, et al. Evaluation of viral concentration methods for SARS-CoV-2 recovery from wastewaters. *Sci Total Environ* 2021; 756: 144105.
- Birnbaum DP, Vilardi KJ, Anderson CL, Pinto AJ, Joshi NS. Simple Affinity-Based Method for Concentrating Viruses from Wastewater Using Engineered Curli Fibers. *ACS ES T Water* 2022; 2: 1836-1843.
- Boogaerts T, Jacobs L, De Roeck N, Van den Bogaert S, Aertgeerts B, Lahousse L, et al. An alternative approach for bioanalytical assay optimization for wastewater-based epidemiology of SARS-CoV-2. *Sci Total Environ* 2021; 789: 148043.
- Boogaerts T, Van den Bogaert S, Van Poelbroeck LAE, El Masri D, De Roeck N, Roosens NHC, et al. Optimization and Application of a Multiplex Digital PCR Assay for the Detection of SARS-CoV-2 Variants of Concern in Belgian Influent Wastewater. *Viruses* 2022; 14: 17.
- Buscarini E, Manfredi G, Brambilla G, Menozzi F, Londoni C, Alicante S, et al. GI symptoms as early signs of COVID-19 in hospitalised Italian patients. *Gut* 2020; 69: 1547-1548.
- Calderon-Franco D, Orsini L, Lackner S, Agrawal S, Weissbrodt DG. Monitoring SARS-CoV-2 in sewage: Toward sentinels with analytical accuracy. *Sci Total Environ* 2022; 804: 150244.
- Cao Y, Francis R. On forecasting the community-level COVID-19 cases from the concentration of SARS-CoV-2 in wastewater. *Sci Total Environ* 2021; 786: 147451.
- Carducci A, Federigi I, Liu D, Thompson JR, Verani M. Making Waves: Coronavirus detection, presence and persistence in the water environment: State of the art and knowledge needs for public health. *Water Res* 2020; 179: 115907.

- Castiglioni S, Schiarea S, Pellegrinelli L, Primache V, Galli C, Bubba L, et al. SARS-CoV-2 RNA in urban wastewater samples to monitor the COVID-19 pandemic in Lombardy, Italy (March-June 2020). *Sci Total Environ* 2022; 806: 150816.
- CDC. CDC 2019-Novel Coronavirus (2019-nCoV) Real-Time RT-PCR Diagnostic Panel Catalog # 2019-nCoV-EUA-01 1000 reactions. CDC, Atlanta GA, 2020.
- Chaibun T, Puenpa J, Ngamdee T, Boonapatcharoen N, Athamanolap P, O'Mullane AP, et al. Rapid electrochemical detection of coronavirus SARS-CoV-2. *Nat Commun* 2021; 12: 802.
- Cheung KS, Hung IFN, Chan PPY, Lung KC, Tso E, Liu R, et al. Gastrointestinal Manifestations of SARS-CoV-2 Infection and Virus Load in Fecal Samples From a Hong Kong Cohort: Systematic Review and Meta-analysis. *Gastroenterology* 2020; 159: 81-95.
- Claro ICM, Cabral AD, Augusto MR, Duran AFA, Caciola MCP, Fonseca FLA, et al. Long-term monitoring of SARS-COV-2 RNA in wastewater in Brazil: A more responsive and economical approach. *Water Res* 2021; 203: 117534.
- Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* 2020; 25: 2000045.
- Cusker J. Elsevier Compendex and Google Scholar: A Quantitative Comparison of Two Resources for Engineering Research and an Update to Prior Comparisons. *Journal of Academic Librarianship* 2013; 39: 241-243.
- D'Aoust PM, Mercier E, Montpetit D, Jia JJ, Alexandrov I, Neault N, et al. Quantitative analysis of SARS-CoV-2 RNA from wastewater solids in communities with low COVID-19 incidence and prevalence. *Water Res* 2021; 188: 116560.
- Davenport TH, Godfrey AB, Redman TC. To Fight Pandemics, We Need Better Data. *MIT Sloan Management Review* 2020; 62: 1-4.
- de Freitas Bueno R, Claro ICM, Augusto MR, Duran AFA, Camillo LMB, Cabral AD, et al. Wastewater-based epidemiology: A Brazilian SARS-COV-2 surveillance experience. *J Environ Chem Eng* 2022; 10: 108298.
- de Sousa ARV, do Carmo Silva L, de Curcio JS, da Silva HD, Eduardo Anunciacao C, Maria Salem Izacc S, et al. "pySewage": a hybrid approach to predict the number

- of SARS-CoV-2-infected people from wastewater in Brazil. *Environ Sci Pollut Res Int* 2022; 29: 67260-67269.
- Denyer D, Tranfield D. Producing a systematic review. *The Sage handbook of organizational research methods*. Sage Publications Ltd, Thousand Oaks, CA, 2009, pp. 671-689.
- Ding S, Liang TJ. Is SARS-CoV-2 Also an Enteric Pathogen With Potential Fecal-Oral Transmission? A COVID-19 Virological and Clinical Review. *Gastroenterology* 2020; 159: 53-61.
- Dumke R, de la Cruz Barron M, Oertel R, Helm B, Kallies R, Berendonk TU, et al. Evaluation of Two Methods to Concentrate SARS-CoV-2 from Untreated Wastewater. *Pathogens* 2021; 10: 1-7.
- Falagas ME, Pitsouni EI, Malietzis GA, Pappas G. Comparison of PubMed, Scopus, Web of Science, and Google Scholar: strengths and weaknesses. *FASEB J* 2008; 22: 338-42.
- Farkas K, Hillary LS, Thorpe J, Walker DI, Lowther JA, McDonald JE, et al. Concentration and Quantification of SARS-CoV-2 RNA in Wastewater Using Polyethylene Glycol-Based Concentration and qRT-PCR. *Methods Protoc* 2021; 4: 9.
- Fernandez-Cassi X, Scheidegger A, Banziger C, Cariti F, Tunas Corzon A, Ganesanandamoorthy P, et al. Wastewater monitoring outperforms case numbers as a tool to track COVID-19 incidence dynamics when test positivity rates are high. *Water Res* 2021; 200: 117252.
- Fitzgerald SF, Rossi C, Low AS, McAteer SP, O'Keefe B, Findlay D, et al. Site Specific Relationships between COVID-19 Cases and SARS-CoV-2 Viral Load in Wastewater Treatment Plant Influent. *Environ Sci Technol* 2021; 55: 15276-15286.
- Flood MT, D'Souza N, Rose JB, Aw TG. Methods Evaluation for Rapid Concentration and Quantification of SARS-CoV-2 in Raw Wastewater Using Droplet Digital and Quantitative RT-PCR. *Food Environ Virol* 2021; 13: 303-315.

- Fongaro G, Stoco PH, Souza DSM, Grisard EC, Magri ME, Rogovski P, et al. The presence of SARS-CoV-2 RNA in human sewage in Santa Catarina, Brazil, November 2019. *Sci Total Environ* 2021; 778: 146198.
- Forthofer RN, Lee ES, Hernandez M. *Biostatistics*: Elsevier, 2007.
- Galani A, Aalizadeh R, Kostakis M, Markou A, Alygizakis N, Lytras T, et al. SARS-CoV-2 wastewater surveillance data can predict hospitalizations and ICU admissions. *Sci Total Environ* 2022; 804: 150151.
- Gendron L, Verreault D, Veillette M, Moineau S, Duchaine C. Evaluation of Filters for the Sampling and Quantification of RNA Phage Aerosols. *Aerosol Science and Technology* 2010; 44: 893-901.
- Giraud-Billoud M, Cuervo P, Altamirano JC, Pizarro M, Aranibar JN, Catapano A, et al. Monitoring of SARS-CoV-2 RNA in wastewater as an epidemiological surveillance tool in Mendoza, Argentina. *Sci Total Environ* 2021; 796: 148887.
- Gómez-Pastora J, Bringas E, Ortiz I. Recent progress and future challenges on the use of high performance magnetic nano adsorbents in environmental applications. *Chemical Engineering Journal* 2014; 256: 187-204.
- Gonzalez-Reyes JR, Hernandez-Flores ML, Paredes-Zarco JE, Tellez-Jurado A, Fayad-Meneses O, Carranza-Ramirez L. Detection of SARS-CoV-2 in Wastewater Northeast of Mexico City: Strategy for Monitoring and Prevalence of COVID-19. *Int J Environ Res Public Health* 2021; 18: 14.
- Gupta S, Parker J, Smits S, Underwood J, Dolwani S. Persistent viral shedding of SARS-CoV-2 in faeces - a rapid review. *Colorectal Dis* 2020; 22: 611-620.
- Gusenbauer M, Haddaway NR. Which academic search systems are suitable for systematic reviews or meta-analyses? Evaluating retrieval qualities of Google Scholar, PubMed, and 26 other resources. *Res Synth Methods* 2020; 11: 181-217.
- Haque MFU, Bukhari SS, Ejaz R, Zaman FU, Sreejith KR, Rashid N, et al. A novel RdRp-based colorimetric RT-LAMP assay for rapid and sensitive detection of SARS-CoV-2 in clinical and sewage samples from Pakistan. *Virus Res* 2021; 302: 198484.

- Haramoto E, Malla B, Thakali O, Kitajima M. First environmental surveillance for the presence of SARS-CoV-2 RNA in wastewater and river water in Japan. *Sci Total Environ* 2020; 737: 140405.
- Hasing M, Yu JA, Qiu YY, Maal-Bared R, Bhavanam S, Lee B, et al. Comparison of Detecting and Quantitating SARS-CoV-2 in Wastewater Using Moderate-Speed Centrifuged Solids versus an Ultrafiltration Method. *Water* 2021; 13: 16.
- Hata A, Honda R, Hara-Yamamura H, Meuchi Y. Detection of SARS-CoV-2 in wastewater in Japan by multiple molecular assays-implication for wastewater-based epidemiology (WBE). *medRxiv* 2020: 2020.06.03.20126417.
- He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med* 2020; 26: 672-675.
- Hellmer M, Paxeus N, Magnus L, Enache L, Arnholm B, Johansson A, et al. Detection of pathogenic viruses in sewage provided early warnings of hepatitis A virus and norovirus outbreaks. *Appl Environ Microbiol* 2014; 80: 6771-81.
- Huang WE, Lim B, Hsu CC, Xiong D, Wu W, Yu Y, et al. RT-LAMP for rapid diagnosis of coronavirus SARS-CoV-2. *Microb Biotechnol* 2020; 13: 950-961.
- Jiang G, Wu J, Weidhaas J, Li X, Chen Y, Mueller J, et al. Artificial neural network-based estimation of COVID-19 case numbers and effective reproduction rate using wastewater-based epidemiology. *Water Res* 2022; 218: 118451.
- Jmii H, Gharbi-Khelifi H, Assaoudi R, Aouni M. Detection of SARS-CoV-2 in the sewerage system in Tunisia: a promising tool to confront COVID-19 pandemic. *Future Virol* 2021; 16: 751-759.
- Jones-Lepp CGDTL. Pharmaceuticals and Personal Care Products in the Environment: Scientific and Regulatory Issues: American Chemical Society, 2001.
- Jones DL, Baluja MQ, Graham DW, Corbishley A, McDonald JE, Malham SK, et al. Shedding of SARS-CoV-2 in feces and urine and its potential role in person-to-person transmission and the environment-based spread of COVID-19. *Sci Total Environ* 2020; 749: 141364.
- Kabdasli I, Tunay O. Concentration techniques tailored for the detection of SARS-CoV-2 genetic material in domestic wastewater and treatment plant sludge: A review. *J Environ Chem Eng* 2021; 9: 106296.

- Karthikeyan S, Ronquillo N, Belda-Ferre P, Alvarado D, Javidi T, Longhurst CA, et al. High-Throughput Wastewater SARS-CoV-2 Detection Enables Forecasting of Community Infection Dynamics in San Diego County. *mSystems* 2021; 6.
- Kaya D, Niemeier D, Ahmed W, Kjellerup BV. Evaluation of multiple analytical methods for SARS-CoV-2 surveillance in wastewater samples. *Sci Total Environ* 2022; 808: 152033.
- Kim S, Kennedy LC, Wolfe MK, Criddle CS, Duong DH, Topol A, et al. SARS-CoV-2 RNA is enriched by orders of magnitude in primary settled solids relative to liquid wastewater at publicly owned treatment works. *Environ. Sci (Camb)* 2022; 8: 757-770.
- Kitajima M, Ahmed W, Bibby K, Carducci A, Gerba CP, Hamilton KA, et al. SARS-CoV-2 in wastewater: State of the knowledge and research needs. *Sci Total Environ* 2020; 739: 139076.
- Kitamura K, Sadamasu K, Muramatsu M, Yoshida H. Efficient detection of SARS-CoV-2 RNA in the solid fraction of wastewater. *Sci Total Environ* 2021; 763: 144587.
- Klein S, Cortese M, Winter SL, Wachsmuth-Melm M, Neufeldt CJ, Cerikan B, et al. SARS-CoV-2 structure and replication characterized by in situ cryo-electron tomography. *Nat Commun* 2020; 11: 5885.
- Kocamemi BA, Kurt H, Sait A, Sarac F, Saatci AM, Pakdemirli B. SARS-CoV-2 Detection in Istanbul Wastewater Treatment Plant Sludges. *medRxiv* 2020: 2020.05.12.20090352.
- Krivosakova N, Solysova A, Tamas M, Takac Z, Krahulec J, Ficek A, et al. Mathematical modeling based on RT-qPCR analysis of SARS-CoV-2 in wastewater as a tool for epidemiology. *Sci Rep* 2021; 11: 19456.
- Kuhn KG, Jarshaw J, Jeffries E, Adesigbin K, Maytubby P, Dundas N, et al. Predicting COVID-19 cases in diverse population groups using SARS-CoV-2 wastewater monitoring across Oklahoma City. *Sci Total Environ* 2022; 812: 151431.
- Kumar N, Shetti NP, Jagannath S, Aminabhavi TM. Electrochemical sensors for the detection of SARS-CoV-2 virus. *Chem Eng J* 2022; 430: 132966.

- Kumblathan T, Liu YM, Qiu YY, Pang LLY, Hrudehy SE, Le XC, et al. An efficient method to enhance recovery and detection of SARS-CoV-2 RNA in wastewater. *Journal of Environmental Sciences* 2023; 130: 139-148.
- La Rosa G, Iaconelli M, Mancini P, Bonanno Ferraro G, Veneri C, Bonadonna L, et al. First detection of SARS-CoV-2 in untreated wastewaters in Italy. *Sci Total Environ* 2020; 736: 139652.
- Lamers MM, Beumer J, van der Vaart J, Knoop K, Puschhof J, Breugem TI, et al. SARS-CoV-2 productively infects human gut enterocytes. *Science* 2020; 369: 50-54.
- Lara-Jacobo LR, Islam G, Desaulniers JP, Kirkwood AE, Simmons DBD. Detection of SARS-CoV-2 Proteins in Wastewater Samples by Mass Spectrometry. *Environ Sci Technol* 2022; 56: 5062-5070.
- LaTurner ZW, Zong DM, Kalvapalle P, Gamas KR, Terwilliger A, Crosby T, et al. Evaluating recovery, cost, and throughput of different concentration methods for SARS-CoV-2 wastewater-based epidemiology. *Water Res* 2021; 197: 117043.
- Layton BA, Kaya D, Kelly C, Williamson KJ, Alegre D, Bachhuber SM, et al. Evaluation of a Wastewater-Based Epidemiological Approach to Estimate the Prevalence of SARS-CoV-2 Infections and the Detection of Viral Variants in Disparate Oregon Communities at City and Neighborhood Scales. *Environ Health Perspect* 2022; 130: 67010.
- Lescure FX, Bouadma L, Nguyen D, Parisey M, Wicky PH, Behillil S, et al. Clinical and virological data of the first cases of COVID-19 in Europe: a case series. *Lancet Infect Dis* 2020; 20: 697-706.
- Li B, Di DYW, Saingam P, Jeon MK, Yan T. Fine-Scale Temporal Dynamics of SARS-CoV-2 RNA Abundance in Wastewater during A COVID-19 Lockdown. *Water Res* 2021a; 197: 117093.
- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med* 2020; 382: 1199-1207.

- Li X, Kulandaivelu J, Zhang S, Shi J, Sivakumar M, Mueller J, et al. Data-driven estimation of COVID-19 community prevalence through wastewater-based epidemiology. *Sci Total Environ* 2021b; 789: 147947.
- Li X, Zhang S, Sherchan S, Orive G, Lertxundi U, Haramoto E, et al. Correlation between SARS-CoV-2 RNA concentration in wastewater and COVID-19 cases in community: A systematic review and meta-analysis. *J Hazard Mater* 2023; 441: 129848.
- Li X, Zhang S, Shi J, Luby SP, Jiang G. Uncertainties in estimating SARS-CoV-2 prevalence by wastewater-based epidemiology. *Chem Eng J* 2021c; 415: 129039.
- Li Z, Jones C, Ejigu GS, George N, Geller AL, Chang CC, et al. Countries with delayed COVID-19 introduction - characteristics, drivers, gaps, and opportunities. *Global Health* 2021d; 17: 28.
- Markt R, Endler L, Amman F, Schedl A, Penz T, Buchel-Marxer M, et al. Detection and abundance of SARS-CoV-2 in wastewater in Liechtenstein, and the estimation of prevalence and impact of the B.1.1.7 variant. *J Water Health* 2022; 20: 114-125.
- Materón EM, Miyazaki CM, Carr C, Joshi N, Picciani PHS, Dalmaschio CJ, et al. Magnetic nanoparticles in biomedical applications: A review. *Applied Surface Science Advances* 2021; 6: 100163.
- McMahan CS, Self S, Renner L, Kalbaugh C, Kriebel D, Graves D, et al. COVID-19 wastewater epidemiology: a model to estimate infected populations. *Lancet Planet Health* 2021; 5: e874-e881.
- McMinn BR, Korajkic A, Kelleher J, Herrmann MP, Pemberton AC, Ahmed W, et al. Development of a large volume concentration method for recovery of coronavirus from wastewater. *Sci Total Environ* 2021; 774: 145727.
- Medema G, Heijnen L, Elsinga G, Italiaander R, Brouwer A. Presence of SARS-Coronavirus-2 RNA in Sewage and Correlation with Reported COVID-19 Prevalence in the Early Stage of the Epidemic in The Netherlands. *Environmental Science & Technology Letters* 2020; 7: 511-516.
- Mesoraca A, Margiotti K, Viola A, Cima A, Sparacino D, Giorlandino C. Evaluation of SARS-CoV-2 viral RNA in fecal samples. *Virol J* 2020; 17: 86.

- Miura F, Kitajima M, Omori R. Duration of SARS-CoV-2 viral shedding in faeces as a parameter for wastewater-based epidemiology: Re-analysis of patient data using a shedding dynamics model. *Sci Total Environ* 2021; 769: 144549.
- Nasseri S, Yavarian J, Baghani AN, Azad TM, Nejati A, Nabizadeh R, et al. The presence of SARS-CoV-2 in raw and treated wastewater in 3 cities of Iran: Tehran, Qom and Anzali during coronavirus disease 2019 (COVID-19) outbreak. *J Environ Health Sci Eng* 2021; 19: 573-584.
- Navarro A, Gomez L, Sanseverino I, Niegowska M, Roka E, Pedraccini R, et al. SARS-CoV-2 detection in wastewater using multiplex quantitative PCR. *Sci Total Environ* 2021; 797: 148890.
- Nemudryi A, Nemudraia A, Wiegand T, Surya K, Buyukyoruk M, Cicha C, et al. Temporal Detection and Phylogenetic Assessment of SARS-CoV-2 in Municipal Wastewater. *Cell Rep Med* 2020; 1: 100098.
- Ni GF, Lu J, Maulani N, Tian W, Yang L, Harl wong I, et al. Novel Multiplexed Amplicon-Based Sequencing to Quantify SARS-CoV-2 RNA from Wastewater. *Environmental Science & Technology Letters* 2021; 8: 683-690.
- Nourbakhsh S, Fazil A, Li M, Mangat CS, Peterson SW, Daigle J, et al. A wastewater-based epidemic model for SARS-CoV-2 with application to three Canadian cities. *Epidemics* 2022; 39: 100560.
- Omori R, Miura F, Kitajima M. Age-dependent association between SARS-CoV-2 cases reported by passive surveillance and viral load in wastewater. *Sci Total Environ* 2021; 792: 148442.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Syst Rev* 2021; 10: 89.
- Peccia J, Zulli A, Brackney DE, Grubaugh ND, Kaplan EH, Casanovas-Massana A, et al. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nat Biotechnol* 2020; 38: 1164-1167.
- Peinado B, Martinez-Garcia L, Martinez F, Nozal L, Sanchez MB. Improved methods for the detection and quantification of SARS-CoV-2 RNA in wastewater. *Sci Rep* 2022; 12: 7201.

- Pellegrinelli L, Castiglioni S, Cocuzza C, Bertasi B, Primache V, Schiarea S, et al. Evaluation of Pre-Analytical and Analytical Methods for Detecting SARS-CoV-2 in Municipal Wastewater Samples in Northern Italy. *Water* 2022; 14: 12.
- Peng L, Liu J, Xu W, Luo Q, Chen D, Lei Z, et al. SARS-CoV-2 can be detected in urine, blood, anal swabs, and oropharyngeal swabs specimens. *J Med Virol* 2020; 92: 1676-1680.
- Pereira IG, Guerin JM, Silva Júnior AG, Garcia GS, Piscitelli P, Miani A, et al. Forecasting Covid-19 Dynamics in Brazil: A Data Driven Approach. *International Journal of Environmental Research and Public Health*. 17, 2020.
- Perez-Cataluna A, Cuevas-Ferrando E, Randazzo W, Falco J, Allende A, Sanchez G. Comparing analytical methods to detect SARS CoV-2 in wastewater. *Sci Total Environ* 2021; 758: 143870.
- Philo SE, Keim EK, Swanstrom R, Ong AQW, Burnor EA, Kossik AL, et al. A comparison of SARS-CoV-2 wastewater concentration methods for environmental surveillance. *Sci Total Environ* 2021; 760: 144215.
- Pillay L, Amoah ID, Deepnarain N, Pillay K, Awolusi OO, Kumari S, et al. Monitoring changes in COVID-19 infection using wastewater-based epidemiology: A South African perspective. *Sci Total Environ* 2021; 786: 147273.
- Pino NJ, Rodriguez DC, Cano LC, Rodriguez A. Detection of SARS-CoV-2 in wastewater is influenced by sampling time, concentration method, and target analyzed. *J Water Health* 2021; 19: 775-784.
- Prakash C. Sewage Analysis as a Tool for Environmental Surveillance of SARS-CoV-2: Experience from Delhi, India. *Journal of Communicable Diseases* 2021; 53: 1-13.
- Prevost B, Lucas FS, Goncalves A, Richard F, Moulin L, Wurtzer S. Large scale survey of enteric viruses in river and waste water underlines the health status of the local population. *Environ Int* 2015; 79: 42-50.
- Proverbio D, Kemp F, Magni S, Ogorzaly L, Cauchie HM, Goncalves J, et al. Model-based assessment of COVID-19 epidemic dynamics by wastewater analysis. *Sci Total Environ* 2022; 827: 154235.

- Qiu Y, Yu J, Pabbaraju K, Lee BE, Gao T, Ashbolt NJ, et al. Validating and optimizing the method for molecular detection and quantification of SARS-CoV-2 in wastewater. *Sci Total Environ* 2022; 812: 151434.
- Ramanujam A, Almodovar S, Botte GG. Ultra-Fast Electrochemical Sensor for Point-of-Care COVID-19 Diagnosis Using Non-Invasive Saliva Sampling. *Processes*. 9, 2021.
- Randazzo W, Truchado P, Cuevas-Ferrando E, Simon P, Allende A, Sanchez G. SARS-CoV-2 RNA in wastewater anticipated COVID-19 occurrence in a low prevalence area. *Water Res* 2020; 181: 115942.
- Reynolds LJ, Gonzalez G, Sala-Comorera L, Martin NA, Byrne A, Fennema S, et al. SARS-CoV-2 variant trends in Ireland: Wastewater-based epidemiology and clinical surveillance. *Sci Total Environ* 2022; 838: 155828.
- Rimoldi SG, Stefani F, Gigantiello A, Polesello S, Comandatore F, Mileto D, et al. Presence and infectivity of SARS-CoV-2 virus in wastewaters and rivers. *Sci Total Environ* 2020; 744: 140911.
- Robotto A, Lembo D, Quaglino P, Brizio E, Polato D, Civra A, et al. Wastewater-based SARS-CoV-2 environmental monitoring for Piedmont, Italy. *Environ Res* 2022; 203: 111901.
- Rodriguez Rasero FJ, Moya Romano LA, Rasero Del Real P, Cuberos Gomez L, Lorusso N. Associations between SARS-CoV-2 RNA concentrations in wastewater and COVID-19 rates in days after sampling in small urban areas of Seville: A time series study. *Sci Total Environ* 2022; 806: 150573.
- Roka E, Khayer B, Kiss Z, Kovacs LB, Schuler E, Magyar N, et al. Ahead of the second wave: Early warning for COVID-19 by wastewater surveillance in Hungary. *Sci Total Environ* 2021; 786: 147398.
- Rosario K, Symonds EM, Sinigalliano C, Stewart J, Breitbart M. Pepper mild mottle virus as an indicator of fecal pollution. *Appl Environ Microbiol* 2009; 75: 7261-7.
- Salvo M, Moller A, Alvareda E, Gamazo P, Colina R, Victoria M. Evaluation of low-cost viral concentration methods in wastewaters: Implications for SARS-CoV-2 pandemic surveillances. *J Virol Methods* 2021; 297: 114249.

- Saththasivam J, El-Malah SS, Gomez TA, Jabbar KA, Remanan R, Krishnankutty AK, et al. COVID-19 (SARS-CoV-2) outbreak monitoring using wastewater-based epidemiology in Qatar. *Sci Total Environ* 2021; 774: 145608.
- Schmitz BW, Innes GK, Prasek SM, Betancourt WQ, Stark ER, Foster AR, et al. Enumerating asymptomatic COVID-19 cases and estimating SARS-CoV-2 fecal shedding rates via wastewater-based epidemiology. *Sci Total Environ* 2021; 801: 149794.
- Schober P, Boer C, Schwarte LA. Correlation Coefficients: Appropriate Use and Interpretation. *Anesth Analg* 2018; 126: 1763-1768.
- Scott LC, Aubee A, Babahaji L, Vigil K, Tims S, Avetisyan TG. Targeted wastewater surveillance of SARS-CoV-2 on a university campus for COVID-19 outbreak detection and mitigation. *Environ Res* 2021; 200: 111374.
- Shah S, Gwee SXW, Ng JQX, Lau N, Koh J, Pang J. Wastewater surveillance to infer COVID-19 transmission: A systematic review. *Sci Total Environ* 2022; 804: 150060.
- Tagliabue F, Galassi L, Mariani P. The "Pandemic" of Disinformation in COVID-19. *SN Compr Clin Med* 2020; 2: 1287-1289.
- Tandukar S, Sthapit N, Thakali C, Mallia B, Sherchan SP, Shakya BM, et al. Detection of SARS-CoV-2 RNA in wastewater, river water, and hospital wastewater of Nepal. *Sci Total Environ* 2022; 824: 153816.
- Tanhaei M, Mohebbi SP, Hosseini SM, Rafieepoor M, Kazemian S, Ghaemi A, et al. The first detection of SARS-CoV-2 RNA in the wastewater of Tehran, Iran. *Environ Sci Pollut Res Int* 2021; 28: 38629-38636.
- Tanimoto Y, Ito E, Miyamoto S, Mori A, Nomoto R, Nakanishi N, et al. SARS-CoV-2 RNA in Wastewater Was Highly Correlated With the Number of COVID-19 Cases During the Fourth and Fifth Pandemic Wave in Kobe City, Japan. *Front Microbiol* 2022; 13: 892447.
- Thongpradit S, Prasongtanakij S, Srisala S, Chanprasertyothin S, Pasomsub E, Ongphiphadhanakul B. The Detection of SARS-CoV2 Antigen in Wastewater Using an Automated Chemiluminescence Enzyme Immunoassay. *Int J Environ Res Public Health* 2022; 19.

- Tiwari A, Lipponen A, Hokajarvi AM, Luomala O, Sarekoski A, Ryttonen A, et al. Detection and quantification of SARS-CoV-2 RNA in wastewater influent in relation to reported COVID-19 incidence in Finland. *Water Res* 2022; 215: 118220.
- Tomasino MP, Semedo M, Vieira EMP, Ferraz E, Rocha A, Carvalho MF, et al. SARS-CoV-2 RNA detected in urban wastewater from Porto, Portugal: Method optimization and continuous 25-week monitoring. *Sci Total Environ* 2021; 792: 148467.
- Torii S, Furumai H, Katayama H. Applicability of polyethylene glycol precipitation followed by acid guanidinium thiocyanate-phenol-chloroform extraction for the detection of SARS-CoV-2 RNA from municipal wastewater. *Sci Total Environ* 2021; 756: 143067.
- Torii S, Oishi W, Zhu Y, Thakali O, Malla B, Yu Z, et al. Comparison of five polyethylene glycol precipitation procedures for the RT-qPCR based recovery of murine hepatitis virus, bacteriophage phiC, and pepper mild mottle virus as a surrogate for SARS-CoV-2 from wastewater. *Sci Total Environ* 2022; 807: 150722.
- Tranfield D, Denyer D, Smart P. Towards a methodology for developing evidence-informed management knowledge by means of systematic review. *British Journal of Management* 2003; 14: 207-222.
- Vallejo JA, Trigo-Tasende M, Rumbero-Feal S, Conde-Perez K, Lopez-Oriona A, Barbeito I, et al. Modeling the number of people infected with SARS-COV-2 from wastewater viral load in Northwest Spain. *Sci Total Environ* 2022; 811: 152334.
- van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med* 2020; 382: 1564-1567.
- Wang P, Zarei-Baygi A, Saucedo C, Iskander SM, Smith AL. Long-term surveillance of wastewater SARS-CoV-2 in Los Angeles County. *Environmental Science-Water Research & Technology* 2021; 7: 2282-2294.
- Wang Q, Waltman L. Large-scale analysis of the accuracy of the journal classification systems of Web of Science and Scopus. *Journal of Informetrics* 2016; 10: 347-364.

- Wehrendt DP, Masso MG, Gonzales Machuca A, Vargas CV, Barrios ME, Campos J, et al. A rapid and simple protocol for concentration of SARS-CoV-2 from sewage. *J Virol Methods* 2021; 297: 114272.
- Wei S, Suryawanshi H, Djandji A, Kohl E, Morgan S, Hod EA, et al. Field-deployable, rapid diagnostic testing of saliva for SARS-CoV-2. *Sci Rep* 2021; 11: 5448.
- Westhaus S, Weber FA, Schiwy S, Linnemann V, Brinkmann M, Widera M, et al. Detection of SARS-CoV-2 in raw and treated wastewater in Germany - Suitability for COVID-19 surveillance and potential transmission risks. *Sci Total Environ* 2021; 751: 141750.
- Wu F, Xiao A, Zhang J, Moniz K, Endo N, Armas F, et al. SARS-CoV-2 RNA concentrations in wastewater foreshadow dynamics and clinical presentation of new COVID-19 cases. *Sci Total Environ* 2022; 805: 150121.
- Wu Y, Guo C, Tang L, Hong Z, Zhou J, Dong X, et al. Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol* 2020; 5: 434-435.
- Xiao A, Wu F, Bushman M, Zhang J, Imakaev M, Chai PR, et al. Metrics to relate COVID-19 wastewater data to clinical testing dynamics. *Water Res* 2022; 212: 118070.
- Xiao F, Tang M, Zheng X, Liu X, Li X, Shan H. Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology* 2020a; 158: 1831-1833 e3.
- Xiao M, Liu X, Ji J, Li M, Li J, Yang L, et al. Multiple approaches for massively parallel sequencing of SARS-CoV-2 genomes directly from clinical samples. *Genome Med* 2020b; 12: 57.
- Xie YW, Challis JK, Oloye FF, Asadi M, Cantin J, Brinkmann M, et al. RNA in Municipal Wastewater Reveals Magnitudes of COVID-19 Outbreaks across Four Waves Driven by SARS-CoV-2 Variants of Concern. *Acs Es&T Water* 2022; 2: 1852-1862.
- Yanac K, Adegoke A, Wang L, Uyaguari M, Yuan Q. Detection of SARS-CoV-2 RNA throughout wastewater treatment plants and a modeling approach to understand COVID-19 infection dynamics in Winnipeg, Canada. *Sci Total Environ* 2022; 825: 153906.

- Ye Y, Ellenberg RM, Graham KE, Wigginton KR. Survivability, Partitioning, and Recovery of Enveloped Viruses in Untreated Municipal Wastewater. *Environ Sci Technol* 2016; 50: 5077-85.
- Yue H, Shin JM, Tegafaw T, Han HS, Chae KS, Chang YM, et al. Magnetic separation of nucleic acids from various biological samples using silica-coated iron oxide nanobeads. *Journal of Nanoparticle Research* 2020; 22: 366.
- Zhang D, Duran SSF, Lim WYS, Tan CKI, Cheong WCD, Suwardi A, et al. SARS-CoV-2 in wastewater: From detection to evaluation. *Mater Today Adv* 2022; 13: 100211.
- Zhang N, Gong Y, Meng F, Shi Y, Wang J, Mao P, et al. Comparative study on virus shedding patterns in nasopharyngeal and fecal specimens of COVID-19 patients. *Sci China Life Sci* 2021a; 64: 486-488.
- Zhang Y, Cen M, Hu M, Du L, Hu W, Kim JJ, et al. Prevalence and Persistent Shedding of Fecal SARS-CoV-2 RNA in Patients With COVID-19 Infection: A Systematic Review and Meta-analysis. *Clin Transl Gastroenterol* 2021b; 12: e00343.
- Zhao L, Zou Y, Li Y, Miyani B, Spooner M, Gentry Z, et al. Five-week warning of COVID-19 peaks prior to the Omicron surge in Detroit, Michigan using wastewater surveillance. *Sci Total Environ* 2022; 844: 157040.
- Zheng X, Deng Y, Xu X, Li S, Zhang Y, Ding J, et al. Comparison of virus concentration methods and RNA extraction methods for SARS-CoV-2 wastewater surveillance. *Sci Total Environ* 2022; 824: 153687.
- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 2020; 382: 727-733.
- Zhu Y, Oishi W, Maruo C, Bandara S, Lin M, Saito M, et al. COVID-19 case prediction via wastewater surveillance in a low-prevalence urban community: a modeling approach. *J Water Health* 2022; 20: 459-470.
- Zuo T, Liu Q, Zhang F, Lui GC, Tso EY, Yeoh YK, et al. Depicting SARS-CoV-2 faecal viral activity in association with gut microbiota composition in patients with COVID-19. *Gut* 2021; 70: 276-284.

Table 1. Search strings and Boolean operators used for each search.

	Boolean operator	Search strings	Category
Analytical methods		<i>COVID?19 OR SARS-CoV-2 OR coronavirus</i>	Topic
	AND	<i>wastewater OR ww or sewage</i>	Topic
	AND	<i>“SARS-CoV-2 RNA” OR RNA OR “ribonucleic acid” OR “nucleic acid” OR genet*</i>	Topic
	AND	<i>analy* OR method* OR procedure OR protocol OR techn*</i>	All Fields
	AND	<i>detect* OR concentrate* OR quantif* OR estimat* OR measur*</i>	All Fields
Mathematical modeling		<i>COVID?19 OR SARS-CoV-2 OR coronavirus</i>	Topic
	AND	<i>wastewater OR ww OR “wastewater-based epidemiology” or WBE</i>	Topic
	AND	<i>surveill* OR monitor* OR track*</i>	All Fields
	AND	<i>predict* OR forecast* OR forecast* OR model* OR correlate* OR relation*</i>	All Fields

Note: “?” denotes a wildcard and was used due to different spellings of the term adopted in the literature. The “*” symbol allows variations of the search string. Quotation marks strictly limit the appearance of the word as it is input.

Table 2. Description of analytical methods from selected literature and their contributions.

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Anderson-Coughlin et al. (2021)	USA August 2020 - March 2021	Raw (sewage)	Filtration using a 0.22µm PES membrane	Centrifugal ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	40 – 45	3
Ahmed et al. (2020a)	Australia March 2020 – April 2020	Raw (influent)	(1) pH adjustment to 3.5/4 using HCl, (2) Centrifugation at 4,750g for 30 minutes	(1) Adsorption-elution with electronegative membranes, (2) Ultrafiltration	QIAgen RNeasy PowerWater Kit and QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR N	100 – 200	5
Ahmed et al. (2020c)	Australia NR	Raw (influent)	(1) Acidification to pH 4 using 2N HCl, (2) NR, (3) MgCl ₂ addition to a final concentration of 25mM MgCl ₂ , (4,5) Centrifugation at 4,500g for 10 minutes at 4°C, (6) Centrifugation at 10,000g for 10 minutes at 4°C, (7) Centrifugation at 10,000g for 1h at 4°C	(1,2,3) Adsorption-elution using electronegative membranes, (4,5) Ultrafiltration, (6) PEG precipitation, (7) Ultracentrifugation	QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR NR	50	1,5
Ahmed et al. (2021a)	Bangladesh July 2020 – August 2020	Raw (sewage)	Centrifugation at 4,500g for 30 minutes, filtration using 0.22µm filters	PEG precipitation	Favor Prep Viral Nucleic Acid Extraction Kit	RT-qPCR N and ORF1ab	50	1,5
Ahmed et al. (2021b)	NR	Raw (influent)	(1) Centrifugation at 4,000g for 30 minutes at 4°C, (2) NR	(1) Concentrating pipette (InnovaPrep), (2) Adsorption-elution with electronegative membranes	QIAgen QIAamp® Viral RNA mini Kit and RNeasy PowerWater Kit	RT-qPCR N1	NR	1
Ahmed et al. (2021c)	Australia February 2020 – May 2020	Raw (influent)	NR	Adsorption-elution using electronegative membranes	QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR N1, N2 and N3	100 – 200	2
Ahmed et al. (2022b)	Australia June 2021	Raw (influent)	Centrifugation at 3,000g for 5 minutes	Concentration Pipette (InnovaPrep)	QIAgen QIAamp® Viral RNA mini Kit and QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR and RT-dPCR N1 and N2	50	4,6

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
					ome Kit (for the solid phase)			
Ai et al. (2021)	USA July 2020 - January 2021	Raw (influent)	Centrifugation at 2,500g for 10 minutes at 4°C, filtration using a 0.45µm sterile filter unit	Sequential concentration using adsorption-elution with positively charged membrane, organic flocculation, and centrifugal ultrafiltration	QIAgen RNeasy PowerMicroome Kit	RT-ddPCR N1, N2 and E	100 – 200	4,5,6
Amereh et al. (2022)	Iran September 2020 – April 2021	Raw (influent)	Centrifugation at 4,000g for 10 minutes	PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N and ORF1ab	50	4
Amoah et al. (2021)	South Africa NR	Raw (influent)	Heat inactivation 60°C for 90 minutes	Centrifugal ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	RT-ddPCR and RT-LAMP E, N, ORF1ab, RdRP and S	250	3
Anneser et al. (2022)	USA March 2020 – March 2021	Raw (influent and sludge)	NR	(1) PEG precipitation, (2) Spectrophotometry	TRIzol-chloroform protocol, RNeasy PowerSoil Total RNA Kit	RT-qPCR N1, N2 and N3	NR	4.5
Arora et al. (2020)	India May 2020 – June 2020	Raw (influent)	Heat inactivation 60°C for 90 minutes, filtration using a 0.45µm membrane	PEG precipitation	Allplex 2019-nCoV Assay Kit	RT-PCR N, S, E, ORF1ab and RdRp	50	2
Bagutti et al. (2022)	Switzerland July 2021 - December 2021	Raw (influent)	NR	NR	Maxwell® RSC Environ Wastewater TNA Kit	RT-qPCR N1, N2 and E	40	4
Baldovin et al. (2021)	Italy April 2020 – May 2020	Raw (influent) and treated (effluent: activated sludge, peracetic acid and UV lamps)	Filtration using a 0.22µm polyether sulfone (PES)	Ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N and ORF1ab	100	1,6
Barbosa et al. (2022)	Brazil May 2020 - October 2020	Raw (influent, sewage)	NR	Ultracentrifugation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	40	2,4,5

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Bar-Or et al. (2021a)	Israel August 2020 - February 2021	Raw (influent)	Centrifugation at 4,696g for 5 minutes	Adsorption-elution with electronegative membranes	NucliSENS EasyMAG	RT-qPCR E	25	2
Bar-Or et al. (2021b)	Israel March 2020 - April 2020	Raw (sewage)	NR	(1) PEG precipitation, (2) Skimmed milk flocculation, (3) Ultrafiltration	QIAgen RNeasy mini Kit, NucliSENS EasyMAG	RT-qPCR N and E	250 – 1,000	1,2,6
Barril et al. (2021)	Argentina March 2020 - October 2020	Raw (influent)	NR	A total of 11 different methods were evaluated.	Maxwell RSC 48 Extraction System	RT-qPCR N1 and N2	Varies from method to method	2,4
Barrios et al. (2021)	Argentina June 2020 – April 2021	Raw (influent)	Heat inactivation 60°C for 90 minutes	PEG precipitation	TRIzol-chloroform protocol	RT-qPCR N1	200	4
Barua et al. (2022)	USA June 2020 – November 2020	Raw (influent)	Heat inactivation 75°C for 40 minutes	HA electronegative filtration	QIAgen QIAamp® Viral RNA mini Kit and NucliSENS EasyMAG	RT-qPCR and RT-ddPCR N1 and N2	20	1
Bertrand et al. (2021)	France April 2020 - May 2020	Raw (influent after decantation)	NR	(1) Ultrafiltration, (2) PEG precipitation	Phenol-chloroform-isoamyl alcohol protocol	RT-PCR and RT-ddPCR E and RdRp	50	1,2
Bivins et al. (2022)	NR	Raw (influent, sewerage)	NR	Centrifugal ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit, AllPrep PowerViral DNA/RNA Kit	RT-ddPCR and RT-LAMP N2 and E	NR	1,3,4
Boogaerts et al. (2021)	Belgium August 2020 – January 2021	Raw (influent)	(1) Centrifugation at 4,600g for 30 minutes at 4°C, (2) Centrifugation at 4,654g for 30 minutes at 4°C	(1) Ultracentrifugation, (2) PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit, RNeasy plus miniKit, and QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR and RT-ddPCR N1, N2, N3 and E	20 – 90	1,4,5
Boogaerts et al. (2022)	Belgium September 2020 – November 2021	Raw (influent)	Centrifugation at 4,000g for 30 minutes	Ultracentrifugation	Maxwell® RSC PureFood GMO and Authentication Kit	RT-qPCR and RT-ddPCR N, S and E	20	3,6

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Calderon-Franco et al. (2022)	Netherlands July 2020 – December 2020	Raw (influent)	Heat inactivation 65°C for 30 minutes	(1) Adsorption-elution with electronegative membranes, (2) Polyethersulfone membranes, (3) Anion-exchange diethylamethyl cellulose columns	Fast RNA Blue Kit, FAST RNA Kit, MagMax CORE Nucleic Acid Purification Kit	RT-qPCR S, N and ORF1ab	50 – 550	1,3
Canh et al. (2021)	Japan January 2021 – February 2021	Raw (influent)	Centrifugation at 3,500g for 15 minutes	(1) Ultrafiltration, (2) PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1	100	1,2
Carrillo-Reyes et al. (2021)	Mexico April 2020 - July 2020	Raw (influent, sewage) and treated (secondary sludge, effluent)	Filtration using 0.2µm polyethersulfone membrane	(1) Ultrafiltration, (2) Adsorption-elution with electronegative membranes	QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR RdRp, S and E	(1) 120, (2) 30 – 100	1,2
Castiglioni et al. (2022)	March 2020 - June 2020	Raw (influent)	Under UV light for 30 minutes, Centrifugation at 4,500g for 30 minutes at 4°C	PEG precipitation	QIAgen QIAamp® MiniElute Virus Spin Kit	RT-PCR N1 and N3	45	1,2,5
Chakraborty et al. (2021)	India September 2020	Raw (influent) and treated (primary sludge, effluent)	NR	Composite, Supernatant, Sediment and Syringe Filtration	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	250	1
Chavarria-Miro et al. (2021)	Spain April 2020 – July 2020	Raw (influent)	NR	PEG precipitation	NucliSENS miniMAG	RT-qPCR N1, N2, RdRp, IP2 and IP4	800	5
Claro et al. (2021)	Brazil June 2020 – April 2021	Raw (influent)	Centrifugation at 8,000g for 120 minutes at 4°C	PEG precipitation	PureLink™ Viral RNA/DNA mini Kit	RT-qPCR N1 and N2	40	4
D'Aoust et al. (2021)	Canada April 2020 - June 2020	Raw (post-grid influence) and treated (primary clarified sludge)	Decantation and serially filtered through a 1.5µm glass fiber filter followed by a 0.45µm GF6 mixed cellulose-ester filter	PEG precipitation	QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR and RT-ddPCR N1 and N2	32	3,5
de Freitas Bueno et al. (2022)	Brazil January 2021 – January 2022	Raw (influent)	NR	PEG precipitation	PureLink Viral RNA/DNA mini Kit	RT-qPCR N1 and N2	40	4.5

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
de Sousa et al. (2022)	Brazil January 2021 – August 2021	Raw (influent) and treated (effluent)	pH adjustment to 3.5 using 1M HCl, shaken at 4°C for 30 minutes, Centrifugation at 2,474g for 30 minutes at 4°C	PEG precipitation	MagMAX Viral/Pathogen Nucleic Acid Isolation Kit	RT-qPCR N1 and N2	50	5
Dimitrakopoulos et al. (2022)	Greece November 2021 - December 2021	Raw (influent)	NR	(1) PEG precipitation, (2) PEG precipitation with glycine, (3) Direct capture, (4) Adsorption-elution with electronegative membranes, (5) Ultrafiltration	Water DNA/RNA magnetic bead Kit, QIAzol RNeasy PowerMicrobiome Kit, AllprepViral DNA/RNA Kit, Manual EnviroWaste water TNA Kit	RT-qPCR and RT-ddPCR N1, N2 and N3	50	2,6
Dumke et al. (2021)	Germany, NR	Raw (influent)	Centrifugation at 3,300g for 30 minutes at 4°C	(1) PEG precipitation, (2) Centrifugation with Vivaspin columns	QIAgen RNeasy kits (not specified what series)	RT-qPCR and RT-ddPCR S and E	40	2
Farkas et al. (2021)	NR	Raw (NR)	Centrifugation at 3,000g for 30 minutes at 4°C or 1,000g for 10 minutes at 4°C, pH adjustment of supernatant to 7-7.5 using 1M NaOH	PEG precipitation	NucliSENS lysis buffer and NucliSENS miniMag extraction system	RT-qPCR N1 and N2	50	1,3
Feng et al. (2021)	USA August 2020 - January 2021	Raw (influent)	Filtration using 0.8µm cellulose-ester filters	Bashing Bead Lysis	QIAgen RNeasy PowerMicrobiome Kit	RT-ddPCR N1 and N2	25	1
Fernandez-Cassi et al. (2021)	Switzerland February 2020 - April 2020	Raw (influent)	Filtered using 2µm glass fiber filters	Centrifugal ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	50	5
Fitzgerald et al. (2021)	Scotland April 2020 – January 2021	Raw (influent)	Centrifugation at 4,000g for 30 minutes at 4°C, filtration using a syringe filter	(1) Ultracentrifugation, (2) PEG precipitation, (3) skimmed milk flocculation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and E	20 – 40	1.2
Flood et al. (2021)	USA March 2020 – September 2020	Raw (influent, sewage)	(1) Centrifugation at 2,500g for 5 minutes at 4°C, (2) Centrifugation at	(1,2) Ultrafiltration, (3) PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR, RT-ddPCR N1, N2 and E	(1) 100, (2) 100, (3) NR	1

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
			4,654g for 30 minutes at 4°C, (3) Centrifugation at 4,700g for 45 minutes at 4°C					
Fongaro et al. (2021)	Brazil October 2019 - March 2020	Raw (sewage)	NR	PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1, S and RdRp	25	4
Fonseca et al. (2022)	Brazil March 2021	Raw (influent, river)	Heat inactivation 60°C for 90 minutes, filtration using 1.2µm pore size microfiber filters, Centrifugation at 4,500g for 30 minutes at 4°C	(1) Ultrafiltration, (2) Adsorption-elution with electronegative membranes (3) Aluminum hydroxide precipitation, (4) PEG precipitation	MonView Viral/Pathogen Mini Kit, KingFisher Duo Purification System	RT-qPCR N1 and N2	40	4,6
Galani et al. (2022)	Greece August 2020 – March 2021	Raw (influent)	Centrifugation at 4,700g for 30 minutes at 4°C	(1) PEG precipitation, (2) centrifugal ultrafiltration	Water DNA/RNA Magnetic Bead Kit, QIAgen RNeasy Power Microbiome Kit, and QIAgen RNeasy Serum/Plasma Advanced Kit	RT-qPCR N1 and N2	50	4,5
Gerrity et al. (2021)	USA March 2020 - May 2020	Raw (influent)	(1) NR, (2,3) Centrifugation at 3,500g for 15-30 minutes at 10°C	(1) Hollow-fiber ultrafiltration, (2) Centrifugal Ultrafiltration, (3) PEG precipitation.	Purelink Viral RNA/DNA mini Kit	RT-qPCR N1, N2, E and ORF1a	50	1,3
Giraud-Billoud et al. (2021)	Argentina July 2020 - November 2020	Raw (influent)	Heat inactivation 60°C for 90 minutes	(1) PEG precipitation, (2): Polyaluminum chloride (PAC) flocculation	NucleoZOL	RT-qPCR N1 and N2	300	5,6
Goncalves et al. (2021)	Slovenia June 2020	Raw (sewage)	Filtration using a 0.70µm glass fiber filter membrane	Ultracentrifugation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR E and RdRp	100	1
Gonzalez et al. (2020)	USA March 2020 – August 2020	Raw (influent)	(1) Centrifugation at 10,000g for 10 minutes, (2) NR	(1) Concentration Pipette (InnovaPrep), (2) Adsorption-	NucliSENS Easy Mag TNA Extraction Kit	RT-ddPCR N1, N2 and N3	125	2

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
				elution using electronegative membranes				
Gonzalez-Reyes et al. (2021)	Mexico June 2020 - July 2020	Raw (influent, sewage)	Heat inactivation 60°C for 90 minutes, filtration using a 0.2µm membrane	PEG precipitation	TRIzol protocol	RT-qPCR N1, N2 and N3	150	2
Haramoto et al. (2020)	Japan March 2020 - May 2020	Raw (influent) and treated (activated sludge before chlorination)	NR	(1) Adsorption-elution with electronegative membranes, (2) Direct adsorption	(1) QIAgen QIAamp [®] Viral RNA mini Kit, (2) QIAgen PowerMicrobiome Kit	RT-qPCR N1, N2, S and ORF1ab	200 – 5000	2
Hasan et al. (2021)	UAE May 2020 - June 2020	Raw (influent) and treated (effluent)	(1,2) Heat inactivation 60°C for 90 minutes, filtration using a 0.22µm polyethersulfone (PES) membrane	(1) Ultrafiltration, (2) PEG precipitation	ASIOpure Viral DNA/RNA Extraction Kit and TRIzol-chloroform protocol	RT-qPCR RdRp	(1,2) 50	1,4
Hasing et al. (2021)	Canada October 2020 - December 2020	Raw (influent)	pH adjustment to 9.6-10 using 5N NaOH, Centrifugation at 4,500g for 10 minutes	Ultrafiltration	MagMAX96 Viral RNA Isolation Kit, King Fisher Flex Purification System	RT-qPCR N2 and E	100	3
Hata et al. (2021)	Japan March 2020 – April 2020	Raw (influent)	NR	PEG precipitation	QIAgen QIAamp [®] Viral RNA mini Kit	RT-qPCR N2 and N3	80	2
Hemalatha et al. (2021)	India July 2020 – August 2020	Raw (influent)	Gravity filtration with 1mm thick blotting sheets to remove debris and larger particles followed by filtration using 0.2µm filtration units	Centrifugal ultrafiltration	QIAamp [®] Viral RNA isolation Kit	RT-qPCR N, E and ORF1ab	100	2,3
Hoar et al. (2022)	USA April 2020 – February 2021	Raw (influent)	Heat inactivation 60°C for 90 minutes, Centrifugation at 5000g for 10 minutes at 4°C, filtration using 0.22µm acetate-cellulose membrane	PEG precipitation	QIAgen QIAamp [®] Viral RNA mini Kit	RT-qPCR N1	40	4,5
Hokajarvi et al. (2021)	Finland April 2020 – May 2020	Raw (influent)	Centrifugation at 4,654g for 30 minutes	Ultrafiltration	Chemagic Viral300 DNA/RNA extraction Kit	RT-qPCR N2 and E	60	2,3

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Huang et al. (2021)	Canada October 2020 – March 2021	Raw (influent)	NR	Ultrafiltration	QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR N1, N2, N3 and E	200	2
Iglesias et al. (2021)	Argentina June 2020 - September 2020	Raw (influent, surface water)	Heat inactivation 60°C for 90 minutes	PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	250	2
Jafferli et al. (2021)	Sweden and Italy May 2020 - June 2020	Raw (influent)	(1) Centrifugation at 4,600g for 30 minutes at 4°C, (2) centrifugation at 1,500g for 15 minutes at 4°C, (3) NR, (4) Centrifugation at 4,600g for 30 minutes at 4°C	(1) Ultrafiltration, (2) Double Ultrafiltration, (3) Adsorption-elution with electronegative membrane (4) Centrifugation combined with adsorption-extraction	TRIzol reagent, RNeasy PowerMicrobiome Kit	RT-qPCR N	40 – 50	1
Jmii et al. (2021)	Tunisia September 2020 - October 2020	Raw (influent)	Coarse filtration and microfiltration, pH adjustment to 6 with aluminum hydroxide	Adsorption-elution with electronegative membranes	QIAgen RNeasy PowerMicrobiome Kit	RT-PCR N, E and RdRp	100	1
Johnson et al. (2021)	South Africa June 2020	Raw (influent)	Centrifugation at 3,500g for 20 minutes	NR	RNeasy PowerSoil Kit	RT-qPCR N1 and N2	50-100	2
Juel et al. (2021)	USA October 2022 – March 2021	Raw (sewage)	NR	(1) Adsorption-elution with electronegative membranes, (2): Concentrating Pipette (InnovaPrep)	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1	40 – 100	1,3
Kevill et al. (2022)	Wales October 2020 – February 2021	Raw (influent)	Centrifugation at 15,000g for 10 minutes at 4°C	(1) PEG precipitation, (2) Ammonium sulfate precipitation, (3) Concentration pipette (Innova Prep)	NucliSENS Lysis Buffer, NucliSENS Extraction Reagent Kit, King-Fisher 96 Flex System	RT-pPCR N1	200	1,3
Kitamura et al. (2021)	Japan June 2020 - August 2020	Raw (influent, sewage)	Centrifugation at 3,000rpm for 30 minutes	(1) Adsorption-elution with electronegative membranes, (2) PEG	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	400	1,3,4,5

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
				precipitation, (3) Ultrafiltration, (4) Solid precipitation				
Koureas et al. (2021)	Greece October 2020 - April 2021	Raw (influent)	NR	PEG precipitation	MagMAX™ Viral/Pathogen Nucleic Acid Isolation Kit	RT-PCR N, S and ORF1ab	105	5
Krivanakova et al. (2021)	Slovakia September 2020 – March 2021	Raw (influent)	Centrifugation at 4,700g for 30 minutes	Ultracentrifugation	Direct-zul RNA miniprep Kit	RT-qPCR E, RdRp and ORF1ab	50	2.4
Kuhn et al. (2022)	USA November 2020 – March 2021	Raw (sewage)	Filtration using a 70µm mesh cell strainer	PEG precipitation	Bio-Cli-Magnetic-Beads platform	RT-qPCR N1	32	5
Kumar et al. (2021)	India August 2020 - September 2020	Raw (influent)	Centrifugation at 4,000g for 40 minutes, filtration using a 0.22µm syringe filter	PEG precipitation	NucleoSpin® RNA Virus isolation Kit	RT-PCR N, S and ORF1ab	30	2
La Rosa et al. (2020)	Italy February 2020 - April 2020	Raw (influent)	Heat inactivation 56°C for 30 minutes	PEG-dextran two-phase separation	NucliSENS miniMAG	RT-qPCR ORF1ab, S and RdRp	250	2,5
Langan et al. (2022)	USA January 2021 - March 2021	Raw (sewage)	Centrifugation at 4,000g for 20 minutes at 4°C	Ultrafiltration	QIAGEN PowerViral DNA/RNA Kit, Zymo EnviroN Water RNA Extraction Kit, Monarch Total RNA miniprep Kit	RT-qPCR N1 and N2	200	1,5
Lara-Jacobo et al. (2022)	Canada October 2020 - April 2021	Raw (influent)	Adding 50 of acetone at 4°C and stored overnight at -2°C to precipitate proteins, Centrifugation at 3,405g for 15 minutes	Protein Precipitation and Digestion	QIAGEN PowerMicrobiome Kit	RT-qPCR N1	40	1,3
LaTurner et al. (2021)	USA October 2020	Raw (influent)	(1) NR, (2) Centrifugation at 4,100g for 10 minutes at 4°C, (3) Centrifugation at 3,000g for 1 minute at 4°C, (4) Centrifugation at 7,140g for 15 minutes at 4°C,	(1) Direct extraction, (2) HA filtration with bead beating, (3) HA filtration with elution, (4) PEG precipitation, (5) Ultrafiltration	Chemagic Prime Viral DNA/RNA 300 Kit H96	RT-qPCR and RT-ddPCR N1 and N2	(1) 1, (2) 50, (3) 50, (4) 200, (5) 50	5

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
			(5) Centrifugation at 4,100g for 10 minutes at 4°C					
Layton et al. (2022)	USA June 2020 – July 2020	Raw (sewage)	Centrifugation at 12,000g for 1 minute	NR	MagMAX Viral Pathogen Kit	RT-ddPCR N1 and N2	30 – 40	4
Li et al. (2022)	USA June 2020 – September 2021	Raw (influent)	Heat inactivation 60°C for 60 minutes, Centrifugation at 3,000g for 15 minutes, and sequential filtration using 1.5, 0.8, and 0.45µm sterile membrane filters	PEG precipitation	AllPrep PowerViral DNA/RNA Kit	RT-qPCR N1 and N2	NR	2,4
Maida et al. (2022)	Italy September 2021 - July 2021	Raw (sewage)	NR	PEG-dextran two-phase separation	NucliSENS miniMAG	RT-qPCR NR	NR	5
Mailepessov et al. (2022)	Singapore April 2020	Raw (sewage)	(1) Centrifugation at 4,000g for 30 minutes, (2) Centrifugation at 2,000g for 5 minutes	(1) PEG precipitation, (2) Ultrafiltration	(1) modified TRIzol-QIAgen protocol, (2) QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR NR	45	1,3
Markt et al. (2022)	Liechtenstein Sept 2020 - March 2021	Raw (influent)	Centrifugation at 4,000g for 30 minutes	PEG precipitation	Monarch total RNA miniprep Kit	RT-qPCR N1	70	1
Masachessi et al. (2022)	Argentina May 2020 - August 2021	Raw (influent)	Centrifugation at 4,750g for 20 minutes at 4°C	PEG precipitation	MagNa Pure 96 DNA and Viral NA Large Volume Kit	RT-qPCR N and E	500	2
McMahan et al. (2021)	USA May 2020 – August 2020	Raw (sewage)	Heat inactivation 60°C for 30 minutes, Centrifugation at 6,500g for 10 minutes at 6°C	PEG precipitation	TRIzol-chloroform protocol	RT-qPCR N	225	5
McMinn et al. (2021)	USA July 2020 - October 2020	Raw (influent, primary treated)	Heat inactivation 121°C for 60 minutes	(1) Ultrafiltration, (2) Concentration Pipette	QIAgen All Prep PowerViral Kit	RT-qPCR N	2000	1
Mlejnkova et al. (2020)	Czech Republic April 2020 - June 2020	Raw (influent)	NR	Skimmed milk flocculation	NucliSENS miniMAG	RT-qPCR NR	500	1

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Mondal et al. (2021)	USA October 2020 - Jan 2021	Raw (influent)	NR	Direct Capture	NR	RT-qPCR N1, N2 and E	40	1
Monteiro et al. (2022)	Portugal April 2020 - December 2020	Raw (influent)	Hollow-fiber filtration	PEG precipitation	QIAgen QIAamp® Fast DNA Stool mini Kit	RT-qPCR N, E and RdRp	1000	1
Nagarkar et al. (2022)	USA May 2020 – November 2020	Raw (influent)	NR	Ultrafiltration	RNeasy PowerWater Kit	dd-PCR N1 and N2	225	4
Nasseri et al. (2021)	Iran April 2020 - May 2020	Raw (influent) and treated (effluent)	Decantation for 5 minutes, Centrifugation at 1,500g for 20 minutes at 4°C, pH adjustment to 7-7.5 using HCl and NaOH	PEG-dextran two-phase separation	Fasture Viral RNA mini Kit	RT-PCR N and ORF1ab	250	2
Navarro et al. (2021)	Italy December 2020 – February 2021	Raw (influent)	Centrifugation at 4,500g for 30 minutes at 4°C	Centrifugal ultrafiltration	Quick-RNA Fecal/Soil Microbe Microprep	RT-qPCR N1, N3 and S	100	1
Ni et al. (2021)	Australia March 2020 - April 2020	Raw (influent)	Centrifugation at 9,000g for 20 minutes at 4°C	Ultrafiltration	QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR N1 and N2	50	3,6
Nourbakhsh et al. (2022)	Canada September 2020 - June 2021	Raw (influent)	(1,2,3) Centrifugation at 4,000g for 20 minutes at 4°C	(1) Centrifugal ultrafiltration, (2) zirconia-silica beads in a Bead Mill 24 Homogenizer, (3) Centrifugation	MagNA Pure 96 DNA, Viral NA Large Volume Kit and QIAgen RNeasy PowerMicrobiome Kit	RT-qPCR N1 and N2	15 – 30	5
Novoa et al. (2022)	Spain May 2020 - May 2021	Raw (influent, sewerage) and treated (effluent)	Filtration using a 20–25µm cellulose filter, pH adjustment to 6	Adsorption-precipitation with AlCl ₃	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1, N2 and E	150	5,6
O'Brien et al. (2021)	USA June 2020	Raw (sewage)	NR	Ultrafiltration	QIAgen All Prep PowerViral DNA/RNA KIT, Monarch RNA miniprep Kit, Zymo Quick RNA-Viral	RT-qPCR N2	250	5
Parra-Guardado et al. (2022)	Canada NR	Raw (influent)	Centrifugation at 5,000 rpm for 5 minutes	NR	Direct Magnetic Bead Extraction	RT-qPCR NR	50	1

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Peinado et al. (2022)	Spain February 2021 – June 2021	Raw (influent)	(1) Centrifugation at 4,600g for 30 minutes, pH adjustment to 6, (2) Centrifugation at 8,000g for 30 minutes at 4°C, (3) Centrifugation at 4,600g for 30 minutes	(1) Adsorption-precipitation with AlOH ₃ , (2) PEG precipitation, (3) Ultrafiltration	NZY Viral RNA Isolation Kit	RT-qPCR N1 and N2	100 – 200	1
Pellegrinelli et al. (2022)	Italy March 2019 - December 2020	Raw (influent)	(1) Centrifugation at 4,500g for 30 minutes, (2) Centrifugation at 4,500g for 30 minutes at 4°C, (3) Centrifugation at 1,200g for 30 minutes at 4°C	(1) PEG-Dextran two-phase separation, (2) PEG precipitation chloroform purification, (3) PEG precipitation chloroform purification	QIAamp® miniElute Virus Spin Kit, NucliSENS EasyMAG	RT-PCR N1, N3 and ORF1ab	(1) 250, (2) 80, (3) 250ML	2
Perez-Cataluna et al. (2021)	NR	NR	(1) NR, (2): Centrifugation at 2,500g for 10 minutes at 4°C	(1) Aluminum-based adsorption-precipitation, (2) PEG precipitation	NucleoSpin RNA Virus Kit	RT-qPCR N1, N2, E, IP2 AND IP4	200	1
Petala et al. (2022)	Greece October 2020 – January 2021	Raw (influent)	pH adjustment to 4 using 2M HCl, Centrifugation at 4,000g for 30 minutes	Adsorption-elution using electronegative membranes	Phenol-chloroform-based RNA extraction protocol	RT-PCR N2 and E	200	5
Philo et al. (2021)	USA March 2020 - July 2020	Raw (influent after sedimentation)	NR	(1) Bag-mediated Filtration System (BMFS), (2) Skimmed milk flocculation, (3) PEG precipitation, (4) Ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1, N2 and N3	(1) 100, (2) 500, (3) 1000	1
Philo et al. (2022)	October 2020 - March 2021	Raw (influent after sedimentation)	NR	Skimmed milk flocculation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	50	1
Pillay et al. (2021)	South Africa July 2020 – October 2020	Raw (influent)	Heat activation 60°C for 90 minutes, Centrifugation at 3,500g for 10 minutes	Ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	dd-PCR N2	250	5

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Pino et al. (2021)	Colombia NR	Raw (influent)	NR	(1) Flocculation with AlCl ₃ , (2) PEG precipitation, (3) Flocculation with skimmed milk, (4) Ultrafiltration	EZNA Total RNA Kit	RT-qPCR N, E and RdRp	200	2
Prakash (2021)	India June 2020 - July 2021	Raw (sewage)	(1) Centrifugation at 4,700g for 30 minutes, (2) Precentrifugation at 5,000rpm for 30 minutes, (3) NR	(1) Ultrafiltration, (2) PEG precipitation (3) PEG-dextran two-phase separation	QIAGEN RNeasy PowerMicrobiome Kit	RT-qPCR N1, N2 and E	200 – 550	1
Qiu et al. (2022)	Canada May 2020	Raw (influent)	pH adjustment to 9.6-10 using 5N NaOH, Centrifugation at 4,500g for 10 minutes, Removal of supernatant and pH readjustment to 7	Centrifugal ultrafiltration	QIAGEN RNeasy PowerMicrobiome Kit, MagMAX-96 Viral RNA Isolation Kit, MagMAX Viral/Pathogen Viral RNA mini Kit, and ReliaPrep RNA miniprep System	RT-qPCR N1, N2, E and RdRp	100	3
Ramos-Mandujano et al. (2021)	Saudi Arabia June 2020	Raw (sewage)	NR	(1) Adsorption-elution with electronegative membranes, (2) Silica-coated Magnetic nanoparticles	QIAamp® RNA mini Kit	RT-PCR N1 and N2	300 – 500	1,3
Randazzo et al. (2020)	Spain March 2020 – April 2020	Raw (influent) and treated (secondary and tertiary effluents)	NR	Aluminum hydroxide adsorption-precipitation	NucleoSpin RNA virus Kit	RT-qPCR N1, N2 and N3	200	1
Reynolds et al. (2022)	Ireland September 2020 – March 2021	Raw (influent)	Centrifugation at 3,200g for 5 minutes	Ultracentrifugation	QIAGEN RNeasy PowerMicrobiome Kit	RT-qPCR and dd-PCR N1	200 – 225	2.6
Robotto et al. (2022)	Italy July 2020 - March 2021	Raw (influent)	NR	NR	Wastewater Large Volume Total Nucleic Acid Capture Kit AX9550	RT-qPCR N1, N2 and E	40	1,3

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Rocha et al. (2022)	USA July 2020 - May 2021	Raw (influent)	2.5M MgCl ₂ was added at a ratio of 1:100 to a final concentration of 25mM	Adsorption-elution with electronegative membranes	QIAgen PowerViral Kit	RT-qPCR N1 and N2	40 – 495	3,4,5
Rodriguez Rasero et al. (2022)	Spain July 2020 – February 2021	Raw (sewage)	pH adjustment to 6 using 2N HCl	AlCl ₃ precipitation	NucleoSpin RNA Virus Kit and QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N, E and IP4	200	5
Roka et al. (2021)	Hungary June 2020 – October 2020	Raw (influent)	(1) NR, (2) Centrifugation at 4,500g for 30 minutes at 4°C	(1) Skimmed milk flocculation, (2) Ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N	(1) 50, (2) 50	1,4
Rondeau et al. (2021)	USA NR	Raw (sewage)	Heat inactivation 60°C for 1h, filtering using 0.22µm filter	Centrifugal ultrafiltration	Quick RNA miniprep Kit	RT-qPCR N1	40	3
Rosiles-Gonzalez et al. (2021)	Mexico August 2020 - January 2021	Treated (primary, biofilter and biological treatment)	(1) Filtration using a 0.45µm cellulose-ester membrane, (2,3) Sequential filtration using 0.8, 0.65, 0.45 and 0.22µm cellulose-ester membranes	(1) Adsorption-elution with electronegative membranes, (2) PEG precipitation, (3) Centrifugal filtration.	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1 and N2	(1) 1.0 - 5.4, (2) 200 – 1000, (3) 0.6-1.3	1,3
Sapula et al. (2021)	Australia NR	Raw (influent)	(1) Centrifugation at 5,000g for 30 minutes at 4°C, (2) adding MgCl ₂ to a final concentration of 25mM	(1) PEG precipitation, (2) Adsorption-elution with electronegative membranes	TRIzol-phenol extraction, NucleoSpin RNA Virus Extraction Kit, RNeasy PowerWater Kit	RT-qPCR N1 and N2	100	1,3
Saththasivam et al. (2021)	Qatar June 2020 – August 2020	Raw (influent)	Heat inactivation 56°C for 30 minutes, Centrifugation 4,500g for 30 minutes at 4°C	PEG precipitation	Quick RNA Viral Kits (Zymo)	RT-qPCR N1, N2 and RdRp	200	2,5
Scott et al. (2021)	USA August 2020 – December 2020	Raw (sewage)	NR	PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR and dd-PCR N1 and N2	200	1
Sharma et al. (2021)	India May 2020 – May 2020	Raw (sewage)	Chloroform was added and mixed thoroughly using a magnetic stirrer for 30 minutes at 4°C, Centrifugation at 3,000g for 20 minutes at 4°C	PEG-dextran phase separation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR E and RdRp	500	2

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Sherchan et al. (2020)	USA January 2020 - April 2020	Raw (influent) and treated (secondary treatment, effluent)	(1) Centrifugation at 3,000g for 30 minutes, (2) NR	(1) Ultrafiltration, (2) Adsorption-elution with electronegative membranes	ZR Viral RNA Kit	RT-qPCR N1 and N2	100 – 1000	1
Song et al. (2021)	USA April 2020 – June 2020	Raw (influent)	(1) Heat inactivation 60°C for 90 minutes, Centrifugation at 4,000g for 30 minutes, filtration using 0.45µm sterile membrane filter, (2) NR	(1) PEG precipitation, (2) Direct extraction method	QIAgen QIAamp® Viral RNA mini Kit and Zymo Quick-RNA Fecal/Soil Microbe Miniprep Kit	RT-qPCR and ddPCR N1 and N2	50	1,2,6
Tandukar et al. (2022)	Nepal July 2020 - February 2021	Raw (influent, sewage)	NR	Electronegative membrane vortex (EMV)	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N1, N2 and E	100	1,3
Tanhaei et al. (2021)	Iran June 2020 - July 2020	Raw (influent) and treated (effluent)	NR	Adsorption-elution with electronegative membranes	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N and ORF1ab	200	2
Tanimoto et al. (2022)	Japan February 2021 – October 2021	Raw (influent)	Centrifugation at 10,000g for 30 minutes	PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N	40	4
Thongpradit et al. (2022b)	Thailand January 2021 - February 2021	Raw (sewage)	Centrifugation at 3,000g for 10 minutes at room temperature	Adsorption-elution using electronegative membranes	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N, S and ORF1ab	100 – 400	2
Tiwari et al. (2022)	Finland August 2020 - May 2021	Raw (influent)	Centrifugation at 3,000g for 25 minutes	Ultrafiltration	Chemagic Viral300 DNA/RNA Extraction Kit	RT-qPCR E and N2	NR	1
Toledo et al. (2022)	USA Sept 2020 - Feb 2021	Raw (influent)	Centrifugation at 4,600g for 30 minutes at 4°C	PEG precipitation	Promega Wastewater Large-Volume TNA Capture Kit	RT-qPCR and RT-ddPCR N1 and N2	45	1
Tomasino et al. (2021)	Portugal May 2020 - March 2021	Raw (influent)	pH adjustment to 3.5/4 using HCl, Heat inactivation 60°C for 90 minutes	(1) NR, (2) Sequential centrifugations followed by PEG precipitation	QIAgen RNeasy Powersoil Total RNA, QIAgen RNeasy PowerMicrobiome Kit, IDXX DNA/RNA Magnetic Bead Kit	RT-qPCR N1 and N2	(1) 10 - 80, (2) 35	3
Torii et al. (2021)	Japan NR	Raw (influent)	(1) Centrifugation at 3,500g for 15 minutes, (2)	(1) Ultracentrifugation, (2) Electronegative	QIAgen QIAamp® Viral RNA mini Kit and	RT-qPCR N1, N2 and N3	40-50	2

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
			filtration using a through 0.45µm cellulose-ester membrane, (3) Centrifugation at 3,500g for 5 minutes	membrane vortex, (3) PEG precipitation	acid guanidium thiocyanate-phenol-chloroform extraction using TRIzol protocol			
Torii et al. (2022)	Japan July 2020 - October 2020	Raw (influent)	(1,2) Centrifugation at 3,500g for 5 minutes, (3) Centrifugation at 4,700g for 30 minutes at 4°C, (4) filtration using a 0.2µm hydrophilic polytetrafluoroethylene membrane (Millipore), (5) NR	PEG precipitation	QIAgen QIAamp Viral RNA mini Kit	RT-qPCR N1 and N2	41	1,2
Trottier et al. (2020)	France May 2020 - July 2020	Treated (effluent)	Centrifugation at 4,500g for 30 minutes at 4°C	Centrifugal filtration	NucleoSpin RNA Virus Kit	RT-qPCR N1, N2 AND RLP27	50	3
Trujillo et al. (2021)	NR	NR	Heat inactivation 60°C for 60 minutes, filtration using a 0.22µm filter	PEG precipitation	TRIzol-chloroform protocol	RT-qPCR N1	40	3
Vallejo et al. (2022)	Spain March 2020 – May 2020	Raw (influent)	Centrifugation at 4,000g for 30 minutes, filtration using 0.22µm membranes	Ultrafiltration	QIAgen QIAamp® Viral RNA mini Kit	RT-qPCR N	100	4,5
Wehrendt et al. (2021)	Argentina April 2021 - May 2021	NR	(1) Centrifugation at 12,000g for 1h at 4°C, (2) pH adjustment to 6-7	(1) PEG precipitation, (2) Centrifugation with polyaluminum chloride (PAC)	High Pure Viral Nucleic Acid Kit, Viral Nucleic Extraction Kit II	RT-qPCR N and ORF1	(1) 200, (2) 40	1,3,4
Westhaus et al. (2021)	Germany April 2020	Raw (influent after sand trap) and treated (activated sludge)	Centrifugation at 4,700g for 30 minutes	Ultracentrifugation	NucleoSpin RNA virus Kit	RT-qPCR N, E and RdRp	45	2
Whitney et al. (2021)	USA NR	Raw (influent)	NR	NR	4S-column and 4S-Milk-of-Silica	RT-qPCR N1	40	1
Wu et al. (2022)	USA January 2020 – May 2020	Raw (influent)	Heat inactivation 60°C for 90 minutes, filtration using a 0.2µm sterile membrane filter	PEG precipitation	TRIzol-chloroform protocol	RT-qPCR N1 and N2	40	4,5

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
Xiao et al. (2022)	USA March 2020 – June 2020	Raw (influent)	Heat inactivation 60°C for 1 hour, filtration using a 0.2µm vacuum-driven filter	Centrifugal ultrafiltration	NR	RT-qPCR N1 and N2	15	5
Xu et al. (2021)	Hong Kong June 2020 – September 2020	Raw (influent and sewage)	Heat inactivation 60°C for 30 minutes, Centrifugation at 4,750g for 30 minutes	Ultrafiltration	TRIzol Plus RNA Purification Kit	RT-qPCR N	50 – 90	1
Yanac et al. (2022)		Raw (influent) and treated (primary sludge, secondary effluent, final effluent)	(1) Cheesecloth and low-protein binding 0.45 and 0.2µm 47-mm Supor-200 membrane disc filters, (2) NR	(1) Ultrafiltration, (2) Skimmed milk flocculation	(1) QIAgen RNeasy PowerMicrobiome Kit, (2) MagMAX Microbiome Kit	RT-qPCR N1 and N2	120	1,4,5
Yaniv et al. (2021)	Israel November 2020 – March 2021	Raw (influent)	Shaken and mixed for 2 minutes manually and left standing 15 minutes to large particle settlement	Ultrafiltration	NucleoSpin RNA Extraction Kit	RT-qPCR N1, N2, N3 and N4	2000 – 5000	1,5
Zhang et al. (2022)	Australia August 2020 – September 2020	Raw (influent)	NR	Adsorption-elution with electronegative membrane	QIAgen RNeasy PowerWater Kit	RT-qPCR N and E	100	2
Zhao et al. (2022)	USA September 2020 – August 2021	Raw (influent)	NR	PEG precipitation	QIAgen QIAamp® Viral RNA mini Kit	RT-ddPCR N1 and N2	NR	4
Zheng et al. (2022)	Hong Kong September 2020 – November 2020	Raw (influent and sewage)	Heat inactivation 60°C for 30 minutes	(1) Ultracentrifugation, (2) PEG precipitation, (3) AlCl ₃ flocculation, (4) MgCl ₂ flocculation, (5) Ultracentrifugation 10kDa, (6) Ultracentrifugation 30kDa, (7) Membrane adsorption with AlCl ₃ , (8) Adsorption-elution using electronegative membranes,	QIAgen QIAamp® Viral RNA mini Kit and TRIzol Plus RNA Purification Kit	RT-qPCR N1	30 – 1000	1

Study	Location and sampling period	Wastewater type and sources	Sample pre-treatment	Concentration methods	RNA extraction Kit/protocol	Quantification method and gene targets	Analyzed initial sample volume (mL)	Type of contribution (Fig. 2)
				(9) Combination of centrifugation and ultracentrifugation, (10) AlCl ₃ precipitation, (11) Membrane adsorption (last 3 methods for 1000)				
Zhu et al. (2022)	Japan August 2020 – February 2021	Raw (influent)	NR	Ultracentrifugation	QIAgen Qiamp® Viral RNA mini Kit	RT-qPCR N1	40	4,5
Note: NR stands for Not Reported.								

Table 3. Summary of variables involved in SARS-CoV-2 WBE modeling and respective modeling techniques.

Study	Location and sampling period	Lowest and highest conc. (solid phase)	Lowest and highest conc. (liquid phase)	Estimated lag period	Statistical correl. and coeff. value*	Modeling technique/algorithm**	Type of contribution (Fig. 2)
Aberi et al. (2021)	Austria (Data collected from databases)	NR	NR	2 – 7 days	NR	Regression models applied to predicting the number of active cases: Linear (LR), Polynomial (PL), K-Nearest Neighbor (KNN), Multilayer Perceptron (MLP), Support Vector Regression (SVR), Generalized Additive Models (GAM), Decision Tree (DT), and Random Forest (RF)	5
Acosta et al. (2022)	Canada June 2020 - May 2021	NR	NR	4 weeks	Pearson's correlation ($r = 0.70$)	NR	4
Ahmed et al. (2021c)	Australia February 2020 - May 2020	NR	1.35E2 - 1.2E4 gc/100mL	NR	NR	NR	2
Ai et al. (2021)	USA July 2020 - January 2021	NR	1E2 – 1E5 gc/L	7 days	Pearson correlation ($r = 0.89$) and Spearman's rank correlation ($r = 0.88$)	Polynomial models	4,5,6
Amereh et al. (2022)	Iran September 2020 - April 2021	NR	1E1 - 1.5E4 gc/L	NR	NR	Monte Carlo simulation to estimate disease prevalence, Linear regression between estimated infected population and confirmed cases ($R^2 = 0.80$, $p < 0.001$)	4
Anneser et al. (2022)	USA March 2020 - March 2021	NR	NR	NR	Spearman's rank correlation ($r = \text{NR}$)	Linear regression ($R^2 = 0.80$), GAM ($R^2 = 0.86$), Poisson ($R^2 = 0.84$), and negative binomial models ($R^2 = 0.15$)	4,5
Bagutti et al. (2022)	Switzerland July 2021 - December 2021	NR	1E2 - 4.13E5 gc/L	14 days	Spearman's rank correlation ($r = 0.9395$)	NR	4
Barrios et al. (2021)	Argentina June 2020 - April 2021	NR	1E-1 - 1E3 gc/L	NR	Spearman rank correlation ($r = 0.812$)	NR	4
Cao and Francis (2021)	USA April 2020 - February 2021	NR	NR	NR	NR	VAR (Vector Autoregression) model	5
Claro et al. (2021)	Brazil June 2020 - April 2021	NR	2.7 - 7.7 \log_{10} gc/L	2 weeks	NR	Monte Carlo simulations to estimate COVID-19 prevalence for each sampling site	4

Study	Location and sampling period	Lowest and highest conc. (solid phase)	Lowest and highest conc. (liquid phase)	Estimated lag period	Statistical correl. and coeff. value*	Modeling technique/algorithm**	Type of contribution (Fig. 2)
de Freitas Bueno et al. (2022)	Brazil January 2021 - January 2022	NR	NR	NR	Spearman's rank correlation ($r = 0.67$)	Monte Carlo statistical model to introduce uncertainty in the virus shedding	4,5
de Sousa et al. (2022)	Brazil January 2021 - August 2021	NR	N1: 2.73 - 3.73 log ₁₀ gc/L; N2: 2.69 - 5.47 log ₁₀ gc/L	NR	NR	Prediction model for infected individuals published by Ahmed et al. (2020a) with Monte Carlo simulations to introduce uncertainties	5
Fernandez-Cassi et al. (2021)	Switzerland February 2020 - April 2020	NR	NR	5.5 days	NR	Incidence estimation by the SEIR model with Gamma distribution to represent virus shedding and time between infection and symptom onset	5
Fitzgerald et al. (2021)	Scotland April 20202 - January 2021	NR	NR	NR	Spearman's rank correlation ($r = 0.91$)	Basic linear mixed model	1,2
Galani et al. (2022)	Greece August 2020 - March 2021	NR	NR	5 - 9 days	Pearson's correlation ($r = 0.947$)	Distributed/fixed lag modeling, linear regression, and artificial neural networks were utilized to build relationships between SARS-CoV-2 RNA load in wastewater and pandemic health indicators	4,5
Gonzalez et al. (2020)	USA March 2020 - August 2020	NR	$10^1 - 10^4$ gc/100mL	NR	NR	NR	2
Hemalatha et al. (2021)	India July 2020 - August 2020	NR	NR	NR	NR	Prediction model for infected individuals published by Ahmed et al. (2020a) and another one by Hellmer et al. (2014)	2,3
Hoar et al. (2022)	USA August 2020 - April 2021	NR	NR	NR	Spearman's rank correlation ($r = 0.81$)	Linear regression ($R^2 = 0.65$)	4,5
Jiang et al. (2022)	USA May 2020 - December 2021	NR	NR	NR	NR	Artificial Neural Network (Best fit with $R^2 = 0.89$)	5
Karthikeyan et al. (2021)	USA July 2020 - October 2020	NR	NR	3 weeks	Pearson's correlation ($r = 0.84$)	ARIMA (Linear Regression model with Autoregressive model)	3,4
Koureas et al. (2021)	Greece October 2020 - April 2021	NR	NR	NR	NR	Linear Regression ($R^2 = 0.9511$) and Random Forest (RF) ($R^2 = 0.9956$)	5
Krivosnakova et al. (2021)	Slovakia September 2020 - March 2021	NR	NR	2 weeks	NR	Regression models to calculate viral concentration: Simple Linear, Double Square Root, and Square Root-Y	2,4

Study	Location and sampling period	Lowest and highest conc. (solid phase)	Lowest and highest conc. (liquid phase)	Estimated lag period	Statistical correl. and coeff. value*	Modeling technique/algorithm**	Type of contribution (Fig. 2)
Kuhn et al. (2022)	USA November 2020 - March 2021	NR	1.6E1 - 7.3E6 gc/L	4 – 10 days	Pearson's correlation and Spearman rank correlation (r = NR)	General Multivariate Linear Regression, multivariate Poisson (best accuracy obtained) and Negative Binomial models	5
Layton et al. (2022)	USA June 2020 - July 2020	NR	2.9 - 5.1 log10 gc/L	NR	Pearson's correlation (r = 0.96)	Monte Carlo simulation to account for the uncertainty in the point estimates for each sampling event	4
Li et al. (2021b)	Australia Used data from seven papers	NR	NR	NR	NR	Three types of data-driven models were applied to a multi-national WBE dataset: multiple linear regression (MLR), artificial neural network (ANN), and adaptive neuro-fuzzy inference system (ANFIS) to predict upcoming new cases	5
Li et al. (2022)	USA June 2020 - September 2021	NR	2.76E3 - 3.86E6 gc/L	7 days	Spearman's rank correlation (r = 0.790)	NR	2,4
Maida et al. (2022)	Italy September 2021 - July 2021	NR	NR	NR	NR	A logistic regression model was calculated to evaluate the association between the active SARS-CoV-2 incidence rates and the probability of positive PCR results of wastewater samples	5
McMahan et al. (2021)	USA May 2020 - August 2020	NR	4.7E3 - 3.3E6 gc/L	NR	NR	Susceptible-Exposed-Infectious-Recovered (SEIR) model to predict the number of infected individuals based on the mass rate (gc/day) of SARS-CoV-2 RNA in WW	5
Nagarkar et al. (2022)	USA May 2020 - November 2020	NR	1E3 – 1E4 gc/L for N1 and N2	NR	Pearson's correlation (r = 0.70)	NR	4
Nourbakhsh et al. (2022)	Canada September 2020 - June 2021	NR	NR	3 – 20 days	NR	Viral transmission is simulated via a standard epidemiological SEIR-like model ("Susceptible - Exposed - Infectious - Recovered") and the fate of SARS-CoV-2 in wastewater using an advection-dispersion-decay model	5
Omori et al. (2021)	USA April 2020 – June 2020	NR	~10 - ~4E2 gc/L	8.4 – 11.6 days	NR	Data fitting using Poisson distribution	5
Peccia et al. (2020)	USA March 2020 - June 2020	NR	1.7E3 - 4.6E5 gc/L	6 – 8 days	NR	Linear regressions were used to estimate the relationship between	4,5

Study	Location and sampling period	Lowest and highest conc. (solid phase)	Lowest and highest conc. (liquid phase)	Estimated lag period	Statistical correl. and coeff. value*	Modeling technique/algorithm**	Type of contribution (Fig. 2)
						SARS-CoV-2 RNA copies per ml results for replicated RNA extractions of each daily sample. Estimation of primary sludge as a potential leading indicator was performed using a distributed lag measurement error time series model	
Petala et al. (2022)	Greece October 2020 - January 2021	NR	NR	NR	NR	Developed a set of parametric equations to estimate the evolution of global virus shedding rate in wastewater	5
Pillay et al. (2021)	South Africa July 2020 - October 2020	NR	0 - 7.12E5 gc/100	NR	NR	Prediction model for infected individuals published by Ahmed et al. (2020a)	5
Proverbio et al. (2022)	Luxembourg NR (Data collected from databases)	NR	NR	NR	NR	Use of Susceptible-Exposed-Infectious-Recovered (SEIR) epidemiological model to the extended Kalman filter – EKF	5
Reynolds et al. (2022)	Ireland June 2020 - August 2021	NR	NR	0 days	Spearman's rank correlation (r = 0.500)	NR	2,6
Rodriguez Rasero et al. (2022)	Spain July 2020 – February 2021	NR	NR	6 days	NR	Data fitting using quasi-Poisson modeling	5
Roka et al. (2021)	Hungary June 2020 - October 2020	NR	~5E3 - ~1E6 gc/L	NR	NR	Data fitting using Linear Regression (Best fit value using a weighted average of viral load against daily new cases, $R^2 = 0.720$ and $p < 0.0001$)	1,4
Saththasivam et al. (2021)	Qatar June 2020 - August 2020	NR	7.889E3 - 5.42E5 gc/L	NR	NR	Conservation principles to estimate the number of infected populations based on measuring RNA concentration	2,5
Scott et al. (2021)	USA August 2020 - December 2020	NR	N1: 22.5 - 5.27E3 gc/100 ; N2: 81.6 - 3.91E4 gc/100	NR	Spearman's rank correlation (r = 0.50)	Multiple Linear Regression, Simple Logistic Regression and Multiple Logistic Regression	1,5
Song et al. (2021)	USA April 2020 – June 2020	NR	~8E0 – 9E5 gc/	NR	NR	NR	1,2,6
Tanimoto et al. (2022)	Japan February 2021 - October 2021	1.5E7 – 2.0E8 gc/L	3.1E7 – 5.5E8 gc/L	NR	NR Solid phase: r = 0.8482, Liquid phase: (r = 0.7803)	Linear Regression	4

Study	Location and sampling period	Lowest and highest conc. (solid phase)	Lowest and highest conc. (liquid phase)	Estimated lag period	Statistical correl. and coeff. value*	Modeling technique/algorithm**	Type of contribution (Fig. 2)
Vallejo et al. (2022)	Spain March 2020 - May 2020	NR	10E3 – 15E4 gc/	NR	NR	Linear regression ($R^2 = 0.8515$), GAM - Generalized Additive with a Cubic Regression Spline ($R^2 = 0.8767$), LOESS Linear ($R^2 = 0.8685$), LOESS Quadratic ($R^2 = 0.8833$)	4,5
Wang et al. (2021)	USA NR	NR	N1: 3.85E5 - 2.55E6 gc/L, N2: 3.79E5 - 2.15E6 gc/L	NR	Pearson's correlation ($r = 0.94$)	Monte Carlo simulations to estimate the number of infected individuals	4
Wu et al. (2022)	USA January 2020 - May 2020	NR	NR	4 – 10 days	Pearson's correlation ($r = \text{NR}$)	Wastewater data was modeled as a convolution of new clinical cases and used Markov Chain Monte Carlo (MCMC) simulation to quantify uncertainty in the shedding model	4,5
Wurtzer et al. (2022)	France March 2020 - June 2021	NR	0 – 1e6 gc/L	3 days	Spearman's rank correlation ($r = \text{NR}$)	Linear regression	4
Xiao et al. (2022)	USA March 2020 - June 2020	NR	NR	6.4 days	NR	Approximate Bayesian computation for estimating delay distribution, convolution to estimate the transfer function model, and Markov chain Monte Carlo (MCMC) simulation to quantify uncertainty in transfer functions	5
Zhao et al. (2022)	USA Sept 2020 - August 2021	N1: 7.14E2 – 7.14E3 gc/L, N2: 1.02E2 – 6.1E3 gc/L	NR	5 weeks	Pearson's correlation (N1: $r = 0.62$ and N2: $r = 0.64$)	Linear regression, ARIMA, SARIMA (Regression Model with Autoregressive Model with Seasonal Pattern), and (4) VAR (Vector Autoregressive model)	4
Zhu et al. (2022)	Japan August 2020 - February 2021	NR	~10 - ~70 gc/	NR	NR	Generalized linear model, Artificial neural network and random forest to predict the cumulative number of cases	4,5

Notes: NR stands for Not Reported;

* The indicated value corresponds to the best fit obtained in the respective study.

** The designated models aim to associate viral concentration signals from wastewater to clinical testing data.

Figure 1. Publication selection process: PRISMA-based flowchart for evidence-based research. 1.5 column image





Figure 2. Types of contributions of the selected studies to SARS-CoV-2 WBE: findings from the analysis of the selected pool of publications. 1 column image

- (1) Operation of wastewater analysis:** quantitative comparisons between concentration methods, RNA isolation kits, and different PCR approaches in terms of viral recovery, sensitivity, and specificity.
- (2) Reporting SARS-CoV-2 detection in wastewater:** description of analytical procedures toward detection of SARS-CoV-2 genomic material in wastewater worldwide.
- (3) Development, modifications and/or optimization of analysis protocols:** detailed descriptions of analytical procedures for concentrating, isolating, detecting, and quantifying SARS-CoV-2 RNA in wastewater samples.
- (4) Building statistical correlations between viral concentration signals and clinical testing data:** development and application of correlations to assess how strongly temporal series of wastewater viral concentrations and disease prevalence are associated.
- (5) WBE modeling, simulation and/or parameter estimation:** details of different modeling techniques and data analysis toward understanding the spread of SARS-CoV-2 in domestic wastewater, and building complex models to predict outbreaks based on viral concentration signals.
- (6) Phylogenetics, genotyping and/or identification of variants of concern (VOC):** descriptions of genetic sequencing to support the hypothesis of VOCs prevalence in domestic wastewater, as well as tracking the same VOCs through the WBE approach.

Figure 3. SARS-CoV-2 WBE: Overall framework for sample analysis and epidemiological modeling. 1.5 column image



Author contributions

Stefano Ciannella: Methodology, Formal analysis, Investigation, Writing - Original Draft, Visualization. **Cristina González-Fernández:** Writing - Original Draft, Writing - Review & Editing, Visualization. **Jenifer Gomez-Pastora:** Conceptualization, Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

Journal Pre-proof


Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof

Graphical abstract



Research highlights:

- COVID-19 community cases can be predicted by wastewater-based epidemiology (WBE);
- WBE correlates SARS-CoV-2 levels in wastewater with COVID-19 clinical cases;
- Six major contribution areas to the development of WBE for COVID-19 were identified;
- Standardization of analytic procedures for SARS-CoV-2 detection is urgently needed;
- Opportunities to improve accuracy in WBE for COVID-19 are emphasized.