

Letter

K_S^0 and $\Lambda(\bar{\Lambda})$ two-particle femtoscopic correlations in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV

The CMS Collaboration ^{*}

CERN, Geneva, Switzerland

ARTICLE INFO

Editor: M. Doser

Keywords:

CMS

Heavy ion

V0

Femtoscopy

Scattering parameters

ABSTRACT

Two-particle correlations are presented for K_S^0 , Λ , and $\bar{\Lambda}$ strange hadrons as a function of relative momentum in lead-lead collisions at a nucleon-nucleon center-of-mass energy of 5.02 TeV. The dataset corresponds to an integrated luminosity of 0.607 nb^{-1} and was collected using the CMS detector at the CERN LHC. These correlations are sensitive to quantum statistics and to final-state interactions between the particles. The source size extracted from the $K_S^0 K_S^0$ correlations is found to decrease from 4.6 to 1.6 fm in going from central to peripheral collisions. Strong interaction scattering parameters (i.e., scattering length and effective range) are determined from the ΛK_S^0 and $\Lambda\Lambda$ (including their charge conjugates) correlations using the Lednický–Lyuboshitz model and are compared to theoretical and other experimental results.

1. Introduction

Two-particle correlations in relative momentum, so-called femtoscopic correlations, arising from relativistic heavy ion collisions provide a powerful tool for studying the spatiotemporal characteristics of the particle emitting sources created in these collisions and the subsequent interactions of the emitted particles [1]. All two-particle correlations involving hadrons are affected by final-state interaction (FSI) effects, and correlations of identical particles are also sensitive to the constraints of quantum statistics (QS). The correlations among the neutral K_S^0 , Λ , and $\bar{\Lambda}$ particles, collectively referred to as V^0 particles, are of special interest. Because of their relatively large mass, femtoscopy based on K_S^0 particles is less affected by resonance decays than is the case with more commonly studied pion and charged kaon pairs [2]. The results from $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ correlation studies can be used to determine the strong-interaction scattering parameters, i.e., the scattering length and the effective range, that are impossible to obtain from currently achievable scattering experiments [3–7]. This information can be used to constrain hadron-hadron correlation models that are used, for example, in modeling the composition of neutron stars [8,9].

Regarding the scattering parameters, of particular interest is establishing whether the interaction between two Λ particles allows for the existence of the H-dibaryon, a bound state with quantum numbers $I = 0$, $J^P = 0^+$, and $S = -2$. In 1977, R. L. Jaffe predicted the existence of such a six-quark (uuddss) state having a mass about 81 MeV below the thresh-

old of twice the Λ mass by considering the strong attraction resulting from color magnetic interactions [10]. Although a double hypernucleus, ${}^6_{\Lambda\Lambda}\text{He}$, was subsequently observed in the NAGARA event from the E313 hybrid emulsion experiment at KEK [11,12], the observed $\Lambda\Lambda$ binding energy was not consistent with the conjectured H-dibaryon [13]. A study of $\Lambda\Lambda$ correlations may provide additional information on whether the baryon-baryon interaction can lead to the formation of the conjectured H-dibaryon.

Recently, the ALICE Collaboration reported on ΛK correlations in lead-lead (PbPb) collisions at a center-of-mass energy per nucleon pair of $\sqrt{s_{NN}} = 2.76$ TeV [14]. According to their findings, the strong force is repulsive in ΛK^+ interactions, yet attractive in ΛK^- interactions. For the ΛK_S^0 pairs, the uncertainty of the ALICE results does not permit a definite conclusion on whether the associated strong interaction is repulsive or attractive. A more precise measurement of ΛK_S^0 correlations should improve our understanding of the strong interaction in baryon-meson systems.

This Letter presents $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ femtoscopic correlations as a function of relative momentum in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, using data recorded by the CMS experiment during the 2018 LHC run. The $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ correlations are measured in an integrated centrality range 0–80%, where centrality refers to the percentage of the total inelastic hadronic nucleus-nucleus cross section [15], and 0% corresponds to the maximum overlap of the colliding nuclei. This is the first measurement of the $\Lambda\Lambda$ femtoscopic correlation in PbPb colli-

^{*} E-mail address: cms-publication-committee-chair@cern.ch.

sions at the LHC. In addition, the $K_S^0 K_S^0$ correlations are measured in six centrality intervals in the range of 0–60%. The source size and strong interaction parameters are determined using the Lednický–Lyuboshitz (LL) model [16]. Unless otherwise indicated, all measurements include the charge conjugate states, so ΛK_S^0 and $\Lambda\Lambda$ include $\bar{\Lambda} K_S^0$ and $\bar{\Lambda}\bar{\Lambda}$, respectively. Tabulated results are provided in the HEPData record for this analysis [17].

2. Experimental setup and data sample

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume there is a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter, and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. The silicon pixel detector [18] is composed of 1856 silicon pixel modules distributed in four 54 cm long barrel layers at radii of 2.9–16.0 cm plus three pairs of endcap disks covering radii of 4.5–16.1 cm at longitudinal distances of 31–51 cm from the origin. The 15148 silicon strip modules are arranged in 10 barrel layers at radii of 20–116 cm plus 3 pairs of small and 9 pairs of large endcap disk layers. Charged particles of pseudorapidity $|\eta| < 3$ are reconstructed with the combined system. For particles with transverse momentum of $1 < p_T < 10$ GeV, the track resolutions are typically 1.5% in p_T and 20–75 μm in the transverse impact parameter [19]. The barrel and endcap detectors are extended to the forward region with two calorimeters which use steel as the absorber and quartz fibers as the sensitive material. These hadron forward (HF) calorimeters are located 11.2 m from the interaction region, one on each side, and provide coverage in the range $3.0 < |\eta| < 5.2$. These detectors are segmented into multiple 0.175×0.175 ($\Delta\eta \times \Delta\phi$) “towers”, where ϕ is azimuthal angle in radians. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4 μs [20]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [21]. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [22].

With an integrated luminosity of 0.607 nb^{-1} [23,24], this analysis uses 4.27×10^9 minimum bias events that are triggered by requiring signals above the readout threshold of 3 GeV in each of the HF calorimeters [21]. Background events due to beam-gas interactions and nonhadronic collisions are filtered offline by applying the procedure described in Ref. [25]. The events used in this analysis are required to have at least one primary interaction vertex determined using two or more tracks [26] within a distance of 15 cm from the center of the nominal interaction point along the beam axis and to have at least two calorimeter towers in each HF detector with energy deposits of more than 4 GeV per tower. The shapes of the clusters in the pixel detector are required to be compatible with those expected in PbPb collisions in order to suppress the contamination from events with multiple collisions [27]. The combined trigger and offline selection efficiency for inelastic events is greater than 95%. The event centrality is obtained from the transverse energy deposited in both HF calorimeters, using the methodology described in Ref. [28]. The analysis makes use of a minimum bias Monte Carlo PbPb sample, based on the HYDJET 1.9 [29] event generator with a full detector simulation using GEANT4 [30].

3. Reconstruction of K_S^0 and Λ candidates

The K_S^0 and Λ candidates, denoted as V^0 candidates, used in this study are reconstructed as in previous CMS analyses [31–33]. The V^0 candidates are found by combining oppositely charged tracks that pass criteria based on the “loose” selection discussed in Ref. [26]. The charged tracks are assumed to be $\pi^+\pi^-$ in K_S^0 reconstruction and $\pi^-\bar{p}$ in Λ reconstruction. For the latter, the higher momentum track is assumed to be a proton since the proton carries nearly all of the momentum in the Λ decay. Each of the oppositely charged tracks must have hits in at least three layers of the silicon tracker, and both tracks must have transverse and longitudinal impact parameter significances (defined as the parameter value divided by its uncertainty) with respect to the primary vertex greater than 1. The two tracks are fitted to a common vertex and the χ^2 per degree of freedom (dof) from the fit must be less than 7. The distance of closest approach between the two tracks is required to be less than 1 cm. As a consequence of the long lifetime of K_S^0 and Λ particles, the significance of the V^0 decay length, which is the three-dimensional distance between the primary and V^0 vertices divided by its uncertainty, is required to be greater than 2.5 to reduce combinatorial background contributions. To remove K_S^0 candidates misidentified as Λ particles and vice versa, the Λ (K_S^0) candidates must have a corresponding $\pi^+\pi^-$ ($p\pi^-$) mass more than 14 (7) MeV (corresponding to approximately 3 times the average resolution) away from the world-average value [34] of the K_S^0 (Λ) mass. The angle θ between the V^0 momentum vector and the vector connecting the primary and V^0 vertices is required to satisfy $\cos \theta > 0.999$. This reduces the contribution from nuclear interactions, random combinations of tracks, and Λ particles originating from weak decays of Ξ and Ω particles.

Further selection of V^0 candidates is performed with a boosted decision tree (BDT) [35]. The selection is optimized separately for K_S^0 and Λ candidates. The discriminating variables include: the collision centrality, the V^0 candidate p_T and rapidity (y), the distance of closest approach of the track pair, the three-dimensional decay length and significance, $\cos \theta$, and the V^0 vertex fit χ^2 . The included variables related to the V^0 daughters are p_T , uncertainty in p_T , η , the number of hits in the silicon tracker, the number of pixel detector layers with hits, and the transverse and longitudinal impact parameter significances with respect to the primary vertex. The BDT training is performed using the simulated minimum bias sample separated into the signal and background subsamples using the generator-level information. The K_S^0 mesons are selected with $1 < p_T < 8.5$ GeV and $|y| < 1$, while the Λ baryons are required to have $1.8 < p_T < 8.5$ GeV and $|y| < 1$. The minimum p_T and maximum y requirements are used to reduce background while the maximum p_T requirement is to reduce contributions from jets. The combined V^0 reconstruction and selection efficiencies are strongly dependent on the centrality of the event and the p_T of the V^0 . Integrating over the selected p_T ranges, the combined efficiencies from the most central to peripheral PbPb collisions are 1–3% for K_S^0 and 1–2% for Λ . The V^0 reconstruction algorithm does not prevent a track from being used for more than one V^0 . While this is normally an infrequent occurrence, selecting pairs of V^0 particles close together in phase space makes it a significant contribution. To resolve this problem, for each correlation measurement, a check of each pair of V^0 candidates is performed and if two V^0 candidates are found to share one or both daughter tracks, one of the V^0 candidates is randomly selected to be removed from the event.

Fits to the invariant mass spectrum are performed using a sum of three Gaussian functions with a common mean to describe the signal distribution and a fourth-order polynomial to describe the background. These empirical functions were chosen to provide a good description of the data. Peak and sideband invariant mass regions are defined to select events dominated by signal and background, respectively. Defining σ as the average resolution based on the Gaussian sum, the peak regions are selected to be within $\pm 2\sigma$ from the nominal V^0 mass and are given by $486 < M(\pi^+\pi^-) < 509$ MeV and $1111.5 < M(p\pi^-) < 1120.4$ MeV




Fig. 1. The invariant mass of K_S^0 (left) and Λ (right), and their corresponding fits in the 0–80% centrality range. The circles are the data, and the fit is shown with a solid (red) line for the total fit, and a dashed (green) line for the background fit. The vertical dashed-dotted (pink) lines indicate the peak region and the vertical dashed (blue) lines indicate the sideband regions.

for K_S^0 and Λ candidates, respectively. The sideband regions are selected to be more than 4σ from the nominal V^0 mass and are given by $23.5 < |M(\pi^+\pi^-) - 497.5| < 62.5$ MeV for K_S^0 candidates and $1080 < M(p\pi^-) < 1107.5$ MeV together with $1124.2 < M(p\pi^-) < 1160$ MeV for Λ candidates. Examples of invariant mass distributions for $\pi^+\pi^-$ and $p\pi^-$ pairs, and their corresponding fits in the 0–80% centrality range, are shown in Fig. 1.

4. Analysis method

The two-particle correlation is constructed as

$$C^{\text{obs}}(q_{\text{inv}}) = \mathcal{N} \frac{A^{\text{obs}}(q_{\text{inv}})}{B^{\text{obs}}(q_{\text{inv}})}, \quad (1)$$

where $C^{\text{obs}}(q_{\text{inv}})$ is the observed normalized pair yield as a function of the invariant relative momentum q_{inv} in the pair rest frame, defined as [1]

$$q_{\text{inv}} = \sqrt{-Q^\mu Q_\mu}, \quad (2)$$

$$Q^\mu = (k_1 - k_2)^\mu - \frac{(k_1 - k_2)^\mu P_\mu}{P^\mu P_\mu} P^\mu,$$

where $P = k_1 + k_2$, and k_1 and k_2 are the four momenta of the V^0 particles. Note that for two particles of the same mass, the second term of Q^μ is zero.

The distribution $A^{\text{obs}}(q_{\text{inv}})$ is the signal distribution, which includes the femtoscopic and nonfemtoscopic correlations formed by pairing the selected V^0 particles from a given event. The reference distribution $B^{\text{obs}}(q_{\text{inv}})$ is constructed by mixing the V^0 particles from different events, as described in Ref. [36]. In this procedure, the V^0 particle from one event is paired with V^0 particles from 30 different events. To ensure that the 30 events used in the mixing are similar to the signal events, the centrality and primary vertex of each mixed event must be within 5% and 2 cm, respectively, of those in the corresponding signal event. By dividing $A^{\text{obs}}(q_{\text{inv}})$ by $B^{\text{obs}}(q_{\text{inv}})$, CMS detector and reconstruction artifacts are strongly suppressed. The normalization factor $\mathcal{N}$ is the ratio of the number of pairs in the reference distribution to that in the signal distribution. Because of the background in the peak region of the invariant mass distributions, the measured signal distribution ($A^{\text{obs}}(q_{\text{inv}})$) contains contributions from signal-signal ($A^{\text{ss}}(q_{\text{inv}})$), signal-background ($A^{\text{sb}}(q_{\text{inv}})$), and background-background ($A^{\text{bb}}(q_{\text{inv}})$) correlations. The measured $A^{\text{obs}}(q_{\text{inv}})$ distribution can be written as

$$A^{\text{obs}}(q_{\text{inv}}) = f^{\text{ss}} A^{\text{ss}}(q_{\text{inv}}) + f^{\text{sb}} A^{\text{sb}}(q_{\text{inv}}) + f^{\text{bb}} A^{\text{bb}}(q_{\text{inv}}). \quad (3)$$

Table 1

The values of the f^{ss} , f^{sb} , and f^{bb} parameters are based on the number of counts in the peak and sideband regions. The values are given for the $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ pairs in the integrated 0–80% centrality range. The uncertainties are statistical.

Parameter	$K_S^0 K_S^0$	ΛK_S^0	$\Lambda\Lambda$
f^{ss}	0.9569 ± 0.0006	0.9332 ± 0.0006	0.9054 ± 0.0020
f^{sb}	0.0426 ± 0.0005	0.0655 ± 0.0006	0.0922 ± 0.0018
f^{bb}	0.0005 ± 0.0001	0.0013 ± 0.0002	0.0024 ± 0.0002

The distributions $A^{\text{sb}}(q_{\text{inv}})$ and $A^{\text{bb}}(q_{\text{inv}})$ are obtained from the peak-sideband and sideband-sideband combinations, respectively. The small amount of background (signal) contamination in the signal (sideband) region has a negligible effect on the shape of $A^{\text{sb}}(q_{\text{inv}})$ ($A^{\text{bb}}(q_{\text{inv}})$). All distributions, $A^{\text{ss}}(q_{\text{inv}})$, $A^{\text{sb}}(q_{\text{inv}})$, and $A^{\text{bb}}(q_{\text{inv}})$ are normalized to unity. The parameters, f^{ss} , f^{sb} , and f^{bb} are the signal-signal, signal-background, and background-background fractions, extracted using an invariant mass fit based on combinatorial analyses with

$$f^{\text{ss}} = \frac{\binom{s}{2}}{\binom{s+b}{2}}, \quad (4)$$

$$f^{\text{bb}} = \frac{\binom{b}{2}}{\binom{s+b}{2}}, \quad \text{and}$$

$$f^{\text{sb}} = 1 - f^{\text{ss}} - f^{\text{bb}},$$

where $\binom{n}{2} = \frac{n(n-1)}{2}$ is the binomial coefficient, which returns the number of ways that a pair can be chosen from n objects. The quantities s and b in the binomial coefficients are the number of signal and background particles, respectively, obtained by integrating the appropriate function from the fit to the invariant mass distribution. The resulting values of the f^{ss} , f^{sb} , and f^{bb} parameters are presented in Table 1.

Once we have all the distributions ($A^{\text{obs}}(q_{\text{inv}})$, $A^{\text{sb}}(q_{\text{inv}})$, and $A^{\text{bb}}(q_{\text{inv}})$) and the parameters (f^{ss} , f^{sb} , and f^{bb}), the $A^{\text{ss}}(q_{\text{inv}})$ distribution can be extracted using Eq. (3), with

$$A^{\text{ss}}(q_{\text{inv}}) = \left[A^{\text{obs}}(q_{\text{inv}}) - f^{\text{sb}} A^{\text{sb}}(q_{\text{inv}}) - f^{\text{bb}} A^{\text{bb}}(q_{\text{inv}}) \right] / f^{\text{ss}}. \quad (5)$$

The same procedure is followed for the reference distribution. After extracting the $A^{\text{ss}}(q_{\text{inv}})$ and $B^{\text{ss}}(q_{\text{inv}})$ distributions, the correlation distribution is calculated as

$$C(q_{\text{inv}}) = \mathcal{N} \frac{A^{\text{ss}}(q_{\text{inv}})}{B^{\text{ss}}(q_{\text{inv}})}. \quad (6)$$

The measured fractions, reported in Table 1, confirm the high purity of the V^0 pairs suggested by Fig. 1. The effect of the non- V^0 background

correction on the measured physics parameters is small compared to either the statistical or systematic uncertainties. While the $C(q_{\text{inv}})$ distribution is corrected for non- V^0 backgrounds as well as nonuniform efficiency and acceptance, it still includes nonfemtoscopic background correlations, such as those associated with elliptic flow [37], minijets [2], resonance decays [2], and energy-momentum conservation [38]. The nonfemtoscopic background contribution is modeled using an empirically determined double Gaussian function

$$\Omega(q_{\text{inv}}) = N \left(1 + \alpha_1 e^{-q_{\text{inv}}^2 R_1^2} \right) \left(1 - \alpha_2 e^{-q_{\text{inv}}^2 R_2^2} \right), \quad (7)$$

where N , α_1 , α_2 , R_1 , and R_2 are fit parameters. This function was selected for its reproduction of the distributions in both real data at high q_{inv} and simulated data that do not include the correlations being measured.

Fits are performed to the $C(q_{\text{inv}})$ distributions to extract the source size and strong interaction scattering parameters. As the V^0 particles are neutral, the Coulomb interaction is absent. However, the correlations are sensitive to QS and FSI effects, with s -wave interactions assumed to dominate for the small relative momenta of the particle pairs analyzed. The correlation distribution for all pairs ($K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$) is interpreted in the LL model. This model relates the two-particle correlation function to the source size and also takes into account FSI effects [16]. The general correlation function is

$$C_{\text{total}}(q_{\text{inv}}) = \left[1 + \lambda \left(C_{\text{QS}}(q_{\text{inv}}) + C_{\text{FSI}}(q_{\text{inv}}) \right) \right] \Omega(q_{\text{inv}}), \quad (8)$$

where $C_{\text{QS}}(q_{\text{inv}})$ is the QS function and $C_{\text{FSI}}(q_{\text{inv}})$ is the FSI function. The parameter λ is referred to as the correlation strength. In the absence of FSI effects, λ equals unity for a perfectly incoherent Gaussian source. However, effects such as resonance decay, minijets, coherent sources, non-Gaussian source contributions, and FSI between particles can lead to deviations of the λ parameter from unity.

Neglecting CP violation, the $K_S^0 K_S^0$ system can be written as

$$|K_S^0 K_S^0\rangle = \frac{1}{2} \left(|K^0 K^0\rangle + |K^0 \bar{K}^0\rangle + |\bar{K}^0 K^0\rangle + |\bar{K}^0 \bar{K}^0\rangle \right). \quad (9)$$

It can be shown [16,39] that the resulting correlations follow Bose-Einstein quantum statistics, with

$$C_{\text{QS}}(q_{\text{inv}}) = e^{(-q_{\text{inv}}^2 R_{\text{inv}}^2)}, \quad (10)$$

where the source radius R_{inv} reflects the size of the region over which the particles are emitted.

The FSI for the $K_S^0 K_S^0$ correlations is modeled by [16,39]

$$C_{\text{FSI}}(q_{\text{inv}}) = \frac{1}{2} \left[\left| \frac{f(k)}{R_{\text{inv}}} \right|^2 + \frac{4\Re f(k)}{\sqrt{\pi} R_{\text{inv}}} F_1(q_{\text{inv}} R_{\text{inv}}) - \frac{2\Im f(k)}{R_{\text{inv}}} F_2(q_{\text{inv}} R_{\text{inv}}) \right], \quad (11)$$

where

$$k = q_{\text{inv}}/2, \quad (12)$$

$$F_1(z) = \frac{1}{z} e^{-z^2} \int_0^z e^{x^2} dx, \text{ and}$$

$$F_2(z) = \frac{1 - e^{-z^2}}{z}.$$

The function $f(k)$ is the $K^0 \bar{K}^0$ s -wave scattering amplitude, with real and imaginary parts $\Re f(k)$ and $\Im f(k)$, respectively. This amplitude is dominated by the near-threshold s -wave isoscalar resonance $f_0(980)$ and the s -wave isovector resonance $a_0(980)$, with the total scattering amplitude given by an average of these contributions: $f(k) = (f_{f_0(980)}(k) + f_{a_0(980)}(k))/2$. The individual resonance amplitudes depend on the res-

onance mass m_r , with $r = f_0(980)$ or $a_0(980)$, the kaon mass m_K , and the resonance couplings γ_r (γ_r') to the $K^0 \bar{K}^0$ ($\pi\pi$ for $f_0(980)$ and $\pi^0\eta$ for $a_0(980)$) channels. Then, $f_r(k) = \gamma_r / [m_r^2 - \zeta - i\gamma_r k - i\gamma_r' k']$, where $\zeta = 4(m_K^2 + k^2)$ and k' denotes the momentum in the second ($\pi\pi$ or $\pi^0\eta$) decay channel with the corresponding partial width $\Gamma' = \gamma_r' k' / m_r$ (more details can be found in Ref. [39]). The scattering amplitude is calculated using the resonance mass and the coupling parameters from Refs. [40–43], taken from row C of Table 1 of Ref. [39].

For the correlations involving Λ baryons, the $C_{\text{QS}}(q_{\text{inv}})$ and $C_{\text{FSI}}(q_{\text{inv}})$ functions are [16]

$$C_{\text{QS}}(q_{\text{inv}}) = \alpha e^{(-q_{\text{inv}}^2 R_{\text{inv}}^2)}, \text{ and}$$

$$C_{\text{FSI}}(q_{\text{inv}}) = (1 + \alpha) \left[\frac{1}{2} \frac{|f(k)|^2}{R_{\text{inv}}^2} \left(1 - \frac{1}{2\sqrt{\pi}} \frac{d_0}{R_{\text{inv}}} \right) + \frac{2\Re f(k)}{\sqrt{\pi} R_{\text{inv}}} F_1(q_{\text{inv}} R_{\text{inv}}) - \frac{\Im f(k)}{R_{\text{inv}}} F_2(q_{\text{inv}} R_{\text{inv}}) \right], \quad (13)$$

where $\alpha = -1/2$ for $\Lambda\Lambda$ correlations for two identical fermions and $\alpha = 0$ for ΛK_S^0 correlations as there are no QS effects for nonidentical particles [16]. The scattering amplitude $f(k)$ is parameterized by a complex scattering length (f_0) and an effective range (d_0) with $f(k) = [1/f_0 + d_0 k^2/2 - ik]^{-1}$ [16]. The imaginary part of f_0 is responsible for inelastic processes (annihilation). For an attractive interaction that is not strong enough to produce a bound state, the real part of f_0 is positive, while a repulsive interaction corresponds to a negative $\Re f_0$ of the order of the range of the repulsive potential. In the presence of a bound state, $\Re f_0$ is also negative, but with a much larger magnitude. The femtoscopic sign convention and notation for the scattering length differ from those used in nuclear physics, where the corresponding scattering length $a_0 = -f_0$. As the ΛK_S^0 and $\Lambda\Lambda$ correlations each have only one spin state that contributes to the s -wave scattering, Eq. (13) suffices to describe the FSI effects.

Fits to the correlation distribution of all the pairs were performed using Eq. (8) with the nonfemtoscopic background parameters (N , α_1 , α_2 , R_1 , and R_2) treated as free parameters. For $K_S^0 K_S^0$ correlations, the parameters of interest are R_{inv} and λ , with the scattering amplitude based on previous measurements [40–43]. The ΛK_S^0 and $\Lambda\Lambda$ correlations include additional parameters: d_0 , $\Re f_0$, and $\Im f_0$. The $\Im f_0$ term for $\Lambda\Lambda$ correlations is set to zero since there are no baryon-baryon annihilation processes.

Histograms of the correlation distributions are generated in the range $0 < q_{\text{inv}} < 6$ GeV with 0.02 GeV wide bins for the $K_S^0 K_S^0$ and ΛK_S^0 correlations and 0.04 GeV wide bins for the $\Lambda\Lambda$ correlations. The fits exclude the first q_{inv} bin to avoid a potential bias from the method used to address cases where V^0 candidates share daughter tracks. Studies using simulated events indicate that only this first bin is affected by this remediation. Least-square fits are performed to the experimental data with the uncertainties in the fit parameters calculated using the MINOS technique [44]. Examples of correlation measurements and their fits and corresponding χ^2/dof values, are presented in Figs. 2 and 3. The $K_S^0 K_S^0$ correlations, shown in Fig. 2, are independently fitted for each of the six centrality bins with $0 < k_T < 2$ GeV, where $k_T \equiv |\vec{p}_{T,1} + \vec{p}_{T,2}|/2$ is the average transverse momentum of the particle pair. The ΛK_S^0 (left) and $\Lambda\Lambda$ (right) correlations, shown in Fig. 3, involve fewer events and, therefore, only a single fit is performed for each, with the data integrated over the centrality range 0–80% and with no restriction on k_T .

Using events simulated with the HYDJET and HIJING 1.3 [45] event generators, we find that 85% and 39% of the selected Λ baryons are directly produced, respectively. Electromagnetic Σ^0 baryon decays account for 8% and 13%, respectively, while weak decays of Ξ/Ω baryons account for 7% and 48% of the observed yields. Given the wide variation in the feed-down contribution and the lack of experimental mea-




Fig. 2. The correlation distributions and fits for $K_S^0 K_S^0$ pairs in different centrality ranges, starting from 0–10% centrality to 50–60% centrality, with $0 < k_T < 2$ GeV. In each plot, the red circles are the data, the blue solid line is the fit using Eq. (8), and the green dashed line is the nonfemtoscopic background from Eq. (7). The χ^2 and dof values are for the full q_{inv} range. The insert plots show the data and the fit for the $q_{\text{inv}} < 0.4$ GeV region, with the χ^2 and number of bins evaluated in that region.




Fig. 3. The correlation distributions and fits for ΛK_S^0 (left) and $\Lambda\Lambda$ (right) pairs with 0–80% centrality and no restriction on k_T . In each plot, the red circles are the data, the blue solid line is the fit using Eq. (8), and the green dashed line is the nonfemtoscopic background from Eq. (7). The χ^2 and dof values are for the full q_{inv} range. The insert plots show the data and the fit for the $q_{\text{inv}} < 0.4$ GeV region, with the χ^2 and number of bins evaluated in that region.

measurements for these contributions, our results involving Λ baryons are presented without being corrected for potential, and possibly significant, feed-down effects and residual correlations.

5. Systematic uncertainties

The systematic uncertainties for the fit parameters are based on the changes found in the parameter values after individually varying each of the analysis criteria, as discussed below. In cases with more than one variation for a single source, the maximum deviation from the nominal value is used. The total systematic uncertainty is obtained by adding the uncertainties from each source in quadrature. The BDT discriminant is varied so as to adjust the signal-to-background ratio, with the signal yield changing by $\pm 15\%$ in the process. The nominal method to account for V^0 candidates sharing daughter tracks is to remove one of the V^0 candidates at random, which is then not used by any pair.

Two alternative approaches are used, one in which both V^0 candidates are removed and another in which, for events with multiple V^0 candidate pair combinations, only the pairs in which the two particles share a daughter are removed. The systematic uncertainties related to V^0 signal and background modeling are investigated by varying the background shape from a fourth- to a third-order polynomial and the signal shape from a sum of three Gaussian functions to a sum of two or four Gaussian functions. An alternative nonfemtoscopic background function $\Omega(q_{\text{inv}}) = N(1 + Be^{-|q_{\text{inv}}/\sigma|^2})(1 + \epsilon q_{\text{inv}})$ is used to assess the uncertainty associated with the choice of the nonfemtoscopic background function. The selection requirements used to construct the mixed event sample are varied to require centrality matching of 3% instead of the nominal 5% and the primary vertex position matching with 1 cm instead of the nominal 2 cm. The peak region requirement is changed from $< 2.0\sigma$ to $< 1.5\sigma$ and $< 2.5\sigma$ and the sideband region selection from $> 4.0\sigma$ to $> 3.5\sigma$ and $> 4.5\sigma$. The upper limit of the correlation distributions fit range is

Table 2

Summary of absolute systematic uncertainties in $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ correlation measurements. The values for R_{inv} , d_0 , $\Re f_0$, and $\Im f_0$ are in fm.

Uncertainty source	$K_S^0 K_S^0$		ΛK_S^0					$\Lambda\Lambda$			
	R_{inv}	λ	R_{inv}	d_0	$\Re f_0$	$\Im f_0$	λ	R_{inv}	d_0	$\Re f_0$	λ
BDT cut	0.06–0.19	0.01–0.05	0.19	0.75	0.10	0.07	0.03	0.06	0.43	0.05	0.31
Duplicate V^0 removal	0.07–0.35	0.01–0.05	0.35	0.92	0.10	0.19	0.11	0.01	1.14	0.05	0.14
Mass fit function	0.00–0.03	0.00–0.02	0.09	0.05	0.01	0.03	0.03	0.02	0.04	0.01	0.02
Nonfemtoscopic func.	0.01–0.13	0.01–0.13	0.02	0.17	0.05	0.07	0.03	0.02	1.02	0.14	0.93
Reference sample	0.02–0.09	0.01–0.06	0.22	0.48	0.12	0.12	0.03	0.10	1.12	0.20	0.76
Peak region	0.02–0.17	0.00–0.07	0.43	0.10	0.05	0.17	0.08	0.22	1.21	0.08	0.35
Sideband region	0.01–0.03	0.00–0.01	0.02	0.02	0.01	0.01	0.00	0.01	0.03	0.01	0.04
Fitting range	0.01–0.32	0.00–0.11	0.20	0.18	0.03	0.08	0.04	0.04	1.79	0.20	0.60
Momentum resolution	0.02–0.07	0.02–0.08	0.41	0.00	0.03	0.09	0.08	0.01	0.08	0.01	0.02
$f_0(980)/a_0(980)$ param.	0.11–0.16	0.03–0.05	—	—	—	—	—	—	—	—	—
Total uncertainty	0.30–0.51	0.10–0.18	0.78	1.31	0.20	0.32	0.17	0.25	2.91	0.33	1.43

changed to 5 and 7 GeV, and the lower limit is changed to include the first bin. The possible influence of momentum resolution on the calculated parameters has been studied following the procedure described in Ref. [46], with an associated systematic uncertainty quoted in Table 2. For the $K_S^0 K_S^0$ correlations, an additional systematic uncertainty is found by varying the mass and coupling parameters for the $f_0(980)$ and $a_0(980)$ resonances based on values found in rows A, B, and D of Table 1 of Ref. [39]. The systematic uncertainties are summarized in Table 2.

Dividing the signal distribution by the reference distribution is expected to correct for efficiency-dependent effects in the correlations. A variety of checks were performed with simulated samples to confirm this assumption. It was verified that the reconstruction efficiency of a pair of V^0 particles is well described by the product of the efficiencies for each V^0 , so the efficiency of a pair of V^0 particles from mixed events should be the same as from the same event. It was also verified that the efficiency versus q_{inv} is the same for both the signal and reference distributions. Therefore, no efficiency correction is applied, and no systematic uncertainty is assessed for the V^0 reconstruction efficiency. For the centrality ranges used in this analysis, the effect of the centrality calibration and resolution is negligible [47].

6. Results

The size of the particle emitting source R_{inv} and the λ parameter extracted from the $K_S^0 K_S^0$ correlations for $0 < k_T < 2$ GeV are shown as a function of centrality in Fig. 4. It is observed that the R_{inv} value decreases from central to peripheral events, as expected from a simple geometric picture of the collisions. Over the full centrality range of 0–80%, $R_{\text{inv}} = 3.40 \pm 0.11$ (stat) ± 0.37 (syst) fm. The transverse mass can be calculated as $m_T = \sqrt{(m_{\text{inv}}/2)^2 + k_T^2}$, where m_{inv} is the invariant mass of the two-particle system [14]. The average $\langle m_T \rangle$ is evaluated from the transverse mass distribution using two-particle pairs with $q_{\text{inv}} < 0.4$ GeV, accounting for background using the binomial analysis as done for the q_{inv} distributions, and is found to be 1.50 GeV and is nearly independent of centrality. Fig. 5 shows CMS and ALICE results for R_{inv} and λ as a function of m_T in three centrality bins. Extrapolations of the m_T dependence found for the ALICE results at $\sqrt{s_{\text{NN}}} = 2.76$ TeV [46] to the m_T values of the CMS results at $\sqrt{s_{\text{NN}}} = 5.02$ TeV show consistent results for the two experiments. The λ parameter is seen to decrease from about 0.55 to 0.37 as the collisions become more peripheral. Possible explanations of this decrease include having an increased contribution of minijets and having a source function that becomes increasingly non-Gaussian as the collisions become more peripheral.

Table 3 includes the extracted R_{inv} and λ parameters as well as $\langle m_T \rangle$ for $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ combinations in the 0–80% centrality range. A significant decrease is seen in R_{inv} as the $\langle m_T \rangle$ increases. Qualitatively similar results have been found, both for a given pair type in bins of m_T

Table 3

Extracted values of the R_{inv} , $\Re f_0$, $\Im f_0$, d_0 , λ , and $\langle m_T \rangle$ parameters from the $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ combinations in the 0–80% centrality range. The first and second uncertainties are statistical and systematic, respectively.

Parameter	$K_S^0 K_S^0$	ΛK_S^0	$\Lambda\Lambda$
R_{inv} (fm)	$3.40 \pm 0.11 \pm 0.37$	$2.1^{+1.4}_{-0.5} \pm 0.8$	$1.3^{+0.4}_{-0.2} \pm 0.3$
$\Re f_0$ (fm)	—	$-0.76^{+0.29}_{-0.19} \pm 0.20$	$0.74^{+0.59}_{-0.16} \pm 0.33$
$\Im f_0$ (fm)	—	$-0.07^{+0.48}_{-0.11} \pm 0.32$	—
d_0 (fm)	—	$2.3^{+0.7}_{-0.5} \pm 1.3$	$4.2^{+5.7}_{-2.1} \pm 2.9$
λ	$0.43 \pm 0.03 \pm 0.13$	$0.34^{+0.41}_{-0.12} \pm 0.17$	$1.5^{+1.2}_{-1.1} \pm 1.4$
$\langle m_T \rangle$ (GeV)	1.50	2.09	2.60

and when comparing multiple pair types [1]. Because of the different minimum p_T requirements for K_S^0 and Λ particles, the variation in $\langle m_T \rangle$ includes both $\langle p_T \rangle$ and particle mass differences. The anticorrelation of R_{inv} and $\langle m_T \rangle$ has been interpreted as indicating the presence of an expanding source [1]. Although the λ parameter obtained from the $\Lambda\Lambda$ correlation is consistent, within statistical and systematic uncertainties, with the absence of a femtoscopic correlation, fitting $C(q_{\text{inv}})$ under the assumption of $\lambda = 0$ leads to a χ^2 of 32 with 9 bins in the range $0.04 < q_{\text{inv}} < 0.4$ GeV, compared to a χ^2 of 5, with the same number of bins, when the femtoscopic correlation is taken to be present (shown in the right plot of Fig. 3).

Table 3 also includes the strong interaction scattering parameters d_0 , $\Re f_0$, and $\Im f_0$ obtained from the ΛK_S^0 and $\Lambda\Lambda$ correlations. Fig. 6 shows d_0 and $\Im f_0$ versus $\Re f_0$ in the left and right panels, respectively, along with comparisons to previous results. The current CMS results are shown as red stars and squares for ΛK_S^0 and $\Lambda\Lambda$, respectively. The boxes and lines indicate one-dimensional statistical and systematic uncertainties, respectively. For the CMS results, a two-dimensional contour is also plotted to show where the χ^2 value changes by one unit with respect to its minimum value. This corresponds to a 39% confidence level, including only the statistical uncertainties.

The negative value of $\Re f_0$ observed for the ΛK_S^0 correlations, combined with its relatively small magnitude, suggests a repulsive ΛK_S^0 interaction. The uncertainty associated with the $\Im f_0$ value for the ΛK_S^0 correlations prevents any claim concerning possible inelastic processes. The measured ΛK_S^0 values of $\Re f_0$, $\Im f_0$, and d_0 differ by about 2, 1, and 1 standard deviations, respectively, from the reported ALICE values (teal diamonds) [14], which are obtained from PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV. In contrast to this analysis, the ALICE result is corrected for the effect on the correlations caused by feed-down, which may be a source of the differences in the measured values.

The positive $\Re f_0$ value obtained for the $\Lambda\Lambda$ correlations suggests an attractive interaction that is not strong enough to produce a bound state such as the H-dibaryon [48,49]. This result disagrees with the




Fig. 4. The R_{inv} (left) and λ parameter (right) as a function of centrality. For each data point, the lines and the shaded areas indicate the statistical and systematic uncertainties, respectively.




Fig. 5. The R_{inv} (left) and λ (right) parameters as a function of $\langle m_T \rangle$ for the three centrality bins (0–10%, 10–30%, and 30–50%) from this analysis (dark filled symbols) and from the ALICE experiment (light open symbols) [46]. For each data point, the lines and the shaded areas indicate the statistical and systematic uncertainties, respectively.




Fig. 6. The measured values of d_0 versus $\Re f_0$ (left) and $\Im f_0$ versus $\Re f_0$ (right) from this analysis along with other measurements and predictions as described in the text. For each data point, the lines and the boxes indicate the one-dimensional statistical and systematic uncertainties, respectively. For the CMS points, the dotted lines show two-dimensional contours corresponding to a 39% confidence level with just the statistical uncertainties included.

finding from the STAR Collaboration in gold-gold (AuAu) collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV (blue circle). The negative $\Re f_0$ value of -1.10 ± 0.37 (stat) $^{+0.68}_{-0.08}$ (syst) fm found by STAR, combined with its magnitude, implies a repulsive interaction. It is noted, however, that a theoretical study of the STAR data which considers collective flow and feed-down effects (shown as a shaded region at $d_0 \approx 5$ fm, $\Re f_0 \approx 0.9$ fm) suggests that these data are consistent with the $\Lambda\Lambda$ interaction being attractive [48]. An analysis by the ALICE Collaboration of the $\Lambda\Lambda$ scattering

parameters, using $\Lambda\Lambda$ correlations from pp collisions at $\sqrt{s} = 7$ and 13 TeV, as well as pPb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, excludes certain regions of the d_0 and $\Re f_0$ parameter space, while also suggesting an attractive interaction [50]. Our measured values fall within their allowed regions and provide further evidence for an attractive interaction. In addition, our results are consistent with two theoretical calculations (black triangles) that reproduce the $\Lambda\Lambda$ binding energy of ${}^6_{\Lambda\Lambda}\text{He}$, as extracted from the NAGARA event [51,52].

7. Summary

The $K_S^0 K_S^0$, ΛK_S^0 , and $\Lambda\Lambda$ femtoscopic correlations are studied using lead-lead (PbPb) collision data at a center-of-mass energy per nucleon pair of $\sqrt{s_{NN}} = 5.02$ TeV, collected by the CMS Collaboration. This is the first report on $\Lambda\Lambda$ correlations in PbPb collisions at the CERN LHC. The source size R_{inv} and the correlation strength parameter λ were extracted for $K_S^0 K_S^0$ correlations in six centrality bins covering the 0–60% range. The value of R_{inv} decreases from 4.6 to 1.6 fm going from central to peripheral collisions and agrees with results from the ALICE Collaboration at a similar transverse mass. Along with the R_{inv} and λ parameters, the strong interaction scattering parameters, i.e., the complex scattering length and effective range, were extracted from ΛK_S^0 and $\Lambda\Lambda$ correlations in the 0–80% centrality range. These scattering parameters indicate that the ΛK_S^0 interaction is repulsive and that the $\Lambda\Lambda$ interaction is attractive. The scattering parameters obtained from ΛK_S^0 correlations differ from those reported by the ALICE Collaboration. The positive real scattering length obtained from the $\Lambda\Lambda$ correlation disfavors the existence of a bound H-dibaryon state. The $\Lambda\Lambda$ scattering parameters help to constrain baryon-baryon and, more specifically, hyperon-hyperon interaction models. These measurements provide an additional input to understand the nature of the strong interaction between pairs of strange hadrons.

Acknowledgements

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MOST, and NSFC (China); Minciencias (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MoSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 884104, and COST Action CA16108 (European Union); the Leventis Foundation; The Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science – EOS” – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010; The Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Deutsche Forschungsgemeinschaft (DFG), under Germany's Excellence Strategy – EXC 2121 “Quantum Universe”

– 390833306, and under project number 400140256 – GRK2497; the Hungarian Academy of Sciences, the New National Excellence Program – ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B05F650021 (Thailand); the Kavli Foundation; the Nvidia Corporation; the Super-Micro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the [CMS data preservation, re-use, and open access policy](#).

References

- [1] M.A. Lisa, S. Pratt, R. Soltz, U. Wiedemann, Femtoscopy in relativistic heavy ion collisions: two decades of progress, *Annu. Rev. Nucl. Part. Sci.* 55 (2005) 357, <https://doi.org/10.1146/annurev.nucl.55.090704.151533>, arXiv:nucl-ex/0505014.
- [2] CMS Collaboration, Bose–Einstein correlations in pp, pPb, and PbPb collisions at $\sqrt{s_{NN}} = 0.9$ –7 TeV, *Phys. Rev. C* 97 (2018) 064912, <https://doi.org/10.1103/PhysRevC.97.064912>, arXiv:1712.07198.
- [3] V.G.J. Stoks, R.A.M. Klomp, M.C.M. Rentmeester, J.J. de Swart, Partial-wave analysis of all nucleon-nucleon scattering data below 350 MeV, *Phys. Rev. C* 48 (1993) 792, <https://doi.org/10.1103/PhysRevC.48.792>.
- [4] J.J. de Swart, C. Dullemond, Effective range theory and the low energy hyperon-nucleon interactions, *Ann. Phys.* 19 (1962) 458, [https://doi.org/10.1016/0003-4916\(62\)90185-9](https://doi.org/10.1016/0003-4916(62)90185-9).
- [5] R. Engelmann, H. Filthuth, V. Hepp, E. Kluge, Inelastic Σ^- -p-interactions at low momenta, *Phys. Lett.* 21 (1966) 587, [https://doi.org/10.1016/0031-9163\(66\)91310-2](https://doi.org/10.1016/0031-9163(66)91310-2).
- [6] F. Eisele, H. Filthuth, W. Föhlich, V. Hepp, G. Zech, Elastic Σ^+p scattering at low energies, *Phys. Lett. B* 37 (1971) 204, [https://doi.org/10.1016/0370-2693\(71\)90053-0](https://doi.org/10.1016/0370-2693(71)90053-0).
- [7] B. Sechi-Zorn, B. Kehoe, J. Twitty, R.A. Burnstein, Low-energy Λ -proton elastic scattering, *Phys. Rev.* 175 (1968) 1735, <https://doi.org/10.1103/PhysRev.175.1735>.
- [8] D.B. Kaplan, A.E. Nelson, Kaon condensation in dense matter, *Nucl. Phys. A* 479 (1988) 273, [https://doi.org/10.1016/0375-9474\(88\)90442-3](https://doi.org/10.1016/0375-9474(88)90442-3).
- [9] J. Schaffner-Bielich, M. Hanauske, H. Stöcker, W. Greiner, Phase transition to hyperon matter in neutron stars, *Phys. Rev. Lett.* 89 (2002) 171101, <https://doi.org/10.1103/PhysRevLett.89.171101>, arXiv:astro-ph/0005490.
- [10] R.L. Jaffe, Perhaps a stable dihyperon, *Phys. Rev. Lett.* 38 (1977) 195, <https://doi.org/10.1103/PhysRevLett.38.195>.
- [11] H. Takahashi, et al., Observation of a ${}_{\Lambda\Lambda}^6\text{He}$ double hypernucleus, *Phys. Rev. Lett.* 87 (2001) 212502, <https://doi.org/10.1103/PhysRevLett.87.212502>.
- [12] K. Nakazawa, H. Takahashi, Experimental study of double- Λ hypernuclei with nuclear emulsion, *Prog. Theor. Phys. Suppl.* 185 (2010) 335, <https://doi.org/10.1143/PTPS.185.335>.
- [13] B.H. Kim, et al., Belle, Search for an H -dibaryon with a mass near $2m_{\Lambda}$ in $Y(1S)$ and $Y(2S)$ decays, *Phys. Rev. Lett.* 110 (2013) 222002, <https://doi.org/10.1103/PhysRevLett.110.222002>, arXiv:1302.4028.

- [14] ALICE Collaboration, AK femtoscopy in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. C 103 (2021) 055201, <https://doi.org/10.1103/PhysRevC.103.055201>, arXiv:2005.11124.
- [15] C. Loizides, J. Kamin, D. d'Enterria, Improved Monte Carlo Glauber predictions at present and future nuclear colliders, Phys. Rev. C 97 (2018) 054910, <https://doi.org/10.1103/PhysRevC.97.054910>, arXiv:1710.07098.
- [16] R. Lednický, V.L. Lyuboshitz, Final state interaction effect on pairing correlations between particles with small relative momenta, Sov. J. Nucl. Phys. 35 (1982) 770.
- [17] HEPData record for this analysis, <https://doi.org/10.17182/hepdata.133573>, 2022.
- [18] W. Adam, et al., Tracker Group of the CMS, The CMS phase-1 pixel detector upgrade, J. Instrum. 16 (2021) P02027, <https://doi.org/10.1088/1748-0221/16/02/P02027>, arXiv:2012.14304.
- [19] CMS Collaboration, Track impact parameter resolution for the full pseudorapidity coverage in the 2017 dataset with the CMS phase-1 pixel detector, CMS Detector Performance Note CMS-DP-2020-049, <https://cds.cern.ch/record/2743740>, 2020.
- [20] CMS Collaboration, Performance of the CMS level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV, J. Instrum. 15 (2020) P10017, <https://doi.org/10.1088/1748-0221/15/10/P10017>, arXiv:2006.10165.
- [21] CMS Collaboration, The CMS trigger system, J. Instrum. 12 (2017) 01020, <https://doi.org/10.1088/1748-0221/12/01/P01020>, arXiv:1609.02366.
- [22] CMS Collaboration, The CMS experiment at the CERN LHC, J. Instrum. 3 (2008) S08004, <https://doi.org/10.1088/1748-0221/3/08/S08004>.
- [23] CMS Collaboration, Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS, Eur. Phys. J. C 81 (2021) 800, <https://doi.org/10.1140/epjc/s10052-021-09538-2>, arXiv:2104.01927.
- [24] CMS Collaboration, CMS luminosity measurement using nucleus-nucleus collisions at $\sqrt{s_{NN}} = 5.02$ TeV, in: 2018, CMS Physics Analysis Summary CMS-PAS-LUM-18-001, 2022, <https://cds.cern.ch/record/2809613>.
- [25] CMS Collaboration, Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, J. High Energy Phys. 04 (2017) 039, [https://doi.org/10.1007/JHEP04\(2017\)039](https://doi.org/10.1007/JHEP04(2017)039), arXiv:1611.01664.
- [26] CMS Collaboration, Description and performance of track and primary-vertex reconstruction with the CMS tracker, J. Instrum. 9 (2014) P10009, <https://doi.org/10.1088/1748-0221/9/10/p10009>, arXiv:1405.6569.
- [27] CMS Collaboration, Transverse-momentum and pseudorapidity distributions of charged hadrons in pp collisions at $\sqrt{s} = 0.9$ and 2.36 TeV, J. High Energy Phys. 02 (2010) 041, [https://doi.org/10.1007/JHEP02\(2010\)041](https://doi.org/10.1007/JHEP02(2010)041), arXiv:1002.0621.
- [28] CMS Collaboration, Observation and studies of jet quenching in PbPb collisions at nucleon-nucleon $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. C 84 (2011) 024906, <https://doi.org/10.1103/PhysRevC.84.024906>, arXiv:1102.1957.
- [29] C. Gale, S. Jeon, B. Schenke, Hydrodynamic modeling of heavy ion collisions, Int. J. Mod. Phys. A 28 (2013) 1340011, <https://doi.org/10.1142/S0217751X13400113>, arXiv:1301.5893.
- [30] S. Agostinelli, et al., GEANT4, Geant4—a simulation toolkit, Nucl. Instrum. Methods A 506 (2003) 250, [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8).
- [31] CMS Collaboration, Strange hadron collectivity in pPb and PbPb collisions, J. High Energy Phys. 05 (2023) 007, [https://doi.org/10.1007/JHEP05\(2023\)007](https://doi.org/10.1007/JHEP05(2023)007), arXiv:2205.00080.
- [32] CMS Collaboration, Strange particle production in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV, J. High Energy Phys. 05 (2011) 064, [https://doi.org/10.1007/JHEP05\(2011\)064](https://doi.org/10.1007/JHEP05(2011)064), arXiv:1102.4282.
- [33] CMS Collaboration, Long-range two-particle correlations of strange hadrons with charged particles in pPb and PbPb collisions at LHC energies, Phys. Lett. B 742 (2015) 200, <https://doi.org/10.1016/j.physletb.2015.01.034>, arXiv:1409.3392.
- [34] R.L. Workman, et al., Particle Data Group, Review of particle physics, Prog. Theor. Exp. Phys. 2022 (2022) 083C01, <https://doi.org/10.1093/ptep/ptac097>.
- [35] H. Voss, A. Höcker, J. Stelzer, F. Tegenfeldt, TMVA, the toolkit for multivariate data analysis with ROOT, in: Xth International Workshop on Advanced Computing and Analysis Techniques in Physics Research (ACAT), 2007, p. 40, arXiv:physics/0703039.
- [36] G.I. Kopylov, Like particle correlations as a tool to study the multiple production mechanism, Phys. Lett. B 50 (1974) 472, [https://doi.org/10.1016/0370-2693\(74\)90263-9](https://doi.org/10.1016/0370-2693(74)90263-9).
- [37] A. Kisiel, Non-identical particle correlation analysis in the presence of non-femtoscopic correlations, Acta Phys. Pol. B 48 (2017) 717, <https://doi.org/10.5506/APhysPolB.48.717>.
- [38] ALICE Collaboration, pp, p-Λ, and Λ-Λ correlations studied via femtoscopy in pp reactions at $\sqrt{s} = 7$ TeV, Phys. Rev. C 99 (2019) 024001, <https://doi.org/10.1103/PhysRevC.99.024001>, arXiv:1805.12455.
- [39] B.I. Abelev, et al., STAR, Neutral kaon interferometry in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. C 74 (2006) 054902, <https://doi.org/10.1103/PhysRevC.74.054902>, arXiv:nucl-ex/0608012.
- [40] A.D. Martin, E.N. Ozmutlu, Analyses of K K production and scalar mesons, Nucl. Phys. B 158 (1979) 520, [https://doi.org/10.1016/0550-3213\(79\)90180-9](https://doi.org/10.1016/0550-3213(79)90180-9).
- [41] A. Antonelli, Radiative ϕ decays, eConfC 020620, 10.48550/HEP-EX/0209069, 2002, THAT06, arXiv:hep-ex/0209069, 2002.
- [42] N.N. Achasov, V.V. Gubin, Analysis of the nature of the $\phi \rightarrow \gamma\pi\eta$ and $\phi \rightarrow \gamma\pi^0\pi^0$ decays, Phys. Rev. D 63 (2001) 094007, <https://doi.org/10.1103/PhysRevD.63.094007>, arXiv:hep-ph/0101024.
- [43] N.N. Achasov, A.V. Kiselev, New analysis of the KLOE data on the $\phi \rightarrow \eta\pi^0\gamma$ decay, Phys. Rev. D 68 (2003) 014006, <https://doi.org/10.1103/PhysRevD.68.014006>, arXiv:hep-ph/0212153.
- [44] F. James, M. Roos, Minuit: a system for function minimization and analysis of the parameter errors and correlations, Comput. Phys. Commun. 10 (1975) 343, [https://doi.org/10.1016/0010-4655\(75\)90039-9](https://doi.org/10.1016/0010-4655(75)90039-9).
- [45] G. Bóró, G.G. Barnaföldi, G. Papp, M. Gyulassy, P. Lévai, X.-N. Wang, B.-W. Zhang, Introducing hijing++: the heavy ion Monte Carlo generator for the high-luminosity LHC era, <https://doi.org/10.48550/arXiv.1901.04220>, arXiv:1901.04220, 2019.
- [46] ALICE Collaboration, One-dimensional pion, kaon, and proton femtoscopy in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. C 92 (2015) 054908, <https://doi.org/10.1103/PhysRevC.92.054908>, arXiv:1506.07884.
- [47] CMS Collaboration, Charged-particle angular correlations in XeXe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, Phys. Rev. C 100 (2019) 044902, <https://doi.org/10.1103/PhysRevC.100.044902>, arXiv:1901.07997.
- [48] K. Morita, T. Furumoto, A. Ohnishi, $\Lambda\Lambda$ interaction from relativistic heavy-ion collisions, Phys. Rev. C 91 (2015) 024916, <https://doi.org/10.1103/PhysRevC.91.024916>, arXiv:1408.6682.
- [49] L. Adamczyk, et al., STAR, $\Lambda\Lambda$ correlation function in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. Lett. 114 (2015) 022301, <https://doi.org/10.1103/PhysRevLett.114.022301>, arXiv:1408.4360.
- [50] ALICE Collaboration, Study of the Λ - Λ interaction with femtoscopy correlations in pp and pPb collisions at the LHC, Phys. Lett. B 797 (2019) 134822, <https://doi.org/10.1016/j.physletb.2019.134822>, arXiv:1905.07209.
- [51] E. Hiyama, M. Kamimura, T. Motoba, T. Yamada, Y. Yamamoto, Four-body cluster structure of $A = 7$ –10 double- Λ hypernuclei, Phys. Rev. C 66 (2002) 024007, <https://doi.org/10.1103/physrevc.66.024007>, arXiv:nucl-th/0204059.
- [52] I.N. Filikhin, A. Gal, Faddeev–Yakubovsky calculations for light $\Lambda\Lambda$ hypernuclei, Nucl. Phys. A 707 (2002) 491, [https://doi.org/10.1016/S0375-9474\(02\)01008-4](https://doi.org/10.1016/S0375-9474(02)01008-4), arXiv:nucl-th/0203036.

The CMS collaboration

A. Tumasyan ¹

Yerevan Physics Institute, Yerevan, Armenia

W. Adam , J.W. Andrejkovic , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic ,
A. Escalante Del Valle , P.S. Hussain , M. Jeitler ², N. Krammer , L. Lechner , D. Liko , I. Mikulec ,
P. Paulitsch, J. Schieck ², R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger ,
C.-E. Wulz ²

Institut für Hochenergiephysik, Vienna, Austria

M.R. Darwish ³, T. Janssen , T. Kello ⁴, H. Rejeb Sfar, P. Van Mechelen 

Universiteit Antwerpen, Antwerpen, Belgium

E.S. Bols^{ID}, J. D'Hondt^{ID}, A. De Moor^{ID}, M. Delcourt^{ID}, H. El Faham^{ID}, S. Lowette^{ID}, A. Morton^{ID},
D. Müller^{ID}, A.R. Sahasransu^{ID}, S. Tavernier^{ID}, W. Van Doninck, S. Van Putte^{ID}, D. Vannerom^{ID}

Vrije Universiteit Brussel, Brussel, Belgium

B. Clerbaux^{ID}, G. De Lentdecker^{ID}, L. Favart^{ID}, D. Hohov^{ID}, J. Jaramillo^{ID}, K. Lee^{ID},
M. Mahdavihorrani^{ID}, I. Makarenko^{ID}, A. Malara^{ID}, S. Paredes^{ID}, L. Pétré^{ID}, N. Postiau, L. Thomas^{ID},
M. Vanden Bemden^{ID}, C. Vander Velde^{ID}, P. Vanlaer^{ID}

Université Libre de Bruxelles, Bruxelles, Belgium

D. Dobur^{ID}, J. Knolle^{ID}, L. Lambrecht^{ID}, G. Mestdach, C. Rendón, A. Samalan, K. Skovpen^{ID}, M. Tytgat^{ID},
N. Van Den Bossche^{ID}, B. Vermassen, L. Wezenbeek^{ID}

Ghent University, Ghent, Belgium

A. Benecke^{ID}, G. Bruno^{ID}, F. Bury^{ID}, C. Caputo^{ID}, P. David^{ID}, C. Delaere^{ID}, I.S. Donertas^{ID},
A. Giammanco^{ID}, K. Jaffel^{ID}, Sa. Jain^{ID}, V. Lemaître, K. Mondal^{ID}, A. Taliencio^{ID}, T.T. Tran^{ID}, P. Vischia^{ID},
S. Wertz^{ID}

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

G.A. Alves^{ID}, E. Coelho^{ID}, C. Hensel^{ID}, A. Moraes^{ID}, P. Rebello Teles^{ID}

Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

W.L. Aldá Júnior^{ID}, M. Alves Gallo Pereira^{ID}, M. Barroso Ferreira Filho^{ID}, H. Brandao Malbouisson^{ID},
W. Carvalho^{ID}, J. Chinellato⁵, E.M. Da Costa^{ID}, G.G. Da Silveira^{ID,6}, D. De Jesus Damiao^{ID},
V. Dos Santos Sousa^{ID}, S. Fonseca De Souza^{ID}, J. Martins^{ID,7}, C. Mora Herrera^{ID}, K. Mota Amarilo^{ID},
L. Mundim^{ID}, H. Nogima^{ID}, A. Santoro^{ID}, S.M. Silva Do Amaral^{ID}, A. Sznajder^{ID}, M. Thiel^{ID},
A. Vilela Pereira^{ID}

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

C.A. Bernardes^{ID,6}, L. Calligaris^{ID}, T.R. Fernandez Perez Tomei^{ID}, E.M. Gregores^{ID}, P.G. Mercadante^{ID},
S.F. Novaes^{ID}, Sandra S. Padula^{ID}

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

A. Aleksandrov^{ID}, G. Antchev^{ID}, R. Hadjiiska^{ID}, P. Iaydjiev^{ID}, M. Misheva^{ID}, M. Rodozov, M. Shopova^{ID},
G. Sultanov^{ID}

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Dimitrov^{ID}, T. Ivanov^{ID}, L. Litov^{ID}, B. Pavlov^{ID}, P. Petkov^{ID}, A. Petrov^{ID}, E. Shumka^{ID}

University of Sofia, Sofia, Bulgaria

S. Thakur^{ID}

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

T. Cheng^{ID}, T. Javaid^{ID,8}, M. Mittal^{ID}, L. Yuan^{ID}

Beihang University, Beijing, China

M. Ahmad^{ID}, G. Bauer⁹, Z. Hu^{ID}, S. Lezki^{ID}, K. Yi^{ID,9,10}

Department of Physics, Tsinghua University, Beijing, China

G.M. Chen^{ID,8}, H.S. Chen^{ID,8}, M. Chen^{ID,8}, F. Iemmi^{ID}, C.H. Jiang, A. Kapoor^{ID}, H. Liao^{ID}, Z.-A. Liu^{ID,11},
V. Milosevic^{ID}, F. Monti^{ID}, R. Sharma^{ID}, J. Tao^{ID}, J. Thomas-Wilsker^{ID}, J. Wang^{ID}, H. Zhang^{ID}, J. Zhao^{ID}

Institute of High Energy Physics, Beijing, China

A. Agapitos^{ID}, Y. An^{ID}, Y. Ban^{ID}, A. Levin^{ID}, C. Li^{ID}, Q. Li^{ID}, X. Lyu, Y. Mao, S.J. Qian^{ID}, X. Sun^{ID},
D. Wang^{ID}, J. Xiao^{ID}, H. Yang

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

M. Lu^{ID}, Z. You^{ID}

Sun Yat-Sen University, Guangzhou, China

N. Lu^{ID}

University of Science and Technology of China, Hefei, China

X. Gao^{ID,4}, D. Leggat, H. Okawa^{ID}, Y. Zhang^{ID}

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Z. Lin^{ID}, C. Lu^{ID}, M. Xiao^{ID}

Zhejiang University, Hangzhou, Zhejiang, China

C. Avila^{ID}, D.A. Barbosa Trujillo, A. Cabrera^{ID}, C. Florez^{ID}, J. Fraga^{ID}

Universidad de Los Andes, Bogota, Colombia

J. Mejia Guisao^{ID}, F. Ramirez^{ID}, M. Rodriguez^{ID}, J.D. Ruiz Alvarez^{ID}

Universidad de Antioquia, Medellin, Colombia

D. Giljanovic^{ID}, N. Godinovic^{ID}, D. Lelas^{ID}, I. Puljak^{ID}

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

Z. Antunovic, M. Kovac^{ID}, T. Sculac^{ID}

University of Split, Faculty of Science, Split, Croatia

V. Brigljevic^{ID}, B.K. Chitroda^{ID}, D. Ferencek^{ID}, S. Mishra^{ID}, M. Roguljic^{ID}, A. Starodumov^{ID,12}, T. Susa^{ID}

Institute Rudjer Boskovic, Zagreb, Croatia

A. Attikis^{ID}, K. Christoforou^{ID}, S. Konstantinou^{ID}, J. Mousa^{ID}, C. Nicolaou, F. Ptochos^{ID}, P.A. Razis^{ID},
H. Rykaczewski, H. Saka^{ID}, A. Stepennov^{ID}

University of Cyprus, Nicosia, Cyprus

M. Finger^{ID}, M. Finger Jr.^{ID}, A. Kveton^{ID}

Charles University, Prague, Czech Republic

E. Ayala^{ID}

Escuela Politecnica Nacional, Quito, Ecuador

E. Carrera Jarrin^{ID}

Universidad San Francisco de Quito, Quito, Ecuador

A.A. Abdelalim^{ID,13,14}, E. Salama^{ID,15,16}

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

M.A. Mahmoud^{ID}, Y. Mohammed^{ID}

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

S. Bhowmik^{ID}, R.K. Dewanjee^{ID}, K. Ehataht^{ID}, M. Kadastik, T. Lange^{ID}, S. Nandan^{ID}, C. Nielsen^{ID}, J. Pata^{ID},
M. Raidal^{ID}, L. Tani^{ID}, C. Veelken^{ID}

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

P. Eerola^{ID}, H. Kirschenmann^{ID}, K. Osterberg^{ID}, M. Voutilainen^{ID}

Department of Physics, University of Helsinki, Helsinki, Finland

S. Bharthuar^{ID}, E. Brücken^{ID}, F. Garcia^{ID}, J. Havukainen^{ID}, M.S. Kim^{ID}, R. Kinnunen, T. Lampén^{ID},
K. Lassila-Perini^{ID}, S. Lehti^{ID}, T. Lindén^{ID}, M. Lotti, L. Martikainen^{ID}, M. Myllymäki^{ID}, M.m. Rantanen^{ID},
H. Siikonen^{ID}, E. Tuominen^{ID}, J. Tuominiemi^{ID}

Helsinki Institute of Physics, Helsinki, Finland

P. Luukka^{ID}, H. Petrow^{ID}, T. Tuuva[†]

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

C. Amendola^{ID}, M. Besancon^{ID}, F. Couderc^{ID}, M. Dejardin^{ID}, D. Denegri, J.L. Faure, F. Ferri^{ID}, S. Ganjour^{ID},
P. Gras^{ID}, G. Hamel de Monchenault^{ID}, V. Lohezic^{ID}, J. Malcles^{ID}, J. Rander, A. Rosowsky^{ID}, M.Ö. Sahin^{ID},
A. Savoy-Navarro^{ID,17}, P. Simkina^{ID}, M. Titov^{ID}

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

C. Baldenegro Barrera^{ID}, F. Beaudette^{ID}, A. Buchot Perraguin^{ID}, P. Busson^{ID}, A. Cappati^{ID}, C. Charlot^{ID},
F. Damas^{ID}, O. Davignon^{ID}, B. Diab^{ID}, G. Falmagne^{ID}, B.A. Fontana Santos Alves^{ID}, S. Ghosh^{ID},
R. Granier de Cassagnac^{ID}, A. Hakimi^{ID}, B. Harikrishnan^{ID}, G. Liu^{ID}, J. Motta^{ID}, M. Nguyen^{ID},
C. Ochando^{ID}, L. Portales^{ID}, R. Salerno^{ID}, U. Sarkar^{ID}, J.B. Sauvan^{ID}, Y. Sirois^{ID}, A. Tarabini^{ID},
E. Vernazza^{ID}, A. Zabi^{ID}, A. Zghiche^{ID}

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

J.-L. Agram^{ID,18}, J. Andrea^{ID}, D. Apparu^{ID}, D. Bloch^{ID}, G. Bourgatte^{ID}, J.-M. Brom^{ID}, E.C. Chabert^{ID},
C. Collard^{ID}, D. Darej, U. Goerlach^{ID}, C. Grimault, A.-C. Le Bihan^{ID}, P. Van Hove^{ID}

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

S. Beauceron^{ID}, B. Blancon^{ID}, G. Boudoul^{ID}, A. Carle, N. Chanon^{ID}, J. Choi^{ID}, D. Contardo^{ID}, P. Depasse^{ID},
C. Dozen^{ID,19}, H. El Mamouni, J. Fay^{ID}, S. Gascon^{ID}, M. Gouzevitch^{ID}, G. Grenier^{ID}, B. Ille^{ID}, I.B. Laktineh,
M. Lethuillier^{ID}, L. Mirabito, S. Perries, L. Torterotot^{ID}, M. Vander Donckt^{ID}, P. Verdier^{ID}, S. Viret

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

I. Bagaturia^{ID,20}, I. Lomidze^{ID}, Z. Tsamalaidze^{ID,12}

Georgian Technical University, Tbilisi, Georgia

V. Botta^{ID}, L. Feld^{ID}, K. Klein^{ID}, M. Lipinski^{ID}, D. Meuser^{ID}, A. Pauls^{ID}, N. Röwert^{ID}, M. Teroerde^{ID}

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

S. Diekmann^{ID}, A. Dodonova^{ID}, N. Eich^{ID}, D. Eliseev^{ID}, M. Erdmann^{ID}, P. Fackeldey^{ID}, D. Fasanella^{ID},
B. Fischer^{ID}, T. Hebbeker^{ID}, K. Hoepfner^{ID}, F. Ivone^{ID}, M.y. Lee^{ID}, L. Mastrolorenzo, M. Merschmeyer^{ID},
A. Meyer^{ID}, S. Mondal^{ID}, S. Mukherjee^{ID}, D. Noll^{ID}, A. Novak^{ID}, F. Nowotny, A. Pozdnyakov^{ID}, Y. Rath,
W. Redjeb^{ID}, H. Reithler^{ID}, A. Schmidt^{ID}, S.C. Schuler, A. Sharma^{ID}, A. Stein^{ID},
F. Torres Da Silva De Araujo^{ID,21}, L. Vigilante, S. Wiedenbeck^{ID}, S. Zaleski

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

C. Dziwok^{ID}, G. Flügge^{ID}, W. Haj Ahmad^{ID,22}, O. Hlushchenko, T. Kress^{ID}, A. Nowack^{ID}, O. Pooth^{ID},
A. Stahl^{ID}, T. Ziemons^{ID}, A. Zotz^{ID}

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

H. Aarup Petersen^{ID}, M. Aldaya Martin^{ID}, J. Alimena^{ID}, P. Asmuss, S. Baxter^{ID}, M. Bayatmakou^{ID},
H. Becerril Gonzalez^{ID}, O. Behnke^{ID}, S. Bhattacharya^{ID}, F. Blekman^{ID,23}, K. Borras^{ID,24}, D. Brunner^{ID},
A. Campbell^{ID}, A. Cardini^{ID}, C. Cheng, F. Colombina^{ID}, S. Consuegra Rodríguez^{ID}, G. Correia Silva^{ID},
M. De Silva^{ID}, G. Eckerlin, D. Eckstein^{ID}, L.I. Estevez Banos^{ID}, O. Filatov^{ID}, E. Gallo^{ID,23}, A. Geiser^{ID},
A. Giraldi^{ID}, G. Greau, A. Grohsjean^{ID}, V. Guglielmi^{ID}, M. Guthoff^{ID}, A. Jafari^{ID,25}, N.Z. Jomhari^{ID},
B. Kaech^{ID}, M. Kasemann^{ID}, H. Kaveh^{ID}, C. Kleinwort^{ID}, R. Kogler^{ID}, M. Komm^{ID}, D. Krücker^{ID}, W. Lange,
D. Leyva Pernia^{ID}, K. Lipka^{ID,26}, W. Lohmann^{ID,27}, R. Mankel^{ID}, I.-A. Melzer-Pellmann^{ID},
M. Mendizabal Morentin^{ID}, J. Metwally, A.B. Meyer^{ID}, G. Milella^{ID}, M. Mormile^{ID}, A. Mussgiller^{ID},
A. Nürnberg^{ID}, Y. Otarid, D. Pérez Adán^{ID}, E. Ranken^{ID}, A. Raspereza^{ID}, B. Ribeiro Lopes^{ID}, J. Rübenach,
A. Saggio^{ID}, M. Savitskyi^{ID}, M. Scham^{ID,28,24}, V. Scheurer, S. Schnake^{ID,24}, P. Schütze^{ID},
C. Schwanenberger^{ID,23}, M. Shchedrolosiev^{ID}, R.E. Sosa Ricardo^{ID}, D. Stafford, N. Tonon^{ID,†},
M. Van De Klundert^{ID}, F. Vazzoler^{ID}, A. Ventura Barroso^{ID}, R. Walsh^{ID}, D. Walter^{ID}, Q. Wang^{ID}, Y. Wen^{ID},
K. Wichmann, L. Wiens^{ID,24}, C. Wissing^{ID}, S. Wuchterl^{ID}, Y. Yang^{ID}, A. Zimmermann Castro Santos^{ID}

Deutsches Elektronen-Synchrotron, Hamburg, Germany

A. Albrecht^{ID}, S. Albrecht^{ID}, M. Antonello^{ID}, S. Bein^{ID}, L. Benato^{ID}, M. Bonanomi^{ID}, P. Connor^{ID},
K. De Leo^{ID}, M. Eich, K. El Morabit^{ID}, F. Feindt, A. Fröhlich, C. Garbers^{ID}, E. Garutti^{ID}, M. Hajheidari,
J. Haller^{ID}, A. Hinzmann^{ID}, H.R. Jabusch^{ID}, G. Kasieczka^{ID}, P. Keicher, R. Klanner^{ID}, W. Korcar^{ID},
T. Kramer^{ID}, V. Kutzner^{ID}, F. Labe^{ID}, J. Lange^{ID}, A. Lobanov^{ID}, C. Matthies^{ID}, A. Mehta^{ID}, L. Moureaux^{ID},
M. Mrowietz, A. Nigamova^{ID}, Y. Nissan, A. Paasch^{ID}, K.J. Pena Rodriguez^{ID}, T. Quadfasel^{ID}, M. Rieger^{ID},
O. Rieger, D. Savoie^{ID}, J. Schindler^{ID}, P. Schleper^{ID}, M. Schröder^{ID}, J. Schwandt^{ID}, M. Sommerhalder^{ID},
H. Stadie^{ID}, G. Steinbrück^{ID}, A. Tews, M. Wolf^{ID}

University of Hamburg, Hamburg, Germany

S. Brommer^{ID}, M. Burkart, E. Butz^{ID}, T. Chwalek^{ID}, A. Dierlamm^{ID}, A. Droll, N. Faltermann^{ID}, M. Giffels^{ID},
J.O. Gosewisch, A. Gottmann^{ID}, F. Hartmann^{ID,29}, M. Horzela^{ID}, U. Husemann^{ID}, M. Klute^{ID},
R. Koppenhöfer^{ID}, M. Link, A. Lintuluoto^{ID}, S. Maier^{ID}, S. Mitra^{ID}, Th. Müller^{ID}, M. Neukum, M. Oh^{ID},
G. Quast^{ID}, K. Rabbertz^{ID}, J. Rauser, I. Shvetsov^{ID}, H.J. Simonis^{ID}, N. Trevisani^{ID}, R. Ulrich^{ID},
J. van der Linden^{ID}, R.F. Von Cube^{ID}, M. Wassmer^{ID}, S. Wieland^{ID}, R. Wolf^{ID}, S. Wozniowski^{ID}, S. Wunsch,
X. Zuo^{ID}

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

G. Anagnostou, P. Assiouras^{ID}, G. Daskalakis^{ID}, A. Kyriakis, A. Stakia^{ID}

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

M. Diamantopoulou, D. Karasavvas, P. Kontaxakis^{ID}, A. Manousakis-Katsikakis^{ID}, A. Panagiotou,
I. Papavergou^{ID}, N. Saoulidou^{ID}, K. Theofilatos^{ID}, E. Tziaferi^{ID}, K. Vellidis^{ID}, I. Zisopoulos^{ID}

National and Kapodistrian University of Athens, Athens, Greece

G. Bakas^{ID}, T. Chatzistavrou, G. Karapostoli^{ID}, K. Kousouris^{ID}, I. Papakrivopoulos^{ID}, G. Tsipolitis, A. Zacharopoulou

National Technical University of Athens, Athens, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou^{ID}, C. Foudas, P. Gianneios^{ID}, C. Kamtsikis, P. Katsoulis, P. Kokkas^{ID}, P.G. Kosmoglou Kioseoglou^{ID}, N. Manthos^{ID}, I. Papadopoulos^{ID}, J. Strologas^{ID}

University of Ioánnina, Ioánnina, Greece

M. Bartók^{ID,30}, G. Bencze, C. Hajdu^{ID}, D. Horvath^{ID,31,32}, F. Sikler^{ID}, V. Veszpremi^{ID}

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

M. Csanád^{ID}, K. Farkas^{ID}, M.M.A. Gadallah^{ID,33}, S. Lökös^{ID,34}, P. Major^{ID}, K. Mandal^{ID}, G. Pásztor^{ID}, A.J. Rádl^{ID,35}, O. Surányi^{ID}, G.I. Veres^{ID}

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

N. Beni^{ID}, S. Czellar, J. Karancsi^{ID,30}, J. Molnar, Z. Szillasi, D. Teyssier^{ID}

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

P. Raics, B. Ujvari^{ID,36}, G. Zilizi^{ID}

Institute of Physics, University of Debrecen, Debrecen, Hungary

T. Csorgo^{ID,35}, F. Nemes^{ID,35}, T. Novak^{ID}

Károly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

J. Babbar^{ID}, S. Bansal^{ID}, S.B. Beri, V. Bhatnagar^{ID}, G. Chaudhary^{ID}, S. Chauhan^{ID}, N. Dhingra^{ID,37}, R. Gupta, A. Kaur^{ID}, A. Kaur^{ID}, H. Kaur^{ID}, M. Kaur^{ID}, S. Kumar^{ID}, P. Kumari^{ID}, M. Meena^{ID}, K. Sandeep^{ID}, T. Sheokand, J.B. Singh^{ID,38}, A. Singla^{ID}, A.K. Virdi^{ID}

Panjab University, Chandigarh, India

A. Ahmed^{ID}, A. Bhardwaj^{ID}, A. Chhetri^{ID}, B.C. Choudhary^{ID}, A. Kumar^{ID}, M. Naimuddin^{ID}, K. Ranjan^{ID}, S. Saumya^{ID}

University of Delhi, Delhi, India

S. Baradia^{ID}, S. Barman^{ID,39}, S. Bhattacharya^{ID}, D. Bhowmik, S. Dutta^{ID}, S. Dutta, B. Gomber^{ID,40}, M. Maity^{ID,39}, P. Palit^{ID}, G. Saha^{ID}, B. Sahu^{ID}, S. Sarkar

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

P.K. Behera^{ID}, S.C. Behera^{ID}, S. Chatterjee^{ID}, P. Kalbhor^{ID}, J.R. Komaragiri^{ID,41}, D. Kumar^{ID,41}, A. Muhammad^{ID}, L. Panwar^{ID,41}, R. Pradhan^{ID}, P.R. Pujahari^{ID}, N.R. Saha^{ID}, A. Sharma^{ID}, A.K. Sikdar^{ID}, S. Verma^{ID}

Indian Institute of Technology Madras, Madras, India

K. Naskar^{ID,42}

Bhabha Atomic Research Centre, Mumbai, India

T. Aziz, I. Das^{ID}, S. Dugad, M. Kumar^{ID}, G.B. Mohanty^{ID}, P. Suryadevara

Tata Institute of Fundamental Research-A, Mumbai, India

S. Banerjee^{ID}, M. Guchait^{ID}, S. Karmakar^{ID}, S. Kumar^{ID}, G. Majumder^{ID}, K. Mazumdar^{ID}, S. Mukherjee^{ID}, A. Thachayath^{ID}

Tata Institute of Fundamental Research-B, Mumbai, India

S. Bahinipati^{ID,43}, A.K. Das^{ID}, C. Kar^{ID}, P. Mal^{ID}, T. Mishra^{ID}, V.K. Muraleedharan Nair Bindhu^{ID,44},
A. Nayak^{ID,44}, P. Saha^{ID}, S.K. Swain^{ID}, D. Vats^{ID,44}

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

A. Alpana^{ID}, S. Dube^{ID}, B. Kansal^{ID}, A. Laha^{ID}, S. Pandey^{ID}, A. Rastogi^{ID}, S. Sharma^{ID}

Indian Institute of Science Education and Research (IISER), Pune, India

H. Bakhshiansohi^{ID,45,46}, E. Khazaie^{ID,46}, M. Zeinali^{ID,47}

Isfahan University of Technology, Isfahan, Iran

S. Chenarani^{ID,48}, S.M. Etesami^{ID}, M. Khakzad^{ID}, M. Mohammadi Najafabadi^{ID}

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

M. Grunewald^{ID}

University College Dublin, Dublin, Ireland

M. Abbrescia^{a,b,ID}, R. Aly^{a,c,ID,13}, C. Aruta^{a,b,ID}, A. Colaleo^{a,ID}, D. Creanza^{a,c,ID}, L. Cristella^{a,b,ID},
N. De Filippis^{a,c,ID}, M. De Palma^{a,b,ID}, A. Di Florio^{a,b,ID}, W. Elmetenawee^{a,b,ID}, F. Errico^{a,b,ID}, L. Fiore^{a,ID},
G. Iaselli^{a,c,ID}, G. Maggi^{a,c,ID}, M. Maggi^{a,ID}, I. Margjeka^{a,b,ID}, V. Mastrapasqua^{a,b,ID}, S. My^{a,b,ID}, S. Nuzzo^{a,b,ID},
A. Pellecchia^{a,b,ID}, A. Pompili^{a,b,ID}, G. Pugliese^{a,c,ID}, R. Radogna^{a,ID}, D. Ramos^{a,ID}, A. Ranieri^{a,ID},
G. Selvaggi^{a,b,ID}, L. Silvestris^{a,ID}, F.M. Simone^{a,b,ID}, Ü. Sözbilir^{a,ID}, A. Stamerra^{a,ID}, R. Venditti^{a,ID},
P. Verwilligen^{a,ID}

^a INFN Sezione di Bari, Bari, Italy^b Università di Bari, Bari, Italy^c Politecnico di Bari, Bari, Italy

G. Abbiendi^{a,ID}, C. Battilana^{a,b,ID}, D. Bonacorsi^{a,b,ID}, L. Borgonovi^{a,ID}, L. Brigliadori^a, R. Campanini^{a,b,ID},
P. Capiluppi^{a,b,ID}, A. Castro^{a,b,ID}, F.R. Cavallo^{a,ID}, M. Cuffiani^{a,b,ID}, G.M. Dallavalle^{a,ID}, T. Diotallevi^{a,b,ID},
F. Fabbri^{a,ID}, A. Fanfani^{a,b,ID}, P. Giacomelli^{a,ID}, L. Giommi^{a,b,ID}, C. Grandi^{a,ID}, L. Guiducci^{a,b,ID},
S. Lo Meo^{a,b,ID,49}, L. Lunerti^{a,b,ID}, S. Marcellini^{a,ID}, G. Masetti^{a,ID}, F.L. Navarria^{a,b,ID}, A. Perrotta^{a,ID},
F. Primavera^{a,b,ID}, A.M. Rossi^{a,b,ID}, T. Rovelli^{a,b,ID}, G.P. Siroli^{a,b,ID}

^a INFN Sezione di Bologna, Bologna, Italy^b Università di Bologna, Bologna, Italy

S. Costa^{a,b,ID,50}, A. Di Mattia^{a,ID}, R. Potenza^{a,b}, A. Tricomi^{a,b,ID,50}, C. Tuve^{a,b,ID}

^a INFN Sezione di Catania, Catania, Italy^b Università di Catania, Catania, Italy

G. Barbagli^{a,ID}, G. Bardelli^{a,b,ID}, B. Camaiani^{a,b,ID}, A. Cassese^{a,ID}, R. Ceccarelli^{a,b,ID}, V. Ciulli^{a,b,ID},
C. Civinini^{a,ID}, R. D'Alessandro^{a,b,ID}, E. Focardi^{a,b,ID}, G. Latino^{a,b,ID}, P. Lenzi^{a,b,ID}, M. Lizzo^{a,b,ID},
M. Meschini^{a,ID}, S. Paoletti^{a,ID}, G. Sguazzoni^{a,ID}, L. Viliani^{a,ID}

^a INFN Sezione di Firenze, Firenze, Italy^b Università di Firenze, Firenze, Italy

L. Benussi^{ID}, S. Bianco^{ID}, S. Meola^{ID,51}, D. Piccolo^{ID}

INFN Laboratori Nazionali di Frascati, Frascati, Italy

M. Bozzo^{a,b,ID}, P. Chatagnon^{a,ID}, F. Ferro^{a,ID}, E. Robutti^{a,ID}, S. Tosi^{a,b,ID}

^a INFN Sezione di Genova, Genova, Italy^b Università di Genova, Genova, Italy

A. Benaglia^{a, id}, G. Boldrini^{a, id}, F. Brivio^{a, b, id}, F. Cetorelli^{a, b, id}, F. De Guio^{a, b, id}, M.E. Dinardo^{a, b, id}, P. Dini^{a, id},
 S. Gennai^{a, id}, A. Ghezzi^{a, b, id}, P. Govoni^{a, b, id}, L. Guzzi^{a, b, id}, M.T. Lucchini^{a, b, id}, M. Malberti^{a, id},
 S. Malvezzi^{a, id}, A. Massironi^{a, id}, D. Menasce^{a, id}, L. Moroni^{a, id}, M. Paganoni^{a, b, id}, D. Pedrini^{a, id},
 B.S. Pinolini^a, S. Ragazzi^{a, b, id}, N. Redaelli^{a, id}, T. Tabarelli de Fatis^{a, b, id}, D. Zuolo^{a, b, id}

^a INFN Sezione di Milano-Bicocca, Milano, Italy

^b Università di Milano-Bicocca, Milano, Italy

S. Buontempo^{a, id}, F. Carnevali^{a, b}, N. Cavallo^{a, c, id}, A. De Iorio^{a, b, id}, F. Fabozzi^{a, c, id}, A.O.M. Iorio^{a, b, id},
 L. Lista^{a, b, id, 52}, P. Paolucci^{a, id, 29}, B. Rossi^{a, id}, C. Sciacca^{a, b, id}

^a INFN Sezione di Napoli, Napoli, Italy

^b Università di Napoli 'Federico II', Napoli, Italy

^c Università della Basilicata, Potenza, Italy

^d Università G. Marconi, Roma, Italy

P. Azzi^{a, id}, N. Bacchetta^{a, id, 53}, M. Biasotto^{a, id, 54}, P. Bortignon^{a, id}, A. Bragagnolo^{a, b, id}, R. Carlin^{a, b, id},
 P. Checchia^{a, id}, S. Fantinel^{a, id}, F. Gasparini^{a, b, id}, U. Gasparini^{a, b, id}, G. Grosso^a, L. Layer^{a, 55}, E. Lusiani^{a, id},
 M. Margoni^{a, b, id}, A.T. Meneguzzo^{a, b, id}, J. Pazzini^{a, b, id}, P. Ronchese^{a, b, id}, R. Rossin^{a, b, id}, F. Simonetto^{a, b, id},
 G. Strong^{a, id}, M. Tosi^{a, b, id}, H. Yarar^{a, b}, M. Zanetti^{a, b, id}, P. Zotto^{a, b, id}, A. Zucchetta^{a, b, id}, G. Zumerle^{a, b, id}

^a INFN Sezione di Padova, Padova, Italy

^b Università di Padova, Padova, Italy

^c Università di Trento, Trento, Italy

S. Abu Zeid^{a, id, 16}, C. Aimè^{a, b, id}, A. Braghieri^{a, id}, S. Calzaferri^{a, b, id}, D. Fiorina^{a, b, id}, P. Montagna^{a, b, id},
 V. Re^{a, id}, C. Riccardi^{a, b, id}, P. Salvini^{a, id}, I. Vai^{a, id}, P. Vitulo^{a, b, id}

^a INFN Sezione di Pavia, Pavia, Italy

^b Università di Pavia, Pavia, Italy

P. Asenov^{a, id, 56}, G.M. Bilei^{a, id}, D. Ciangottini^{a, b, id}, L. Fanò^{a, b, id}, M. Magherini^{a, b, id}, G. Mantovani^{a, b},
 V. Mariani^{a, b, id}, M. Menichelli^{a, id}, F. Moscatelli^{a, id, 56}, A. Piccinelli^{a, b, id}, M. Presilla^{a, b, id}, A. Rossi^{a, b, id},
 A. Santocchia^{a, b, id}, D. Spiga^{a, id}, T. Tedeschi^{a, b, id}

^a INFN Sezione di Perugia, Perugia, Italy

^b Università di Perugia, Perugia, Italy

P. Azzurri^{a, id}, G. Bagliesi^{a, id}, V. Bertacchi^{a, c, id}, R. Bhattacharya^{a, id}, L. Bianchini^{a, b, id}, T. Boccali^{a, id},
 E. Bossini^{a, b, id}, D. Bruschini^{a, c, id}, R. Castaldi^{a, id}, M.A. Ciocci^{a, b, id}, V. D'Amante^{a, d, id}, R. Dell'Orso^{a, id},
 S. Donato^{a, id}, A. Giassi^{a, id}, F. Ligabue^{a, c, id}, D. Matos Figueiredo^{a, id}, A. Messineo^{a, b, id}, M. Musich^{a, b, id},
 F. Palla^{a, id}, S. Parolia^{a, id}, G. Ramirez-Sanchez^{a, c, id}, A. Rizzi^{a, b, id}, G. Rolandi^{a, c, id}, S. Roy Chowdhury^{a, id},
 T. Sarkar^{a, id}, A. Scribano^{a, id}, P. Spagnolo^{a, id}, R. Tenchini^{a, id}, G. Tonelli^{a, b, id}, N. Turini^{a, d, id}, A. Venturi^{a, id},
 P.G. Verдини^{a, id}

^a INFN Sezione di Pisa, Pisa, Italy

^b Università di Pisa, Pisa, Italy

^c Scuola Normale Superiore di Pisa, Pisa, Italy

^d Università di Siena, Siena, Italy

P. Barria^{a, id}, M. Campana^{a, b, id}, F. Cavallari^{a, id}, D. Del Re^{a, b, id}, E. Di Marco^{a, id}, M. Diemoz^{a, id}, E. Longo^{a, b, id},
 P. Meridiani^{a, id}, G. Organtini^{a, b, id}, F. Pandolfi^{a, id}, R. Paramatti^{a, b, id}, C. Quaranta^{a, b, id}, S. Rahatlou^{a, b, id},
 C. Rovelli^{a, id}, F. Santanastasio^{a, b, id}, L. Soffi^{a, id}, R. Tramontano^{a, b, id}

^a INFN Sezione di Roma, Roma, Italy

^b Sapienza Università di Roma, Roma, Italy

N. Amapane^{a,b, ID}, R. Arcidiacono^{a,c, ID}, S. Argiro^{a,b, ID}, M. Arneodo^{a,c, ID}, N. Bartosik^{a, ID}, R. Bellan^{a,b, ID},
 A. Bellora^{a,b, ID}, C. Biino^{a, ID}, N. Cartiglia^{a, ID}, M. Costa^{a,b, ID}, R. Covarelli^{a,b, ID}, N. Demaria^{a, ID}, M. Grippo^{a,b, ID},
 B. Kiani^{a,b, ID}, F. Legger^{a, ID}, C. Mariotti^{a, ID}, S. Maselli^{a, ID}, A. Mecca^{a,b, ID}, E. Migliore^{a,b, ID}, M. Monteno^{a, ID},
 R. Mulargia^{a, ID}, M.M. Obertino^{a,b, ID}, G. Ortona^{a, ID}, L. Pacher^{a,b, ID}, N. Pastrone^{a, ID}, M. Pelliccioni^{a, ID},
 M. Ruspa^{a,c, ID}, K. Shchelina^{a, ID}, F. Siviero^{a,b, ID}, V. Sola^{a,b, ID}, A. Solano^{a,b, ID}, D. Soldi^{a,b, ID}, A. Staiano^{a, ID},
 M. Tornago^{a,b, ID}, D. Trocino^{a, ID}, G. Umoret^{a,b, ID}, A. Vagnerini^{a,b, ID}, E. Vlasov^{a,b, ID}

^a INFN Sezione di Torino, Torino, Italy

^b Università di Torino, Torino, Italy

^c Università del Piemonte Orientale, Novara, Italy

S. Belforte^{a, ID}, V. Candelise^{a,b, ID}, M. Casarsa^{a, ID}, F. Cossutti^{a, ID}, G. Della Ricca^{a,b, ID}, G. Sorrentino^{a,b, ID}

^a INFN Sezione di Trieste, Trieste, Italy

^b Università di Trieste, Trieste, Italy

S. Dogra^{ID}, C. Huh^{ID}, B. Kim^{ID}, D.H. Kim^{ID}, G.N. Kim^{ID}, J. Kim, J. Lee^{ID}, S.W. Lee^{ID}, C.S. Moon^{ID},
 Y.D. Oh^{ID}, S.I. Pak^{ID}, M.S. Ryu^{ID}, S. Sekmen^{ID}, Y.C. Yang^{ID}

Kyungpook National University, Daegu, Korea

H. Kim^{ID}, D.H. Moon^{ID}

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

E. Asilar^{ID}, T.J. Kim^{ID}, J. Park^{ID}

Hanyang University, Seoul, Korea

S. Choi^{ID}, S. Han, B. Hong^{ID}, K. Lee, K.S. Lee^{ID}, J. Lim, J. Park, S.K. Park, J. Yoo^{ID}

Korea University, Seoul, Korea

J. Goh^{ID}

Kyung Hee University, Department of Physics, Seoul, Korea

H.S. Kim^{ID}, Y. Kim, S. Lee

Sejong University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi^{ID}, S. Jeon^{ID}, J. Kim^{ID}, J.S. Kim, S. Ko^{ID}, H. Kwon^{ID}, H. Lee^{ID}, S. Lee,
 B.H. Oh^{ID}, S.B. Oh^{ID}, H. Seo^{ID}, U.K. Yang, I. Yoon^{ID}

Seoul National University, Seoul, Korea

W. Jang^{ID}, D.Y. Kang, Y. Kang^{ID}, D. Kim^{ID}, S. Kim^{ID}, B. Ko, J.S.H. Lee^{ID}, Y. Lee^{ID}, J.A. Merlin, I.C. Park^{ID},
 Y. Roh, D. Song, I.J. Watson^{ID}, S. Yang^{ID}

University of Seoul, Seoul, Korea

S. Ha^{ID}, H.D. Yoo^{ID}

Yonsei University, Department of Physics, Seoul, Korea

M. Choi^{ID}, M.R. Kim^{ID}, H. Lee, Y. Lee^{ID}, I. Yu^{ID}

Sungkyunkwan University, Suwon, Korea

T. Beyrouthy, Y. Maghrbi^{ID}

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

K. Dreimanis^{ID}, G. Pikurs, A. Potrebko^{ID}, M. Seidel^{ID}, V. Veckalns^{ID}

Riga Technical University, Riga, Latvia

M. Ambroz^a, A. Carvalho Antunes De Oliveira^a, A. Juodagalvis^a, A. Rinkevicius^a, G. Tamulaitis^a

Vilnius University, Vilnius, Lithuania

N. Bin Norjoharuddeen^b, S.Y. Hoh^{b,57}, I. Yusuff^{b,57}, Z. Zolkapli

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

J.F. Benitez^c, A. Castaneda Hernandez^c, H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello^c,
J.A. Murillo Quijada^c, A. Sehrawat^c, L. Valencia Palomo^c

Universidad de Sonora (UNISON), Hermosillo, Mexico

G. Ayala^d, H. Castilla-Valdez^d, I. Heredia-De La Cruz^{d,58}, R. Lopez-Fernandez^d, C.A. Mondragon Herrera,
D.A. Perez Navarro^d, A. Sánchez Hernández^d

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

C. Oropeza Barrera^e, F. Vazquez Valencia^e

Universidad Iberoamericana, Mexico City, Mexico

I. Pedraza^f, H.A. Salazar Ibarguen^f, C. Uribe Estrada^f

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bubanja, J. Mijuskovic^{g,59}, N. Raicevic^g

University of Montenegro, Podgorica, Montenegro

A. Ahmad^h, M.I. Asghar, A. Awais^h, M.I.M. Awan, M. Gul^h, H.R. Hoorani^h, W.A. Khan^h

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

V. Avati, L. Grzankaⁱ, M. Malawskiⁱ

AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

H. Bialkowska^j, M. Bluj^j, B. Boimska^j, M. Górski^j, M. Kazana^j, M. Szleper^j, P. Zalewski^j

National Centre for Nuclear Research, Swierk, Poland

K. Bunkowski^k, K. Doroba^k, A. Kalinowski^k, M. Konecki^k, J. Krolikowski^k

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

M. Araujo^l, P. Bargassa^l, D. Bastos^l, A. Boletti^l, P. Faccioli^l, M. Gallinaro^l, J. Hollar^l,
N. Leonardo^l, T. Niknejad^l, M. Pisano^l, J. Seixas^l, J. Varela^l

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

P. Adzic^{m,60}, M. Dordevic^m, P. Milenovic^m, J. Milosevic^m

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

M. Aguilar-Benitez, J. Alcaraz Maestreⁿ, M. Barrio Luna, Cristina F. Bedoyaⁿ, M. Cepedaⁿ, M. Cerradaⁿ,
N. Colinoⁿ, B. De La Cruzⁿ, A. Delgado Perisⁿ, D. Fernández Del Valⁿ, J.P. Fernández Ramosⁿ, J. Flixⁿ,
M.C. Fouzⁿ, O. Gonzalez Lopezⁿ, S. Goy Lopezⁿ, J.M. Hernandezⁿ, M.I. Josaⁿ, J. León Holgadoⁿ,
D. Moranⁿ, C. Perez Dengraⁿ, A. Pérez-Calero Yzquierdoⁿ, J. Puerta Pelayoⁿ, I. Redondoⁿ,
D.D. Redondo Ferreroⁿ, L. Romero, S. Sánchez Navasⁿ, J. Sastreⁿ, L. Urda Gómezⁿ, J. Vazquez Escobarⁿ,
C. Willmott

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J.F. de Trocóniz 

Universidad Autónoma de Madrid, Madrid, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero ,
J.R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza ,
A. Soto Rodríguez , A. Trapote , C. Vico Villalba 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez ,
C. Fernandez Madrazo , A. García Alonso, G. Gomez , C. Lasasa García , C. Martinez Rivero ,
P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , J. Piedra Gomez , C. Prieels,
L. Scodellaro , I. Vila , J.M. Vizan Garcia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

M.K. Jayananda , B. Kailasapathy ⁶¹, D.U.J. Sonnadara , D.D.C. Wickramaratna 

University of Colombo, Colombo, Sri Lanka

W.G.D. Dharmaratna ⁶², K. Liyanage , N. Perera , N. Wickramage 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

D. Abbaneo , E. Auffray , G. Auzinger , J. Baechler, P. Baillon [†], D. Barney , J. Bendavid ,
A. Bermúdez Martínez , M. Bianco , B. Bilin , A.A. Bin Anuar , A. Bocci , E. Brondolin , C. Caillol ,
T. Camporesi , G. Cerminara , N. Chernyavskaya , S.S. Chhibra , S. Choudhury, M. Cipriani ,
D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson ,
M. Dünser , N. Dupont, F. Fallavollita ⁶³, A. Florent , L. Forthomme , G. Franzoni , W. Funk ,
S. Ghosh , S. Giani, D. Gigi, K. Gill , F. Glege , L. Gouskos , E. Govorkova , M. Haranko ,
J. Hegeman , V. Innocente , T. James , P. Janot , J. Kaspar , J. Kieseler , N. Kratochwil ,
S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri , M. Mannelli ,
A.C. Marini , F. Meijers , S. Mersi , E. Meschi , F. Moortgat , M. Mulders , S. Orfanelli, L. Orsini,
F. Pantaleo , E. Perez, M. Peruzzi , A. Petrilli , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo ,
M. Pitt , H. Qu , T. Quast, D. Rabadý , A. Racz, G. Reales Gutiérrez, M. Rovere , H. Sakulin ,
J. Salfeld-Nebgen , S. Scarfi , M. Selvaggi , A. Sharma , P. Silva , P. Sphicas ⁶⁴, A.G. Stahl Leiton ,
S. Summers , K. Tatar , D. Treille , P. Tropea , A. Tsiros, J. Wanczyk ⁶⁵, K.A. Wozniak , W.D. Zeuner

CERN, European Organization for Nuclear Research, Geneva, Switzerland

L. Caminada ⁶⁶, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli ,
D. Kotlinski , C. Lange , M. Missiroli ⁶⁶, L. Noehte ⁶⁶, T. Rohe 

Paul Scherrer Institut, Villigen, Switzerland

T.K. Aarrestad , K. Androsov ⁶⁵, M. Backhaus , A. Calandri , K. Datta , A. De Cosa , G. Dissertori ,
M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , T.A. Gómez Espinosa ,
C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , L. Marchese , C. Martin Perez ,
A. Mascellani ⁶⁵, F. Nessi-Tedaldi , J. Niedziela , F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti ,
M. Reichmann , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, D.A. Sanz Becerra ,
R. Seidita , J. Steggemann ⁶⁵, D. Valsecchi , R. Wallny 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

C. Amsler^{ID,67}, P. Bärttschi^{ID}, C. Botta^{ID}, D. Brzhechko, M.F. Canelli^{ID}, K. Cormier^{ID}, A. De Wit^{ID},
R. Del Burgo, J.K. Heikkilä^{ID}, M. Huwiler^{ID}, W. Jin^{ID}, A. Jofrehei^{ID}, B. Kilminster^{ID}, S. Leontsinis^{ID},
S.P. Liechti^{ID}, A. Macchiolo^{ID}, P. Meiring^{ID}, V.M. Mikuni^{ID}, U. Molinatti^{ID}, I. Neutelings^{ID}, A. Reimers^{ID},
P. Robmann, S. Sanchez Cruz^{ID}, K. Schweiger^{ID}, M. Senger^{ID}, Y. Takahashi^{ID}

Universität Zürich, Zurich, Switzerland

C. Adloff⁶⁸, C.M. Kuo, W. Lin, P.K. Rout^{ID}, P.C. Tiwari^{ID,41}, S.S. Yu^{ID}

National Central University, Chung-Li, Taiwan

L. Ceard, Y. Chao^{ID}, K.F. Chen^{ID}, P.s. Chen, H. Cheng^{ID}, W.-S. Hou^{ID}, R. Khurana, G. Kole^{ID}, Y.y. Li^{ID},
R.-S. Lu^{ID}, E. Paganis^{ID}, A. Psallidas, A. Steen^{ID}, H.y. Wu, E. Yazgan^{ID}

National Taiwan University (NTU), Taipei, Taiwan

C. Asawatangtrakuldee^{ID}, N. Srimanobhas^{ID}, V. Wachirapusanand^{ID}

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

D. Agyel^{ID}, F. Boran^{ID}, Z.S. Demiroglu^{ID}, F. Dolek^{ID}, I. Dumanoglu^{ID,69}, E. Eskut^{ID}, Y. Guler^{ID,70},
E. Gurpinar Guler^{ID,70}, C. Isik^{ID}, O. Kara, A. Kayis Topaksu^{ID}, U. Kiminsu^{ID}, G. Onengut^{ID}, K. Ozdemir^{ID,71},
A. Polatoz^{ID}, A.E. Simsek^{ID}, B. Tali^{ID,72}, U.G. Tok^{ID}, S. Turkcapar^{ID}, E. Usan^{ID}, I.S. Zorbakir^{ID}

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

G. Karapinar⁷³, K. Ocalan^{ID,74}, M. Yalvac^{ID,75}

Middle East Technical University, Physics Department, Ankara, Turkey

B. Akgun^{ID}, I.O. Atakisi^{ID}, E. Gülmez^{ID}, M. Kaya^{ID,76}, O. Kaya^{ID,77}, S. Tekten^{ID,78}

Bogazici University, Istanbul, Turkey

A. Cakir^{ID}, K. Cankocak^{ID,69}, Y. Komurcu^{ID}, S. Sen^{ID,79}

Istanbul Technical University, Istanbul, Turkey

O. Aydilek^{ID}, S. Cerci^{ID,72}, B. Hacisahinoglu^{ID}, I. Hos^{ID,80}, B. Isildak^{ID,81}, B. Kaynak^{ID}, S. Ozkorucuklu^{ID},
C. Simsek^{ID}, D. Sunar Cerci^{ID,72}

Istanbul University, Istanbul, Turkey

B. Grynyov^{ID}

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

L. Levchuk^{ID}

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

D. Anthony^{ID}, J.J. Brooke^{ID}, A. Bundock^{ID}, E. Clement^{ID}, D. Cussans^{ID}, H. Flacher^{ID}, M. Glowacki,
J. Goldstein^{ID}, H.F. Heath^{ID}, L. Kreczko^{ID}, B. Krikler^{ID}, S. Paramesvaran^{ID}, S. Seif El Nasr-Storey,
V.J. Smith^{ID}, N. Stylianou^{ID,82}, K. Walkingshaw Pass, R. White^{ID}

University of Bristol, Bristol, United Kingdom

A.H. Ball, K.W. Bell^{ID}, A. Belyaev^{ID,83}, C. Brew^{ID}, R.M. Brown^{ID}, D.J.A. Cockerill^{ID}, C. Cooke^{ID}, K.V. Ellis,
K. Harder^{ID}, S. Harper^{ID}, M.-L. Holmberg^{ID,84}, Sh. Jain^{ID}, J. Linacre^{ID}, K. Manolopoulos, D.M. Newbold^{ID},

E. Olaiya, D. Petyt^{ID}, T. Reis^{ID}, G. Salvi^{ID}, T. Schuh, C.H. Shepherd-Themistocleous^{ID}, I.R. Tomalin^{ID},
T. Williams^{ID}

Rutherford Appleton Laboratory, Didcot, United Kingdom

R. Bainbridge^{ID}, P. Bloch^{ID}, S. Bonomally, J. Borg^{ID}, C.E. Brown^{ID}, O. Buchmuller, V. Cacchio,
C.A. Carrillo Montoya^{ID}, V. Cepaitis^{ID}, G.S. Chahal^{ID,85}, D. Colling^{ID}, J.S. Dancu, P. Dauncey^{ID}, G. Davies^{ID},
J. Davies, M. Della Negra^{ID}, S. Fayer, G. Fedi^{ID}, G. Hall^{ID}, M.H. Hassanshahi^{ID}, A. Howard, G. Iles^{ID},
J. Langford^{ID}, L. Lyons^{ID}, A.-M. Magnan^{ID}, S. Malik, A. Martelli^{ID}, M. Mieskolainen^{ID}, D.G. Monk^{ID},
J. Nash^{ID,86}, M. Pesaresi, B.C. Radburn-Smith^{ID}, D.M. Raymond, A. Richards, A. Rose^{ID}, E. Scott^{ID}, C. Seez^{ID},
R. Shukla^{ID}, A. Tapper^{ID}, K. Uchida^{ID}, G.P. Uttley^{ID}, L.H. Vage, T. Virdee^{ID,29}, M. Vojinovic^{ID}, N. Wardle^{ID},
S.N. Webb^{ID}, D. Winterbottom^{ID}

Imperial College, London, United Kingdom

K. Coldham, J.E. Cole^{ID}, A. Khan, P. Kyberd^{ID}, I.D. Reid^{ID}

Brunel University, Uxbridge, United Kingdom

S. Abdullin^{ID}, A. Brinkerhoff^{ID}, B. Caraway^{ID}, J. Dittmann^{ID}, K. Hatakeyama^{ID}, A.R. Kanuganti^{ID},
B. McMaster^{ID}, M. Saunders^{ID}, S. Sawant^{ID}, C. Sutantawibul^{ID}, M. Toms^{ID}, J. Wilson^{ID}

Baylor University, Waco, TX, USA

R. Bartek^{ID}, A. Dominguez^{ID}, C. Huerta Escamilla, R. Uniyal^{ID}, A.M. Vargas Hernandez^{ID}

Catholic University of America, Washington, DC, USA

R. Chudasama^{ID}, S.I. Cooper^{ID}, D. Di Croce^{ID}, S.V. Gleyzer^{ID}, C. Henderson^{ID}, C.U. Perez^{ID}, P. Rumerio^{ID,87},
C. West^{ID}

The University of Alabama, Tuscaloosa, AL, USA

A. Akpinar^{ID}, A. Albert^{ID}, D. Arcaro^{ID}, C. Cosby^{ID}, Z. Demiragli^{ID}, C. Erice^{ID}, E. Fontanesi^{ID}, D. Gastler^{ID},
S. May^{ID}, J. Rohlf^{ID}, K. Salyer^{ID}, D. Sperka^{ID}, D. Spitzbart^{ID}, I. Suarez^{ID}, A. Tsatsos^{ID}, S. Yuan^{ID}

Boston University, Boston, MA, USA

G. Benelli^{ID}, B. Burke^{ID}, X. Coubez²⁴, D. Cutts^{ID}, M. Hadley^{ID}, U. Heintz^{ID}, J.M. Hogan^{ID,88}, T. Kwon^{ID},
G. Landsberg^{ID}, K.T. Lau^{ID}, D. Li^{ID}, J. Luo^{ID}, M. Narain^{ID}, N. Pervan^{ID}, S. Sagir^{ID,89}, F. Simpson^{ID},
E. Usai^{ID}, W.Y. Wong, X. Yan^{ID}, D. Yu^{ID}, W. Zhang

Brown University, Providence, RI, USA

S. Abbott^{ID}, J. Bonilla^{ID}, C. Brainerd^{ID}, R. Breedon^{ID}, M. Calderon De La Barca Sanchez^{ID}, M. Chertok^{ID},
J. Conway^{ID}, P.T. Cox^{ID}, R. Erbacher^{ID}, G. Haza^{ID}, F. Jensen^{ID}, O. Kukral^{ID}, G. Mocellin^{ID}, M. Mulhearn^{ID},
D. Pellett^{ID}, B. Regnery^{ID}, Y. Yao^{ID}, F. Zhang^{ID}

University of California, Davis, Davis, CA, USA

M. Bachtis^{ID}, R. Cousins^{ID}, A. Datta^{ID}, J. Hauser^{ID}, M. Ignatenko^{ID}, M.A. Iqbal^{ID}, T. Lam^{ID}, E. Manca^{ID},
W.A. Nash^{ID}, D. Saltzberg^{ID}, B. Stone^{ID}, V. Valuev^{ID}

University of California, Los Angeles, CA, USA

R. Clare^{ID}, J.W. Gary^{ID}, M. Gordon, G. Hanson^{ID}, O.R. Long^{ID}, N. Manganello^{ID}, W. Si^{ID}, S. Wimpenny^{ID}

University of California, Riverside, Riverside, CA, USA

J.G. Branson^{ID}, S. Cittolin^{ID}, S. Cooperstein^{ID}, D. Diaz^{ID}, J. Duarte^{ID}, R. Gerosa^{ID}, L. Giannini^{ID},
J. Guiang^{ID}, R. Kansal^{ID}, V. Krutelyov^{ID}, R. Lee^{ID}, J. Letts^{ID}, M. Masciovecchio^{ID}, F. Mokhtar^{ID}, M. Pieri^{ID},
M. Quinnan^{ID}, B.V. Sathia Narayanan^{ID}, V. Sharma^{ID}, M. Tadel^{ID}, E. Vourliotis^{ID}, F. Würthwein^{ID},
Y. Xiang^{ID}, A. Yagil^{ID}

University of California, San Diego, La Jolla, CA, USA

N. Amin, C. Campagnari^{ID}, M. Citron^{ID}, G. Collura^{ID}, A. Dorsett^{ID}, J. Incandela^{ID}, M. Kilpatrick^{ID}, J. Kim^{ID},
A.J. Li^{ID}, P. Masterson^{ID}, H. Mei^{ID}, M. Oshiro^{ID}, J. Richman^{ID}, U. Sarica^{ID}, R. Schmitz^{ID}, F. Setti^{ID},
J. Shephlock^{ID}, P. Siddireddy, D. Stuart^{ID}, S. Wang^{ID}

University of California, Santa Barbara - Department of Physics, Santa Barbara, CA, USA

A. Bornheim^{ID}, O. Cerri, I. Dutta^{ID}, A. Latorre, J.M. Lawhorn^{ID}, J. Mao^{ID}, H.B. Newman^{ID}, T.Q. Nguyen^{ID},
M. Spiropulu^{ID}, J.R. Vlimant^{ID}, C. Wang^{ID}, S. Xie^{ID}, R.Y. Zhu^{ID}

California Institute of Technology, Pasadena, CA, USA

J. Alison^{ID}, S. An^{ID}, M.B. Andrews^{ID}, P. Bryant^{ID}, V. Dutta^{ID}, T. Ferguson^{ID}, A. Harilal^{ID}, C. Liu^{ID},
T. Mudholkar^{ID}, S. Murthy^{ID}, M. Paulini^{ID}, A. Roberts^{ID}, A. Sanchez^{ID}, W. Terrill^{ID}

Carnegie Mellon University, Pittsburgh, PA, USA

J.P. Cumalat^{ID}, W.T. Ford^{ID}, A. Hassani^{ID}, G. Karathanasis^{ID}, E. MacDonald, F. Marini^{ID}, A. Perloff^{ID},
C. Savard^{ID}, N. Schonbeck^{ID}, K. Stenson^{ID}, K.A. Ulmer^{ID}, S.R. Wagner^{ID}, N. Zipper^{ID}

University of Colorado Boulder, Boulder, CO, USA

J. Alexander^{ID}, S. Bright-Thonney^{ID}, X. Chen^{ID}, D.J. Cranshaw^{ID}, J. Fan^{ID}, X. Fan^{ID}, D. Gadkari^{ID},
S. Hogan^{ID}, J. Monroy^{ID}, J.R. Patterson^{ID}, J. Reichert^{ID}, M. Reid^{ID}, A. Ryd^{ID}, J. Thom^{ID}, P. Wittich^{ID},
R. Zou^{ID}

Cornell University, Ithaca, NY, USA

M. Albrow^{ID}, M. Alyari^{ID}, G. Apollinari^{ID}, A. Apresyan^{ID}, L.A.T. Bauerdick^{ID}, D. Berry^{ID}, J. Berryhill^{ID},
P.C. Bhat^{ID}, K. Burkett^{ID}, J.N. Butler^{ID}, A. Canepa^{ID}, G.B. Cerati^{ID}, H.W.K. Cheung^{ID}, F. Chlebana^{ID},
K.F. Di Petrillo^{ID}, J. Dickinson^{ID}, V.D. Elvira^{ID}, Y. Feng^{ID}, J. Freeman^{ID}, A. Gandrakota^{ID}, Z. Gecse^{ID},
L. Gray^{ID}, D. Green, S. Grünendahl^{ID}, D. Guerrero^{ID}, O. Gutsche^{ID}, R.M. Harris^{ID}, R. Heller^{ID},
T.C. Herwig^{ID}, J. Hirschauer^{ID}, L. Horyn^{ID}, B. Jayatilaka^{ID}, S. Jindariani^{ID}, M. Johnson^{ID}, U. Joshi^{ID},
T. Klijnsma^{ID}, B. Klima^{ID}, K.H.M. Kwok^{ID}, S. Lammel^{ID}, D. Lincoln^{ID}, R. Lipton^{ID}, T. Liu^{ID}, C. Madrid^{ID},
K. Maeshima^{ID}, C. Mantilla^{ID}, D. Mason^{ID}, P. McBride^{ID}, P. Merkel^{ID}, S. Mrenna^{ID}, S. Nahn^{ID},
J. Ngadiuba^{ID}, D. Noonan^{ID}, S. Norberg, V. Papadimitriou^{ID}, N. Pastika^{ID}, K. Pedro^{ID}, C. Pena^{ID,90},
F. Ravera^{ID}, A. Reinsvold Hall^{ID,91}, L. Ristori^{ID}, E. Sexton-Kennedy^{ID}, N. Smith^{ID}, A. Soha^{ID}, L. Spiegel^{ID},
S. Stoynev^{ID}, J. Strait^{ID}, L. Taylor^{ID}, S. Tkaczyk^{ID}, N.V. Tran^{ID}, L. Uplegger^{ID}, E.W. Vaandering^{ID}, I. Zoi^{ID}

Fermi National Accelerator Laboratory, Batavia, IL, USA

P. Avery^{ID}, D. Bourilkov^{ID}, L. Cadamuro^{ID}, P. Chang^{ID}, V. Cherepanov^{ID}, R.D. Field, E. Koenig^{ID},
M. Kolosova^{ID}, J. Konigsberg^{ID}, A. Korytov^{ID}, E. Kuznetsova^{ID}, K.H. Lo, K. Matchev^{ID}, N. Menendez^{ID},
G. Mitselmakher^{ID}, A. Muthirakalayil Madhu^{ID}, N. Rawal^{ID}, D. Rosenzweig^{ID}, S. Rosenzweig^{ID}, K. Shi^{ID},
J. Wang^{ID}, Z. Wu^{ID}

University of Florida, Gainesville, FL, USA

T. Adams^{ID}, A. Askew^{ID}, N. Bower^{ID}, R. Habibullah^{ID}, V. Hagopian^{ID}, T. Kolberg^{ID}, G. Martinez,
H. Prosper^{ID}, O. Viazlo^{ID}, M. Wulansatiti^{ID}, R. Yohay^{ID}, J. Zhang

Florida State University, Tallahassee, FL, USA

M.M. Baarmand^{ID}, S. Butalla^{ID}, T. Elkafrawy^{ID,16}, M. Hohlmann^{ID}, R. Kumar Verma^{ID}, M. Rahmani,
F. Yumiceva^{ID}

Florida Institute of Technology, Melbourne, FL, USA

M.R. Adams^{ID}, R. Cavanaugh^{ID}, S. Dittmer^{ID}, O. Evdokimov^{ID}, C.E. Gerber^{ID}, D.J. Hofman^{ID}, D.S. Lemos^{ID},
A.H. Merit^{ID}, C. Mills^{ID}, G. Oh^{ID}, T. Roy^{ID}, S. Rudrabhatla^{ID}, M.B. Tonjes^{ID}, N. Varelas^{ID}, X. Wang^{ID},
Z. Ye^{ID}, J. Yoo^{ID}

University of Illinois Chicago, Chicago, USA

M. Alhusseini^{ID}, K. Dilsiz^{ID,92}, L. Emediato^{ID}, G. Karaman^{ID}, O.K. Köseyan^{ID}, J.-P. Merlo,
A. Mestvirishvili^{ID,93}, J. Nachtman^{ID}, O. Neogi, H. Ogul^{ID,94}, Y. Onel^{ID}, A. Penzo^{ID}, C. Snyder, E. Tiras^{ID,95}

The University of Iowa, Iowa City, IA, USA

O. Amram^{ID}, B. Blumenfeld^{ID}, L. Corcodilos^{ID}, J. Davis^{ID}, A.V. Gritsan^{ID}, S. Kyriacou^{ID}, P. Maksimovic^{ID},
J. Roskes^{ID}, S. Sekhar^{ID}, M. Swartz^{ID}, T.Á. Vámi^{ID}

Johns Hopkins University, Baltimore, MD, USA

A. Abreu^{ID}, L.F. Alcerro Alcerro^{ID}, J. Anguiano^{ID}, P. Baringer^{ID}, A. Bean^{ID}, Z. Flowers^{ID}, J. King^{ID},
G. Krintiras^{ID}, M. Lazarovits^{ID}, C. Le Mahieu^{ID}, C. Lindsey, J. Marquez^{ID}, N. Minafra^{ID}, M. Murray^{ID},
M. Nickel^{ID}, C. Rogan^{ID}, C. Royon^{ID}, R. Salvatico^{ID}, S. Sanders^{ID}, C. Smith^{ID}, Q. Wang^{ID}, G. Wilson^{ID}

The University of Kansas, Lawrence, KS, USA

B. Allmond^{ID}, S. Duric, A. Ivanov^{ID}, K. Kaadze^{ID}, A. Kalogeropoulos^{ID}, D. Kim, Y. Maravin^{ID}, T. Mitchell,
A. Modak, K. Nam, D. Roy^{ID}

Kansas State University, Manhattan, KS, USA

F. Rebassoo^{ID}, D. Wright^{ID}

Lawrence Livermore National Laboratory, Livermore, CA, USA

E. Adams^{ID}, A. Baden^{ID}, O. Baron, A. Belloni^{ID}, A. Bethani^{ID}, S.C. Eno^{ID}, N.J. Hadley^{ID}, S. Jabeen^{ID},
R.G. Kellogg^{ID}, T. Koeth^{ID}, Y. Lai^{ID}, S. Lascio^{ID}, A.C. Mignerey^{ID}, S. Nabili^{ID}, C. Palmer^{ID},
C. Papageorgakis^{ID}, L. Wang^{ID}, K. Wong^{ID}

University of Maryland, College Park, MD, USA

W. Busza^{ID}, I.A. Cali^{ID}, Y. Chen^{ID}, M. D'Alfonso^{ID}, J. Eysermans^{ID}, C. Freer^{ID}, G. Gomez-Ceballos^{ID},
M. Goncharov, P. Harris, M. Hu^{ID}, D. Kovalskyi^{ID}, J. Krupa^{ID}, Y.-J. Lee^{ID}, K. Long^{ID}, C. Mironov^{ID},
C. Paus^{ID}, D. Rankin^{ID}, C. Roland^{ID}, G. Roland^{ID}, Z. Shi^{ID}, G.S.F. Stephans^{ID}, J. Wang, Z. Wang^{ID},
B. Wyslouch^{ID}, T.J. Yang^{ID}

Massachusetts Institute of Technology, Cambridge, MA, USA

R.M. Chatterjee, B. Crossman^{ID}, J. Hiltbrand^{ID}, B.M. Joshi^{ID}, C. Kapsiak^{ID}, M. Krohn^{ID}, Y. Kubota^{ID},
D. Mahon^{ID}, J. Mans^{ID}, M. Revering^{ID}, R. Rusack^{ID}, R. Saradhy^{ID}, N. Schroeder^{ID}, N. Strobbe^{ID},
M.A. Wadud^{ID}

University of Minnesota, Minneapolis, MN, USA

L.M. Cremaldi 

University of Mississippi, Oxford, MS, USA

K. Bloom , M. Bryson , D.R. Claes , C. Fangmeier , L. Finco , F. Golf , C. Joo , R. Kamalieddin ,
 I. Kravchenko , I. Reed , J.E. Siado , G.R. Snow , W. Tabb , A. Wightman , F. Yan ,
 A.G. Zecchinelli 

University of Nebraska-Lincoln, Lincoln, NE, USA

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , C. McLean , M. Morris ,
 D. Nguyen , J. Pekkanen , S. Rappoccio , A. Williams 

State University of New York at Buffalo, Buffalo, NY, USA

G. Alverson , E. Barberis , Y. Haddad , Y. Han , A. Krishna , J. Li , J. Lidrych , G. Madigan ,
 B. Marzocchi , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari ,
 A. Tishelman-Charny , T. Wamorkar , B. Wang , A. Wisecarver , D. Wood 

Northeastern University, Boston, MA, USA

S. Bhattacharya , J. Bueghly , Z. Chen , A. Gilbert , K.A. Hahn , Y. Liu , N. Odell , M.H. Schmitt ,
 M. Velasco 

Northwestern University, Evanston, IL, USA

R. Band , R. Bucci , M. Cremonesi , A. Das , R. Goldouzian , M. Hildreth , K. Hurtado Anampa ,
 C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli , I. Mcalister ,
 T. McCauley , C. Mcgrady , K. Mohrman , C. Moore , Y. Musienko ,¹² R. Ruchti , A. Townsend ,
 M. Wayne , H. Yockey , M. Zarucki , L. Zygala 

University of Notre Dame, Notre Dame, IN, USA

B. Bylsma , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , A. Lesauvage , M. Nunez Ornelas , K. Wei ,
 B.L. Winer , B.R. Yates 

The Ohio State University, Columbus, OH, USA

F.M. Addesa , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich ,
 S. Higginbotham , G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen ,
 D. Stickland , C. Tully 

Princeton University, Princeton, NJ, USA

S. Malik 

University of Puerto Rico, Mayaguez, PR, USA

A.S. Bakshi , V.E. Barnes , R. Chawla , S. Das , L. Gutay , M. Jones , A.W. Jung , D. Kondratyev ,
 A.M. Koshy , M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , A. Purohit ,
 J.F. Schulte , M. Stojanovic , J. Thieman , F. Wang , R. Xiao , W. Xie 

Purdue University, West Lafayette, IN, USA

J. Dolen , N. Parashar 

Purdue University Northwest, Hammond, IN, USA

D. Acosta^{ID}, A. Baty^{ID}, T. Carnahan^{ID}, S. Dildick^{ID}, K.M. Ecklund^{ID}, P.J. Fernández Manteca^{ID}, S. Freed,
P. Gardner, F.J.M. Geurts^{ID}, A. Kumar^{ID}, W. Li^{ID}, B.P. Padley^{ID}, R. Redjimi, J. Rotter^{ID}, S. Yang^{ID},
E. Yigitbasi^{ID}, Y. Zhang^{ID}

Rice University, Houston, TX, USA

A. Bodek^{ID}, P. de Barbaro^{ID}, R. Demina^{ID}, J.L. Dulemba^{ID}, C. Fallon, A. Garcia-Bellido^{ID}, O. Hindrichs^{ID},
A. Khukhunaishvili^{ID}, P. Parygin^{ID}, E. Popova^{ID}, R. Taus^{ID}, G.P. Van Onsem^{ID}

University of Rochester, Rochester, NY, USA

K. Goulios^{ID}

The Rockefeller University, New York, NY, USA

B. Chiarito, J.P. Chou^{ID}, Y. Gershtein^{ID}, E. Halkiadakis^{ID}, A. Hart^{ID}, M. Heindl^{ID}, D. Jaroslawski^{ID},
O. Karacheban^{ID,27}, I. Laflotte^{ID}, A. Lath^{ID}, R. Montalvo, K. Nash, M. Osherson^{ID}, H. Routray^{ID}, S. Salur^{ID},
S. Schnetzer, S. Somalwar^{ID}, R. Stone^{ID}, S.A. Thayil^{ID}, S. Thomas, H. Wang^{ID}

Rutgers, The State University of New Jersey, Piscataway, NJ, USA

H. Acharya, A.G. Delannoy^{ID}, S. Fiorendi^{ID}, T. Holmes^{ID}, E. Nibigira^{ID}, S. Spanier^{ID}

University of Tennessee, Knoxville, TN, USA

O. Bouhali^{ID,96}, M. Dalchenko^{ID}, A. Delgado^{ID}, R. Eusebi^{ID}, J. Gilmore^{ID}, T. Huang^{ID}, T. Kamon^{ID,97},
H. Kim^{ID}, S. Luo^{ID}, S. Malhotra, R. Mueller^{ID}, D. Overton^{ID}, D. Rathjens^{ID}, A. Safonov^{ID}

Texas A&M University, College Station, TX, USA

N. Akchurin^{ID}, J. Damgov^{ID}, V. Hegde^{ID}, K. Lamichhane^{ID}, S.W. Lee^{ID}, T. Mengke, S. Muthumuni^{ID},
T. Peltola^{ID}, I. Volobouev^{ID}, A. Whitbeck^{ID}

Texas Tech University, Lubbock, TX, USA

E. Appelt^{ID}, S. Greene, A. Gurrola^{ID}, W. Johns^{ID}, A. Melo^{ID}, F. Romeo^{ID}, P. Sheldon^{ID}, S. Tuo^{ID},
J. Velkovska^{ID}, J. Viinikainen^{ID}

Vanderbilt University, Nashville, TN, USA

B. Cardwell^{ID}, B. Cox^{ID}, G. Cummings^{ID}, J. Hakala^{ID}, R. Hirosky^{ID}, A. Ledovskoy^{ID}, A. Li^{ID}, C. Neu^{ID},
C.E. Perez Lara^{ID}

University of Virginia, Charlottesville, VA, USA

P.E. Karchin^{ID}

Wayne State University, Detroit, MI, USA

A. Aravind, S. Banerjee^{ID}, K. Black^{ID}, T. Bose^{ID}, S. Dasu^{ID}, I. De Bruyn^{ID}, P. Everaerts^{ID}, C. Galloni, H. He^{ID},
M. Herndon^{ID}, A. Herve^{ID}, C.K. Koraka^{ID}, A. Lanaro, R. Loveless^{ID}, J. Madhusudanan Sreekala^{ID},
A. Mallampalli^{ID}, A. Mohammadi^{ID}, S. Mondal, G. Parida^{ID}, D. Pinna, A. Savin, V. Shang^{ID}, V. Sharma^{ID},
W.H. Smith^{ID}, D. Teague, H.F. Tsoi^{ID}, W. Vetens^{ID}, A. Warden^{ID}

University of Wisconsin - Madison, Madison, WI, USA

S. Afanasiev^{ID}, V. Andreev^{ID}, Yu. Andreev^{ID}, T. Aushev^{ID}, M. Azarkin^{ID}, A. Babaev^{ID}, A. Belyaev^{ID},
V. Blinov^{ID,98}, E. Boos^{ID}, V. Borshch^{ID}, D. Budkouski^{ID}, M. Chadeeva^{ID,98}, V. Chekhovsky, M. Danilov^{ID,98},
A. Dermenev^{ID}, T. Dimova^{ID,98}, I. Dremin^{ID}, V. Epshteyn^{ID}, A. Ershov^{ID}, G. Gavrilo^{ID}, V. Gavrilo^{ID},

S. Gninenko^{1b}, V. Golovtsov^{1b}, N. Golubev^{1b}, I. Golutvin^{1b}, I. Gorbunov^{1b}, A. Gribushin^{1b}, Y. Ivanov^{1b},
V. Kachanov^{1b}, A. Kaminskiy^{1b}, L. Kardapoltsev^{1b,98}, V. Karjavine^{1b}, A. Karneyeu^{1b}, L. Khein, V. Kim^{1b,98},
M. Kirakosyan, D. Kirpichnikov^{1b}, M. Kirsanov^{1b}, O. Kodolova^{1b,99}, D. Konstantinov^{1b}, V. Korenkov^{1b},
V. Korotkikh, A. Kozyrev^{1b,98}, N. Krasnikov^{1b}, A. Lanev^{1b}, P. Levchenko^{1b}, A. Litomin, N. Lychkovskaya^{1b},
V. Makarenko^{1b}, A. Malakhov^{1b}, V. Matveev^{1b,98}, V. Murzin^{1b}, A. Nikitenko^{1b,100,99}, S. Obraztsov^{1b},
I. Ovtin^{1b,98}, V. Palichik^{1b}, V. Perelygin^{1b}, S. Petrushanko^{1b}, V. Popov, O. Radchenko^{1b,98}, V. Rusinov,
M. Savina^{1b}, V. Savrin^{1b}, D. Selivanova^{1b}, V. Shalaev^{1b}, S. Shmatov^{1b}, S. Shulha^{1b}, Y. Skovpen^{1b,98},
S. Slabospitskii^{1b}, V. Smirnov^{1b}, A. Snigirev^{1b}, D. Sosnov^{1b}, V. Sulimov^{1b}, E. Tcherniaev^{1b}, A. Terkulov^{1b},
O. Teryaev^{1b}, I. Tlisova^{1b}, A. Toropin^{1b}, L. Uvarov^{1b}, A. Uzunian^{1b}, I. Vardanyan^{1b}, A. Vorobyev[†],
N. Voytishin^{1b}, B.S. Yuldashev¹⁰¹, A. Zarubin^{1b}, I. Zhizhin^{1b}, A. Zhokin^{1b}

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

[†] Deceased.

¹ Also at Yerevan State University, Yerevan, Armenia.

² Also at TU Wien, Vienna, Austria.

³ Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt.

⁴ Also at Université Libre de Bruxelles, Bruxelles, Belgium.

⁵ Also at Universidade Estadual de Campinas, Campinas, Brazil.

⁶ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil.

⁷ Also at UFMS, Nova Andradina, Brazil.

⁸ Also at University of Chinese Academy of Sciences, Beijing, China.

⁹ Also at Nanjing Normal University, Nanjing, China.

¹⁰ Now at The University of Iowa, Iowa City, Iowa, USA.

¹¹ Also at University of Chinese Academy of Sciences, Beijing, China.

¹² Also at an institute or an international laboratory covered by a cooperation agreement with CERN.

¹³ Also at Helwan University, Cairo, Egypt.

¹⁴ Now at Zewail City of Science and Technology, Zewail, Egypt.

¹⁵ Also at British University in Egypt, Cairo, Egypt.

¹⁶ Now at Ain Shams University, Cairo, Egypt.

¹⁷ Also at Purdue University, West Lafayette, Indiana, USA.

¹⁸ Also at Université de Haute Alsace, Mulhouse, France.

¹⁹ Also at Department of Physics, Tsinghua University, Beijing, China.

²⁰ Also at Ilia State University, Tbilisi, Georgia.

²¹ Also at The University of the State of Amazonas, Manaus, Brazil.

²² Also at Erzincan Binali Yildirim University, Erzincan, Turkey.

²³ Also at University of Hamburg, Hamburg, Germany.

²⁴ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany.

²⁵ Also at Isfahan University of Technology, Isfahan, Iran.

²⁶ Also at Bergische University Wuppertal (BUW), Wuppertal, Germany.

²⁷ Also at Brandenburg University of Technology, Cottbus, Germany.

²⁸ Also at Forschungszentrum Jülich, Juelich, Germany.

²⁹ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland.

³⁰ Also at Institute of Physics, University of Debrecen, Debrecen, Hungary.

³¹ Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary.

³² Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania.

³³ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt.

³⁴ Also at Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary.

³⁵ Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary.

³⁶ Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary.

³⁷ Also at Punjab Agricultural University, Ludhiana, India.

³⁸ Also at UPES - University of Petroleum and Energy Studies, Dehradun, India.

³⁹ Also at University of Visva-Bharati, Santiniketan, India.

⁴⁰ Also at University of Hyderabad, Hyderabad, India.

⁴¹ Also at Indian Institute of Science (IISc), Bangalore, India.

⁴² Also at Indian Institute of Technology (IIT), Mumbai, India.

⁴³ Also at IIT Bhubaneswar, Bhubaneswar, India.

⁴⁴ Also at Institute of Physics, Bhubaneswar, India.

⁴⁵ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany.

⁴⁶ Now at Department of Physics, Isfahan University of Technology, Isfahan, Iran.

⁴⁷ Also at Sharif University of Technology, Tehran, Iran.

⁴⁸ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran.

- ⁴⁹ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy.
- ⁵⁰ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy.
- ⁵¹ Also at Università degli Studi Guglielmo Marconi, Roma, Italy.
- ⁵² Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy.
- ⁵³ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA.
- ⁵⁴ Also at Laboratori Nazionali di Legnaro dell'INFN, Legnaro, Italy.
- ⁵⁵ Also at Università di Napoli 'Federico II', Napoli, Italy.
- ⁵⁶ Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy.
- ⁵⁷ Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia.
- ⁵⁸ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico.
- ⁵⁹ Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France.
- ⁶⁰ Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia.
- ⁶¹ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka.
- ⁶² Also at Saegis Campus, Nugegoda, Sri Lanka.
- ⁶³ Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy.
- ⁶⁴ Also at National and Kapodistrian University of Athens, Athens, Greece.
- ⁶⁵ Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland.
- ⁶⁶ Also at Universität Zürich, Zurich, Switzerland.
- ⁶⁷ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria.
- ⁶⁸ Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France.
- ⁶⁹ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey.
- ⁷⁰ Also at Konya Technical University, Konya, Turkey.
- ⁷¹ Also at Izmir Bakircay University, Izmir, Turkey.
- ⁷² Also at Adiyaman University, Adiyaman, Turkey.
- ⁷³ Also at Istanbul Gedik University, Istanbul, Turkey.
- ⁷⁴ Also at Necmettin Erbakan University, Konya, Turkey.
- ⁷⁵ Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey.
- ⁷⁶ Also at Marmara University, Istanbul, Turkey.
- ⁷⁷ Also at Milli Savunma University, Istanbul, Turkey.
- ⁷⁸ Also at Kafkas University, Kars, Turkey.
- ⁷⁹ Also at Hacettepe University, Ankara, Turkey.
- ⁸⁰ Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey.
- ⁸¹ Also at Ozyegin University, Istanbul, Turkey.
- ⁸² Also at Vrije Universiteit Brussel, Brussel, Belgium.
- ⁸³ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom.
- ⁸⁴ Also at University of Bristol, Bristol, United Kingdom.
- ⁸⁵ Also at IPPP Durham University, Durham, United Kingdom.
- ⁸⁶ Also at Monash University, Faculty of Science, Clayton, Australia.
- ⁸⁷ Also at Università di Torino, Torino, Italy.
- ⁸⁸ Also at Bethel University, St. Paul, Minnesota, USA.
- ⁸⁹ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey.
- ⁹⁰ Also at California Institute of Technology, Pasadena, California, USA.
- ⁹¹ Also at United States Naval Academy, Annapolis, Maryland, USA.
- ⁹² Also at Bingol University, Bingol, Turkey.
- ⁹³ Also at Georgian Technical University, Tbilisi, Georgia.
- ⁹⁴ Also at Sinop University, Sinop, Turkey.
- ⁹⁵ Also at Erciyes University, Kayseri, Turkey.
- ⁹⁶ Also at Texas A&M University at Qatar, Doha, Qatar.
- ⁹⁷ Also at Kyungpook National University, Daegu, Korea.
- ⁹⁸ Also at another institute or international laboratory covered by a cooperation agreement with CERN.
- ⁹⁹ Also at Yerevan Physics Institute, Yerevan, Armenia.
- ¹⁰⁰ Also at Imperial College, London, United Kingdom.
- ¹⁰¹ Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan.