

A Polyhedral Method for Sparse Systems with Many Positive Solutions*

Frédéric Bihan[†], Francisco Santos[‡], and Pierre-Jean Spaenlehauer[§]

Abstract. We investigate a version of Viro’s method for constructing polynomial systems with many positive solutions, based on regular triangulations of the Newton polytope of the system. The number of positive solutions obtained with our method is governed by the size of the largest *positively decorable* subcomplex of the triangulation. Here, positive decorability is a property that we introduce and which is dual to being a subcomplex of some regular triangulation. Using this duality, we produce large positively decorable subcomplexes of the boundary complexes of cyclic polytopes. As a byproduct, we get new lower bounds, some of them being the best currently known, for the maximal number of positive solutions of polynomial systems with prescribed numbers of monomials and variables. We also study the asymptotics of these numbers and observe a log-concavity property.

Key words. polynomial systems, triangulations, cyclic polytopes

AMS subject classifications. 05, 14

DOI. 10.1137/18M1181912

1. Introduction. Positive solutions of multivariate polynomial systems are central objects in many applications of mathematics, as they often contain meaningful information, e.g., in robotics, optimization, algebraic statistics, the study of multistationarity in chemical reaction networks, etc. In the 1970s, foundational results by Kushnirenko [15], Khovanskii [14], and Bernstein [2] laid the theoretical ground for the study of the algebraic structure of polynomial systems with prescribed conditions on the set of monomials appearing with nonzero coefficients. As a particular case of more general bounds, Khovanskii [14] obtained an upper bound on the number of nondegenerate positive solutions which depends only on the dimension of the problem and on the number of monomials.

More precisely, our main object of interest in this paper is the function $\Xi_{d,k}$, defined as the maximal possible number of nondegenerate solutions in $\mathbb{R}_{>0}^d$ of a polynomial system $f_1 = \cdots = f_d = 0$, where $f_1, \dots, f_d \in \mathbb{R}[X_1, \dots, X_d]$ involve at most $d + k + 1$ monomials with nonzero coefficients. Here, *nondegenerate* means that the Jacobian matrix of the system is

*Received by the editors April 19, 2018; accepted for publication (in revised form) October 4, 2018; published electronically December 18, 2018.

<http://www.siam.org/journals/siaga/2-4/M118191.html>

Funding: The second author’s research was partially supported by grant MTM2014-54207-P of the Spanish Ministry of Science, by the Einstein Foundation Berlin, and, while he was in residence at the Mathematical Sciences Research Institute in Berkeley, California during the Fall 2017 semester, by the Clay Institute and the National Science Foundation (grant DMS-1440140).

[†]Laboratoire de Mathématiques, Université Savoie Mont Blanc, Campus Scientifique, 73376 Le Bourget-du-Lac Cedex, France (frederic.bihan@univ-smb.fr).

[‡]Depto. de Matemáticas, Estadística y Computación, Universidad de Cantabria; 39012 Santander, Spain (francisco.santos@unican.es).

[§]CARAMBA project, INRIA Nancy – Grand Est; Université de Lorraine; CNRS, UMR 7503; LORIA, Nancy, France (pierre-jean.spaienlehauer@inria.fr).

invertible at the solution. Finding sharp bounds for $\Xi_{d,k}$ is a notably hard problem; see [21]. The current knowledge can be briefly summarized as follows (see [5, 6]):

$$\forall d, k > 0, \quad \max((\lfloor k/d \rfloor + 1)^d, (\lfloor d/k \rfloor + 1)^k) \leq \Xi_{d,k} \leq (e^2 + 3)2^{\binom{k}{2}} d^k / 4.$$

Another important and recent lower bound is $\Xi_{2,2} \geq 7$ [8].

In this paper, we introduce a new technique to construct fewnomial systems with many positive roots, based on the notion of *positively decorable* subcomplexes in a regular triangulation of the point configuration given by the exponent vectors of the monomials. Using this method, we obtain new lower bounds for $\Xi_{d,k}$. Combining it with a log-concavity property, we obtain systems which admit asymptotically more positive solutions than previous constructions for a large range of parameters.

Main results. Consider a regular full-dimensional pure simplicial complex Γ supported on a point configuration $\mathcal{A} = \{w_1, \dots, w_n\} \subset \mathbb{Z}^d$, by which we mean that Γ is a pure d -dimensional subcomplex of a regular triangulation of \mathcal{A} (see Definition 4.2 and Proposition 4.3). Consider also a map $\phi: \mathcal{A} \rightarrow \mathbb{R}^d$. This map will be used to construct a polynomial system where the coefficients of the monomial w_i will be obtained from $\phi(w_i)$. We call a facet $\tau = \text{conv}(w_{i_1}, \dots, w_{i_{d+1}})$ of Γ *positively decorated* by ϕ if $\phi(\{w_{i_1}, \dots, w_{i_{d+1}}\})$ positively spans \mathbb{R}^d . We are interested in sparse polynomial systems

$$(1.1) \quad f_1(X_1, \dots, X_d) = \dots = f_d(X_1, \dots, X_d) = 0$$

with real coefficients and support contained in \mathcal{A} : this means that all exponent vectors $w \in \mathbb{Z}^d$ of the monomials X^w appearing with a nonzero coefficient in at least one equation are in \mathcal{A} . Our starting point is the following result.

Theorem A (Theorem 3.4). *There is a choice of coefficients—which can be constructed from the map ϕ —which produces a sparse system supported on \mathcal{A} such that the number of nondegenerate positive solutions of (1.1) is bounded below by the number of facets in Γ which are positively decorated by ϕ .*

This theorem is a version of Viro's method which was used by Sturmfels [23] to construct sparse polynomial systems, all solutions of which are real. Viro's method ([25]; see also [3, 19, 24]) is one of the roots of tropical geometry, and it has been used for constructing real algebraic varieties with interesting topological and combinatorial properties.

We then apply this theorem to the problem of constructing fewnomial systems with many positive solutions. For this we construct large simplicial complexes that are regular and *positively decorable* (that is, all their facets can be positively decorated with a certain ϕ), obtained as subcomplexes of the boundary of cyclic polytopes. Combinatorial techniques allow us to count the simplices of these complexes, which gives us new explicit lower bounds on $\Xi_{d,k}$. More precisely, for all $i, j \in \mathbb{Z}_{>0}$, set

$$F_{i,j} = D_{i,j} + D_{i-1,j-1},$$

where $D_{i,j}$ is the (i, j) th *Delannoy number* [1], defined as

$$(1.2) \quad D_{i,j} := \sum_{\ell=0}^{\min\{i,j\}} \frac{(i+j-\ell)!}{(i-\ell)!(j-\ell)! \ell!} = \sum_{\ell=0}^{\min\{i,j\}} 2^\ell \binom{i}{\ell} \binom{j}{\ell}.$$

Theorem B (Corollary 6.9, Remark 6.10). *For every $i, j \in \mathbb{Z}_{>0}$, we have*

$$\Xi_{2i-1,2j} \geq F_{i,j}, \quad \Xi_{2i-1,2j-1} \geq \frac{j}{i+j} F_{i,j}, \quad \Xi_{2i,2j} \geq \frac{i+1}{i+j+1} F_{i+1,j}, \quad \Xi_{2i,2j-1} \geq 2 F_{i,j-1}.$$

We are then interested in the asymptotics of $\Xi_{d,k}$ for big d and k . One way to make sense of this is the following.

Theorem C (Theorem 2.4). *For all $k, d \in \mathbb{Z}_{>0}$ the limit $\xi_{d,k} := \lim_{n \rightarrow \infty} (\Xi_{dn,kn})^{1/(dn+kn)} \in [1, \infty]$ exists. Moreover, this limit depends only on the ratio d/k and it is bounded from below by $\Xi_{d,k}^{1/(d+k)}$.*

Analyzing the asymptotics of Delannoy numbers leads to the following new lower bound, which also depends only on d/k .

Theorem D (Theorem 7.2, Corollary 7.3). *For all $k, d \in \mathbb{Z}_{>0}$, we have*

$$\xi_{d,k} \geq \left(\frac{\sqrt{d^2 + k^2} + k}{d} \right)^{\frac{d}{2(d+k)}} \left(\frac{\sqrt{d^2 + k^2} + d}{k} \right)^{\frac{k}{2(d+k)}}.$$

This statement allows us to improve the lower bounds on $\xi_{d,k}$ for $0.2434 < d/(d+k) < 0.3659$ and for $0.6342 < d/(d+k) < 0.7565$; see Figure 4. In fact, Theorem 2.4 implies that the limit $\xi_{\alpha,\beta} := \lim_{n \rightarrow \infty} (\Xi_{\alpha n, \beta n})^{1/(\alpha n + \beta n)}$ exists for any positive rational numbers α, β . It is convenient to look at ξ along the segment $\alpha + \beta = 1$. This is no loss of generality since $\xi_{d,k} = \xi_{\alpha, 1-\alpha}$ for $\alpha = d/(d+k)$ and it has the nice property that the function $\xi : \alpha \mapsto \xi_{\alpha, 1-\alpha}$, $\alpha \in (0, 1) \cap \mathbb{Q}$, is log-concave (Proposition 2.5). Therefore, convex hulls of lower bounds for $\log \xi_{\alpha, 1-\alpha}$ also produce lower bounds for this function. With this observation, the methods in this paper improve the previously known lower bounds for $\xi_{d,k}$ for all d, k with $d/(d+k) \in (0.2, 0.5) \cup (0.5, 0.8)$; see Figure 5.

Our bounds also raise some important questions about $\xi_{d,k}$. Notice that log-concavity implies that if ξ is infinite somewhere, then it is infinite everywhere (Corollary 2.6). In light of this, we pose the following question.

Question 1.1. *Is $\xi_{d,k}$ finite for some (equivalently, for every) $d, k > 0$? That is to say, is there a global constant c such that $\Xi_{d,k} \leq c^{k+d}$ for all $k, d \in \mathbb{Z}_{>0}$?*

In fact, we do not know whether $\Xi_{d,k}$ admits a singly exponential upper bound since the best known general upper bound (see (2.1)) is only of type $2^{O(k^2 + k \log d)}$. Compare this to Problem 2.8 in Sturmfels [23] (still open), which asks whether $\Xi_{d,k}$ is polynomial for fixed d . In a sense, Sturmfels' formulation is related to the behavior of $\xi_{\alpha,\beta}$ when $\alpha/\beta \approx 0$, although the answer to it might be positive even if ξ is infinite. (Think, e.g., of $\Xi_{d,k}$ growing as $\min\{d^k, k^d\}$). Our formulation looks at Ξ globally and gives the same role to d and k , which is consistent with Proposition 2.1.

Another intriguing question is whether $\xi_{d,k} = \xi_{k,d}$ or, more strongly, whether $\Xi_{d,k} = \Xi_{k,d}$. This symmetry between d and k holds true for all known lower bounds and exact values, including the lower bounds for $\xi_{d,k}$ obtained with our construction where the symmetry is a consequence of a Gale-type duality between regular and positively decorable complexes (see

Corollary 4.7 and Theorem 4.10). This duality is also instrumental in our proof that the complexes used for Theorem B are positively decorable.

Although not needed for the rest of the paper, we also show that positive decorability is related to two classical properties in topological combinatorics.

Theorem E (Theorem 5.5). *For every pure orientable simplicial complex, one has*

$$\text{balanced} \implies \text{positively decorable} \implies \text{bipartite}.$$

Under certain hypotheses (e.g., for complexes that are simply connected manifolds with or without boundary), the reverse implications also hold (Corollary 5.8).

Organization of the paper. In section 2, we present classical bounds for $\Xi_{d,k}$, introduce the quantity $\xi_{d,k}$, and prove Theorem C, plus the log-concavity property. Section 3 describes the Viro construction used throughout the paper and proves Theorem A. In section 4, we show the duality between positively decorable and regular complexes, and section 5 relates positive decorability to balancedness and bipartiteness. Section 6 contains our main construction, based on cyclic polytopes, and shows the lower bounds stated in Theorem B. This bound is analyzed and compared to previous ones in section 7, where we prove Theorem D. In section 8, we investigate the potential of the proposed method and show that the number of positive solutions that can be produced by this method is inherently limited by the upper bound theorem for polytopes.

2. Preliminaries on $\Xi_{d,k}$. Here, we review what is known about the function $\Xi_{d,k}$, defined as the maximum possible number of positive nondegenerate solutions of d -dimensional systems with $d + k + 1$ monomials. The finiteness of $\Xi_{d,k}$ follows from the work of Khovanskii [14]. The currently best known general upper bound for $\Xi_{d,k}$ for arbitrary d and k is proved by Bihan and Sottile [6]:

$$(2.1) \quad \Xi_{d,k} \leq \frac{e^2 + 3}{4} 2^{\binom{k}{2}} d^k \quad \forall k, d \in \mathbb{Z}_{>0}.$$

The following proposition summarizes what is known about lower bounds of $\Xi_{d,k}$.

Proposition 2.1.

1. $\Xi_{d+d',k+k'} \geq \Xi_{d,k} \Xi_{d',k'}$ for all $d, d', k, k' \in \mathbb{Z}_{>0}$.
2. $\Xi_{1,k} = k + 1$ for all $k \in \mathbb{Z}_{>0}$ (Descartes).
3. $\Xi_{d,1} = d + 1$ for all $d \in \mathbb{Z}_{>0}$ (Bihan [4]).
4. $\Xi_{2,2} \geq 7$ (El Hilany [8]).

Proof. Let $A \subset \mathbb{Z}^d$ and $A' \subset \mathbb{Z}^{d'}$ be supports of systems in d and d' variables with $d + k + 1$ and $d' + k' + 1$ monomials achieving the bounds $\Xi_{d,k}$ and $\Xi_{d',k'}$. Without loss of generality, we can assume that both A and A' contain the origin; indeed, translating the supports amounts to multiplying the whole system by a monomial, which does not affect the number of positive roots. Then $(A \times \{0\}) \cup (\{0\} \times A') \subset \mathbb{Z}^{d+d'}$ has $(d+d') + (k+k') + 1$ points and supports a system (the union of the original systems) with $\Xi_{d,k} \Xi_{d',k'}$ nondegenerate positive solutions (the Cartesian product of the solutions sets of the original systems). Therefore, $\Xi_{d+d',k+k'} \geq \Xi_{d,k} \Xi_{d',k'}$.

The equality $\Xi_{1,k} = k + 1$ comes from the fact that a univariate polynomial with $k + 2$ monomials cannot have more than $k + 1$ positive solutions by Descartes' rule of signs (and the polynomial $\prod_{i=1}^{k+1}(x - i)$ reaches this bound).

Finally, $\Xi_{d,1} = d + 1$ was proved in [4, Thm. A] and $\Xi_{2,2} \geq 7$ has recently been shown by El Hilany [8, Thm. 1.2] using tropical geometry. ■

Remark 2.2. It is known that $\Xi_{d,0} = 1$ (see Proposition 3.3). Moreover, $\Xi_{d,k+1} \geq \Xi_{d,k}$ is obvious (adding one monomial with a very small coefficient does not decrease the number of nondegenerate positive solutions). Then, by setting $\Xi_{0,k} = 1$, part 1 of Proposition 2.1 can be extended to allow zero values for d and k . Consequently, $\Xi_{d',k'} \geq \Xi_{d,k}$ if $d' \geq d$ and $k' \geq k$.

The following consequences of Proposition 2.1 have been observed before. Part 1 comes from a system of univariate polynomials in independent variables, and part 2 was proved by Bihan, Rojas, and Sottile in [5].

Corollary 2.3.

1. If $k_1 + \cdots + k_d = k$ is an integer partition of k , then we have $\Xi_{d,k} \geq \prod_{1 \leq i \leq d} (k_i + 1)$. In particular,

$$(2.2) \quad \Xi_{d,k} \geq (\lfloor k/d \rfloor + 1)^d.$$

2. If $d_1 + \cdots + d_k = d$ is an integer partition of d , then $\Xi_{d,k} \geq \prod_{1 \leq i \leq d} (d_i + 1)$. In particular,

$$(2.3) \quad \Xi_{d,k} \geq (\lfloor d/k \rfloor + 1)^k. \quad \blacksquare$$

Observe that both bounds specialize to

$$(2.4) \quad \Xi_{d,d} \geq 2^d$$

for $k = d$, but a better bound of $\Xi_{2d,2d} \geq 7^d$ follows from parts 1 and 4 of Proposition 2.1.

In section 7, we will be interested in the asymptotics of $\Xi_{d,k}$ for big d and k .

Theorem 2.4. Let $d, k \in \mathbb{Z}_{>0}$. Then the following limit exists:

$$\lim_{n \rightarrow \infty} (\Xi_{dn, kn})^{1/(dn+kn)} \in [1, \infty].$$

Moreover, the limit depends only on the ratio d/k and it is bounded from below by $(\Xi_{d,k})^{1/(d+k)}$.

Proof. For each n , let $a_n := \log(\Xi_{dn, kn})$, so that the limit that we want to compute is $\lim_{n \rightarrow \infty} e^{a_n/(d+k)n}$ and we can instead look at $\lim_{n \rightarrow \infty} (a_n/(d+k)n)$. Since $(a_n)_{n \in \mathbb{Z}_{>0}}$ is increasing (Remark 2.2) and $a_{pn_0} \geq pa_{n_0}$ for every positive integer p (Proposition 2.1), we have $a_n \geq \lfloor \frac{n}{n_0} \rfloor a_{n_0}$ for all $n, n_0 \in \mathbb{Z}_{>0}$. Thus,

$$\liminf_{n \rightarrow \infty} \frac{a_n}{n} \geq \liminf_{n \rightarrow \infty} \left\lfloor \frac{n}{n_0} \right\rfloor \frac{a_{n_0}}{n} = \frac{a_{n_0}}{n_0} \quad \forall n_0 \in \mathbb{Z}_{>0}.$$

Consequently,

$$\liminf_{n \rightarrow \infty} \frac{a_n}{(d+k)n} = \frac{1}{d+k} \liminf_{n \rightarrow \infty} \frac{a_n}{n} \geq \frac{1}{d+k} \frac{a_{n_0}}{n_0} \quad \forall n_0 \in \mathbb{Z}_{>0}.$$

In particular,

$$\liminf_{n \rightarrow \infty} \frac{a_n}{(d+k)n} \geq \sup_{n \in \mathbb{Z}_{>0}} \frac{a_n}{(d+k)n} \geq \limsup_{n \rightarrow \infty} \frac{a_n}{(d+k)n},$$

which implies that the limit exists and equals the supremum. To show that the limit depends only on the ratio d/k , observe that if (d, k) and (d', k') are proportional vectors, then the sequences $(a_n/(d+k)n)_{n \in \mathbb{Z}_{>0}}$ and $(a'_n/(d'+k')n)_{n \in \mathbb{Z}_{>0}}$ (where $a'_n := \log(\Xi_{d'n, k'n})$) have a common subsequence. ■

Note that the statement implies the existence of the limit

$$\xi_{\alpha, \beta} = \lim_{\substack{n \rightarrow \infty \\ \alpha n, \beta n \in \mathbb{Z}}} (\Xi_{\alpha n, \beta n})^{1/(\alpha n + \beta n)} \in [1, \infty]$$

for any positive rational numbers $\alpha, \beta \in \mathbb{Q}_{>0}$ and that for $\alpha = \frac{d}{d+k}$ (where $d, k \in \mathbb{Z}_{>0}$) we have $\xi_{\alpha, 1-\alpha} = \lim_{n \rightarrow \infty} (\Xi_{dn, kn})^{1/(dn+kn)}$. Also, since the limit in Theorem 2.4 depends only on d/k , we only need to consider the function ξ for one point along each ray in the positive orthant. We choose the segment defined by $\alpha + \beta = 1$ because along this segment ξ is log-concave.

Proposition 2.5. *The function $\alpha \mapsto \xi_{\alpha, 1-\alpha}$ is log-concave over $(0, 1) \cap \mathbb{Q}$.*

Proof. For any integer n and any $(\alpha, \beta) \in \mathbb{Q}_{>0}^2$ with $\alpha n, \beta n \in \mathbb{Z}$, let $a_n(\alpha, \beta) = \log(\Xi_{\alpha n, \beta n})$. The statement is that for any $(\alpha, \beta), (\alpha', \beta')$ in $\mathbb{Q}_{>0}^2$ and any $\theta \in [0, 1] \cap \mathbb{Q}$, we have

$$(2.5) \quad \lim_{n \rightarrow \infty} \frac{1}{n} a_n(\theta(\alpha, \beta) + (1-\theta)(\alpha', \beta')) \geq \theta \lim_{n \rightarrow \infty} \frac{1}{n} a_n(\alpha, \beta) + (1-\theta) \lim_{n \rightarrow \infty} \frac{1}{n} a_n(\alpha', \beta').$$

Here and in what follows, only values of n where $\alpha n, \theta \alpha n$, etc., are integers are considered. This is enough since they form an infinite sequence and the limit ξ is independent of the subsequence considered.

Using Proposition 2.1, together with Remark 2.2, we get

$$\begin{aligned} a_n(\theta(\alpha, \beta) + (1-\theta)(\alpha', \beta')) &= \log(\Xi_{\theta \alpha n + (1-\theta)\alpha' n, \theta \beta n + (1-\theta)\beta' n}) \\ &\geq \log(\Xi_{\theta \alpha n, \theta \beta n}) + \log(\Xi_{(1-\theta)\alpha' n, (1-\theta)\beta' n}) \\ &= a_n(\theta \alpha, \theta \beta) + a_n((1-\theta)\alpha', (1-\theta)\beta'). \end{aligned}$$

It remains to note that $\lim_{n \rightarrow \infty} \frac{1}{n} a_n(\theta \alpha, \theta \beta) = \theta \lim_{n \rightarrow \infty} \frac{1}{n} a_n(\alpha, \beta)$ for any (α, β) in $\mathbb{Q}_{>0}^2$ and any $\theta \in [0, 1] \cap \mathbb{Q}$. ■

One interesting consequence of log-concavity is the following.

Corollary 2.6. *The function $\xi_{\alpha, \beta}$ is either finite for all $(\alpha, \beta) \in \mathbb{Q}_{>0}^2$ or infinite for all $(\alpha, \beta) \in \mathbb{Q}_{>0}^2$.*

Proof. Since $\xi_{\alpha, \beta}$ depends only on α/β , there is no loss of generality in assuming $\beta = 1 - \alpha$ and $\alpha \in (0, 1) \cap \mathbb{Q}$. Suppose $\xi_{\alpha, 1-\alpha} = \infty$ for some $\alpha \in (0, 1) \cap \mathbb{Q}$, and let us show that $\xi_{\beta, 1-\beta} = \infty$ for every other $\beta \in (0, 1) \cap \mathbb{Q}$. For this, let $\gamma = (1+\epsilon)\beta - \epsilon\alpha$ for a sufficiently small $\epsilon \in \mathbb{Q}_{>0}$, so that $\gamma \in (0, 1) \cap \mathbb{Q}$. Then $\beta = \frac{1}{1+\epsilon}\gamma + \frac{\epsilon}{1+\epsilon}\alpha$. By log-concavity,

$$\xi_{\beta, 1-\beta} \geq \xi_{\gamma, 1-\gamma}^{\frac{1}{1+\epsilon}} \xi_{\alpha, 1-\alpha}^{\frac{\epsilon}{1+\epsilon}} = \infty. \quad \blacksquare$$

Remark 2.7. Although we have defined ξ only for rational values in order to avoid technicalities, log-concavity and Proposition 2.1 easily imply that ξ admits a unique continuous extension to $\alpha, \beta \in \mathbb{R}_{>0}$ and that this extension satisfies

$$\xi_{\alpha, \beta} = \lim_{n \rightarrow \infty} (\Xi_{\lfloor \alpha n \rfloor, \lfloor \beta n \rfloor})^{1/(\alpha n + \beta n)} = \lim_{n \rightarrow \infty} (\Xi_{\lceil \alpha n \rceil, \lceil \beta n \rceil})^{1/(\alpha n + \beta n)}.$$

3. Positively decorated simplices and Viro polynomial systems. We start by considering systems of d equations in d variables whose support $\mathcal{A} = \{w_1, \dots, w_{d+1}\} \subset \mathbb{Z}^d$ is the set of vertices of a d -simplex. This case is a basic building block in our construction.

Definition 3.1. A $d \times (d+1)$ matrix M with real entries is called *positively spanning* if all the values $(-1)^i \text{minor}(M, i)$ are nonzero and have the same sign, where $\text{minor}(M, i)$ is the determinant of the square matrix obtained by removing the i th column.

The terminology “positively spanning” comes from the fact that if $\mathcal{A} = \{w_1, \dots, w_{d+1}\}$ is the set of columns of M , then saying that M is positively spanning is equivalent to saying that any vector in \mathbb{R}^d is a linear combination with positive coefficients of w_1, \dots, w_{d+1} .

Proposition 3.2. Let M be a full rank $d \times (d+1)$ matrix with real coefficients. The following statements are equivalent:

1. the matrix M is positively spanning;
2. for any $L \in \text{GL}_d(\mathbb{R})$, $L \cdot M$ is a positively spanning matrix;
3. for any permutation matrix $P \in \mathfrak{S}_{d+1}$, $M \cdot P$ is a positively spanning matrix;
4. all the coordinates of any nonzero vector in the kernel of the matrix are nonzero and have the same sign;
5. the origin belongs to the interior of the convex hull of the column vectors of M ;
6. every vector in \mathbb{R}^d is a nonnegative linear combination of the columns of M ;
7. there is no $w \in \mathbb{R}^d \setminus \{0\}$ such that $w \cdot M \geq 0$.

Proof. The equivalence (1) \Leftrightarrow (4) follows from Cramer’s rule, while (2) \Rightarrow (1) and (3) \Rightarrow (1) are proved directly by instantiating L and P to the identity matrix. The implication (1) \Rightarrow (2) follows from

$$\text{sign}((-1)^i \text{minor}(L \cdot M, i)) = \text{sign}(\det(L)) \cdot \text{sign}((-1)^i \text{minor}(M, i)),$$

while (3) \Leftrightarrow (4) is a consequence of the fact that permuting the columns of M is equivalent to permuting the coordinates of the kernel vectors. The equivalence between (4) and (5) follows from the definition of convex hull: the origin is in the interior of the convex hull of the column vectors if and only if it can be written as a positive linear combination of these vectors. The equivalence between (5) and (6) is obvious, and the equivalence between (5) and (7) follows from Farkas’ lemma. ■

Proposition 3.3. Assume that $\mathcal{A} = \{w_1, \dots, w_{d+1}\}$ is the set of vertices of a d -simplex in \mathbb{R}^d , and consider the polynomial system with real coefficients

$$f_i(X) = \sum_{j=1}^{d+1} C_{ij} X^{w_j}, \quad 1 \leq i \leq d.$$

The system $f_1(X) = \cdots = f_d(X) = 0$ has at most one nondegenerate positive solution and it has one nondegenerate positive solution if and only if the $d \times (d+1)$ matrix $C = (C_{ij})$ is positively spanning.

Proof. Multiplying the system by $X^{-w_{d+1}}$ (which does not change the set of positive solutions), we can assume without loss of generality that $w_{d+1} = \mathbf{0}$. Consider the monomial map $(X_1, \dots, X_d) \rightarrow (X^{w_1}, \dots, X^{w_d})$ which bijects the positive orthant to itself. The map is invertible since \mathcal{A} is affinely independent, and the inverse map transforms the system $f_1(X) = \cdots = f_d(X) = 0$ into a linear system with C as its coefficient matrix. Then the statement follows from Proposition 3.2: by part 4 of the proposition, the unique solution of the linear system lies in the positive orthant if and only if C is positively spanning. ■

Consider now a set $\mathcal{A} = \{w_1, \dots, w_n\} \subset \mathbb{Z}^d$, and assume that its convex hull is a full-dimensional polytope Q . Let Γ be a triangulation of Q with vertices in \mathcal{A} . Assume that Γ is a *regular* triangulation, which means that there exists a convex function $\nu : Q \rightarrow \mathbb{R}$ which is affine on each simplex of Γ but not affine on the union of two different facets of Γ (such triangulations are sometimes called coherent or convex in the literature; see [7] for extensive information on regular triangulations). We say that ν , which is sometimes called the *lifting* function, *certifies* the regularity of Γ . Let C be a $d \times n$ matrix with real entries. This matrix defines a map $\phi : \mathcal{A} \rightarrow \mathbb{R}^d$ as in Theorem A by setting $\phi(w_i)$ to the i th column of C . We say that C *positively decorates* a facet $\tau = \text{conv}(w_{i_1}, \dots, w_{i_{d+1}}) \in \Gamma$ if the $d \times (d+1)$ submatrix of C given by the columns numbered by $\{i_1, \dots, i_{d+1}\}$ is positively spanning. The associated *Viro polynomial system* is

$$(3.1) \quad f_{1,t}(X) = \cdots = f_{d,t}(X) = 0,$$

where t is a positive parameter and

$$f_{i,t}(X) = \sum_{j=1}^n C_{ij} t^{\nu(w_j)} X^{w_j} \in \mathbb{R}[X_1, \dots, X_d], \quad i = 1, \dots, d.$$

The following result is a variation of the main theorem in [23]. There the number of *real* roots of the system (3.1) is bounded below by the number of *odd* facets in Γ (facets with odd normalized volume). Proposition 3.3 allows us to change that to a lower bound for *positive* roots in terms of *positively decorated* simplices.

Theorem 3.4. *Let Γ be a regular triangulation of $\mathcal{A} = \{w_1, \dots, w_n\} \subset \mathbb{Z}^d$, and let $C \in \mathbb{R}^{d \times n}$. Then there exists $t_0 \in \mathbb{R}_+$ such that for all $0 < t < t_0$ the number of nondegenerate positive solutions of the system (3.1) is bounded from below by the number of facets in Γ which are positively decorated by C .*

Proof. Let τ_1, \dots, τ_m be the facets of Γ which are positively decorated by C . For all $\ell \in \{1, \dots, m\}$, the function ν is affine on τ_ℓ , and thus there exist $\alpha_\ell = (\alpha_{1\ell}, \dots, \alpha_{d\ell}) \in \mathbb{R}^d$ and $\beta_\ell \in \mathbb{R}$ such that $\nu(x) = \langle \alpha_\ell, x \rangle + \beta_\ell$ for any $x = (x_1, \dots, x_d)$ in the simplex τ_ℓ . Set $Xt^{-\alpha_\ell} = (X_1 t^{-\alpha_{1\ell}}, \dots, X_d t^{-\alpha_{d\ell}})$. Since ν is convex and not affine on the union of two distinct facets of Γ , we get

$$(3.2) \quad \frac{f_{i,t}(Xt^{-\alpha_\ell})}{t^{\beta_\ell}} = f_i^{(\ell)}(X) + r_{i,t}^{(\ell)}(X), \quad i = 1, \dots, d,$$

where $f_i^{(\ell)}(X) = \sum_{w_j \in \tau_\ell} C_{ij} X^{w_j}$ and $r_{i,t}^{(\ell)}(X)$ is a polynomial, each of whose coefficients is equal to a positive power of t multiplied by a coefficient of C . Since τ_ℓ is positively decorated by C , the system $f_1^{(\ell)}(X) = \cdots = f_d^{(\ell)}(X) = 0$ has one nondegenerate positive solution z_ℓ by Proposition 3.3. It follows that the system $f_1^{(\ell)}(X) + r_{1,t}^{(\ell)}(X) = \cdots = f_d^{(\ell)}(X) + r_{d,t}^{(\ell)}(X) = 0$ has a nondegenerate solution close to z_ℓ for $t > 0$ small enough. More precisely, for all $\varepsilon > 0$, there exists $t_{\varepsilon,\ell} > 0$ such that for all $0 < t < t_{\varepsilon,\ell}$ there exists a nondegenerate solution $z_{\ell,t}$ of $f_1^{(\ell)}(X) + r_{1,t}^{(\ell)}(X) = \cdots = f_d^{(\ell)}(X) + r_{d,t}^{(\ell)}(X) = 0$ such that $\|z_{\ell,t} - z_\ell\| < \varepsilon$. Then using (3.2) we get $f_{1,t}(z_{\ell,t}t^{-\alpha_\ell}) = \cdots = f_{d,t}(z_{\ell,t}t^{-\alpha_\ell}) = 0$. Now choose ε small enough so that the balls of radius ε centered at z_1, \dots, z_m are contained in a compact set $K \subset \mathbb{R}_{>0}^d$. Since the vectors α_ℓ are distinct, there exists $\tau > 0$ such that for all $0 < t < \tau$ the sets $K \cdot t^{-\alpha_\ell} = \{(X_1 t^{-\alpha_{1\ell}}, \dots, X_d t^{-\alpha_{d\ell}}) \mid (X_1, \dots, X_d) \in K\}$, $\ell = 1, \dots, m$, are pairwise disjoint. Set $t_0 = \min(\tau, t_{\varepsilon,1}, \dots, t_{\varepsilon,m})$. Then, for $0 < t < t_0$, each of these sets $K \cdot t^{-\alpha_\ell}$ contains a nondegenerate positive solution $z_{\ell,t}t^{-\alpha_\ell}$ of the system (3.1). ■

4. Duality between regular and positively decorable complexes. In this section, we study the two combinatorial properties on Γ that are needed in order to apply Theorem 3.4: being (part of) a regular triangulation and having (many, hopefully all) positively decorated simplices. As we will see, these properties turn out to be dual to one another. Our combinatorial framework is that of pure, abstract simplicial complexes.

Definition 4.1. A pure abstract simplicial complex of dimension d on n vertices (abbreviated (n, d) -complex) is a finite set $\Gamma = \{\tau_1, \dots, \tau_\ell\}$, where for any $i \in \{1, \dots, \ell\}$, τ_i is a subset of cardinality $d+1$ of $[n] := \{1, \dots, n\}$. The elements of Γ are called facets, and their number (the number ℓ in our notation) is the size of Γ . A subset of cardinality 2 of a facet is called an edge of Γ .

Let $\mathcal{A} = \{w_1, \dots, w_n\}$ be a configuration of n points in \mathbb{R}^d (by which we mean an ordered set; that is, we implicitly have a bijection between \mathcal{A} and $[n]$). An (n, d) -complex Γ is said to be *supported on \mathcal{A}* if the simplices with vertices in \mathcal{A} indicated by Γ , together with all their faces, form a geometric simplicial complex; see [17, Def. 2.3.5]. Typical examples of (n, d) -complexes supported on point configurations are the boundary complexes of simplicial $(d+1)$ -polytopes, or triangulations of point sets. The following definition and proposition relate these two notions.

Definition 4.2. An (n, d) -complex Γ is said to be *regular* if it is isomorphic to a (perhaps nonproper) subcomplex of a regular triangulation of some point configuration $\mathcal{A} \subset \mathbb{R}^d$.

Proposition 4.3. For a pure abstract simplicial complex Γ of dimension d , the following properties are equivalent: (1) Γ is regular. (2) Γ is (isomorphic to) a proper subcomplex of the boundary complex of a simplicial $(d+1)$ -polytope P .

Proof. This is a well-known fact, the proof of which appears, e.g., in [10, sect. 2.3]. The main tool to show the backwards statement (which is the harder direction) is as follows: let F be a facet of P that does not belong to Γ , and let o be a point outside P but very close to the relative interior of F . Project Γ towards o into F to obtain (part of) a regular triangulation of a d -dimensional configuration in the hyperplane containing F . (This construction is usually called a Schlegel diagram of P in F). ■

For $\tau \in \Gamma$ a facet and C a coefficient matrix associated to the point configuration \mathcal{A} , we let C_τ denote the $d \times (d+1)$ submatrix of C whose columns correspond to the $d+1$ vertices in τ .

Definition 4.4. An (n, d) -complex Γ is positively decorable if there is a $d \times n$ matrix C that positively decorates every facet of Γ , that is, such that every submatrix C_τ corresponding to a facet $\tau \in \Gamma$ is positively spanning.

In this language, Theorem 3.4 says that if there is a regular and positively decorable (n, d) -complex of size ℓ , then $\Xi_{d, n-d-1} \geq \ell$.

We now introduce a notion of complementarity for pure complexes. This notion is closely related to matroid duality, and, in fact, our result that regularity and positive decorability are exchanged by complementarity is an expression of that duality via its geometric (and oriented) version: Gale duality.

Definition 4.5. Let Γ be an (n, d) -complex with facets $\{\tau_1, \dots, \tau_{|\Gamma|}\}$. We call it a complement complex of Γ and denote $\bar{\Gamma}$ the $(n, n-d-2)$ -complex with facets $\{\bar{\tau}_1, \dots, \bar{\tau}_{|\Gamma|}\}$, where $\bar{\tau}_i := [n] \setminus \tau_i$.

Lemma 4.6. An (n, d) -complex Γ is positively decorable if and only if its complement $\bar{\Gamma}$ is a subcomplex of the boundary complex of an $(n-d-1)$ -polytope.

Proof. Recall that an (ordered) set of points $\mathcal{A} = \{w_1, \dots, w_n\} \in \mathbb{R}^{n-d-1}$ and an (ordered) set of vectors $\{b_1, \dots, b_n\} \subset \mathbb{R}^d$ are Gale transforms of one another if the following $(n-d) \times n$ and $d \times n$ matrices have orthogonally complementary row-spaces (that is, if the kernel of one equals the row-space of the other):

$$\tilde{A} = \begin{pmatrix} 1 & \cdots & 1 \\ w_1 & \cdots & w_n \end{pmatrix}, \quad C = (b_1 \quad \cdots \quad b_n).$$

Every set of points affinely spanning \mathbb{R}^{n-d-1} has a Gale transform (construct C by using as rows a basis for the kernel of \tilde{A}), and every set of vectors with $\sum b_i = 0$ has a Gale transform (extend the vector $(1, 1, \dots, 1)$ to a basis of the kernel of C).

By construction, Gale transforms have the property that a vector $\lambda \in \mathbb{R}^n$ is the vector of coefficients of a linear dependence of $\{b_1, \dots, b_n\}$ (kernel of C) if and only if it is the vector of values of some affine functional on $\{w_1, \dots, w_n\}$ (row-space of \tilde{A}). This implies the following (see, e.g., [9, Thm. 1, p. 88]): a subset $\tau \subset [n]$ of size $d+1$ indexes a positively spanning submatrix of C if and only if the polytope $\text{conv}(w_1, \dots, w_n)$ has a facet containing exactly the points $\{w_i : i \notin \tau\}$ (by a dimensionality argument, these points must then be vertices and the facet be a d -simplex). Indeed, both things are equivalent to the existence of a unique (modulo scalar) nonnegative λ with support equal to τ in the kernel of C and the row-space of \tilde{A} .

Now let Γ be an (n, d) -complex. If $\bar{\Gamma}$ is realized as a subcomplex of the boundary complex of an $(n-d-1)$ -polytope $P \subset \mathbb{R}^{n-d-1}$, there is no loss of generality in assuming P to be the convex hull of the vertices of $\bar{\Gamma}$. Let w_1, \dots, w_n be those vertices (together with w_i arbitrarily chosen in the interior of P if $i \in [n]$ happens to not be used as a vertex in $\bar{\Gamma}$). Any matrix C constructed as above positively decorates Γ .

Conversely, if a matrix C positively decorates Γ , then there is a nonnegative vector u_τ in the kernel of C with support τ for each facet τ of Γ . Thus, $\lambda = \sum_{\tau \in \Gamma} u_\tau$ is also in the kernel

of C and all its entries are strictly positive. Rescaling the columns of C by the entries of λ , we get a new matrix that has $(1, \dots, 1)$ in the kernel and still positively decorates Γ . The Gale transform described above can then be applied and results in a set of points whose convex hull P contains $\bar{\Gamma}$ as a subcomplex. ■

Corollary 4.7. *Let Γ be an (n, d) -complex. Then the following hold:*

1. *If $\bar{\Gamma}$ is regular, then Γ is positively decorable.*
2. *If Γ is positively decorable, then $\bar{\Gamma}$ is either regular or the boundary complex of a simplicial polytope. If the latter happens, then $\bar{\Gamma}$ minus a facet is regular.*

The following examples illustrate the need to perhaps remove a facet in part 2 of the corollary. A regular and positively decorable complex may not have a regular complement.

Example 4.8. Let Γ be the $(2^d, d)$ -complex formed by the boundary of a cross-polytope of dimension $d+1$. Observe that $\Gamma = \bar{\Gamma}$, which, by Lemma 4.6, implies Γ is positively decorable. Yet, $\bar{\Gamma}$ is not regular since it is not a *proper* subcomplex of the boundary of a $(d+1)$ -polytope.

Example 4.9. The complex $\Gamma = \{\{1, 2, 3, 4\}, \{2, 3, 4, 5\}, \{3, 4, 5, 6\}, \{1, 4, 5, 6\}, \{1, 2, 5, 6\}, \{1, 2, 3, 6\}\}$ is a proper subcomplex of the boundary of a cyclic 4-polytope. Its complement is a cycle of length six, so Γ is regular and positively decorable, but $\bar{\Gamma}$ is not regular.

The following is the main consequence of Lemma 4.6.

Theorem 4.10. *Let Γ be an (n, d) -complex.*

1. *If Γ is regular and positively decorable, then $\Xi_{d, n-d-1} \geq |\Gamma|$ and $\Xi_{n-d-2, d+1} \geq |\Gamma| - 1$.*
2. *If both Γ and $\bar{\Gamma}$ are regular, then $\Xi_{d, n-d-1} \geq |\Gamma|$ and $\Xi_{n-d-2, d+1} \geq |\Gamma|$.*

Proof. The fact that if Γ is regular and positively decorable then $\Xi_{d, n-d-1} \geq |\Gamma|$ is merely a rephrasing of Theorem 3.4. The rest follows from Lemma 4.6. ■

Example 4.11. The inequality $\Xi_{1, k} \geq k + 1$ from Proposition 2.1 is a special case of Theorem 4.10 since a path with $k + 1$ edges is regular and positively decorable (the decorating matrix alternates 1's and -1 's).

5. Relation to bipartite and balanced complexes. In this section, we relate regularity and positive decorability to the following two familiar notions for pure simplicial complexes.

Definition 5.1. *The adjacency graph of a pure simplicial complex Γ of dimension d is the graph whose vertices are the facets of Γ , with two facets adjacent if they share d vertices. We say Γ is bipartite if its adjacency graph is bipartite.*

Definition 5.2 (see [22, sect. III.4]). *A $(d+1)$ -coloring of an (n, d) -complex Γ is a map $\gamma : [n] \rightarrow [d+1]$ such that $\gamma(w_1) \neq \gamma(w_2)$ for every edge $\{w_1, w_2\}$ of Γ . If such a coloring exists, Γ is called balanced.*

Observe that two complement complexes Γ and $\bar{\Gamma}$ have the same adjacency graph. Thus, if one is bipartite, then so is the other. The same is not true for balancedness: a cycle of length six is balanced, but its complement (the complex Γ of Example 4.9) is not. For instance, the simplices $\{1, 2, 3, 4\}$ and $\{2, 3, 4, 5\}$ are adjacent, which implies that 1 and 5 should get the same color. But this does not work since $\{1, 5\}$ is an edge.

Colorings are sometimes called *foldings* since they can be extended to a map from Γ to the d -dimensional standard simplex which is linear and bijective on each facet of Γ . Similarly,

balanced triangulations are sometimes called *foldable* triangulations; see, e.g., [13].

It is easy to show that *orientable* balanced complexes are bipartite. (For nonorientable ones, the same is not true, as shown by the $(9,2)$ -complex $\{123, 234, 345, 456, 567, 678, 789, 189, 129\}$). We here show that being positively decorable is an intermediate property.

Recall that an *orientation* of an abstract d -simplex $\tau = \{w_1, \dots, w_{d+1}\}$ is a choice of calling “positive” one of the two classes, modulo even permutations, of orderings of its vertices and “negative” the other class. For example, every embedding $\varphi : \tau \rightarrow \mathbb{R}^d$ of τ into $d+1$ points not lying in an affine hyperplane induces a canonical orientation of τ by calling an ordering $w_{\sigma_1}, \dots, w_{\sigma_{d+1}}$ positive or negative according to the sign of the determinant:

$$\begin{vmatrix} \varphi(w_{\sigma_1}) & \dots & \varphi(w_{\sigma_{d+1}}) \\ 1 & \dots & 1 \end{vmatrix}.$$

If τ and τ' are two d -simplices with d common vertices, then respective orientations of them are called *consistent* (along their common $(d-1)$ -face) if replacing in a positive ordering of τ the vertex of $\tau \setminus \tau'$ by the vertex of $\tau' \setminus \tau$ results in a negative ordering of τ' . A pure simplicial complex is called *orientable* if one can orient all facets in a manner that makes orientations of all neighboring pairs of them consistent. In particular, every geometric simplicial complex is orientable since its embedding in \mathbb{R}^d induces consistent orientations.

Observe that if we decorate a (geometric or abstract) d -complex Γ on n vertices with a $d \times n$ matrix C as we have been doing in the previous sections, then each facet inherits a canonical orientation from C . When C positively decorates Γ , these orientations are “as inconsistent as can be.”

Proposition 5.3. *Let (Γ, C) be a positively decorated pure simplicial complex. Then the canonical orientations given by C to the facets of Γ are inconsistent along every common face of two neighboring facets. In particular, if Γ is orientable (e.g., if Γ can be geometrically embedded in $\mathbb{R}^{\dim(\Gamma)}$) and positively decorable, then its adjacency graph is bipartite.*

Proof. We need to check that the submatrices of C corresponding to two adjacent facets τ and τ' , extended with a row of ones, have determinants of the same sign. Without loss of generality, assume the matrices (without the row of ones) to be

$$M_\tau = (c_1 \ \dots \ c_d \ c_{d+1}) \quad \text{and} \quad M_{\tau'} = (c_1 \ \dots \ c_d \ c'_{d+1}).$$

Since C positively decorates τ and τ' , and since $\text{minor}(M_\tau, d+1) = \text{minor}(M_{\tau'}, d+1) = |c_1 \ \dots \ c_d|$, we get that all the signed minors $(-1)^i \text{minor}(M_\tau, i)$ and $(-1)^i \text{minor}(M_{\tau'}, i)$ have one and the same sign. In particular, the determinants

$$\begin{vmatrix} c_1 & \dots & c_d & c_{d+1} \\ 1 & \dots & 1 & 1 \end{vmatrix} \quad \text{and} \quad \begin{vmatrix} c_1 & \dots & c_d & c'_{d+1} \\ 1 & \dots & 1 & 1 \end{vmatrix}$$

have the same sign, so the orientations given to τ and τ' by C are inconsistent.

The last assertion is obvious: the positive decoration gives us orientations for the facets that alternate along the adjacency graph, while orientability gives us one that is preserved along the adjacency graph. This can only happen if every cycle in the graph has even length, that is, if the graph is bipartite. ■




Figure 1. A two-dimensional simplicial complex whose adjacency graph is bipartite (left) and which is positively decorable (right) but not balanced. The white triangle 134 is not part of the complex.

Proposition 5.4. Let e_i be the i th canonical basis vector of \mathbb{R}^d , and let $e_{d+1} = (-1, \dots, -1)$. Let Γ be a balanced (n, d) -complex with $(d+1)$ -coloring $\gamma : [n] \rightarrow [d+1]$. Then the matrix C with column vectors $e_{\gamma(1)}, \dots, e_{\gamma(n)}$ in this order positively decorates Γ .

Proof. By construction, every $d \times (d+1)$ submatrix of C corresponding to a facet of Γ is a column permutation of the $d \times (d+1)$ matrix with column vectors e_1, \dots, e_{d+1} in this order. This latter matrix is positively spanning, so the statement follows from Proposition 3.2. ■

Propositions 5.3 and 5.4 imply the following.

Theorem 5.5. For orientable pure complexes (in particular, for geometric d -complexes in \mathbb{R}^d), one has

$$\text{balanced} \implies \text{positively decorable} \implies \text{bipartite}.$$

None of the reverse implications is true, as the following two examples respectively show.

Example 5.6. The $(7, 2)$ -complex of Figure 1 has a bipartite adjacency graph but is not balanced. The right-hand side of the figure describes a positive decoration of the simplex. Therefore, positively decorable simplicial complexes are not necessarily balanced.

Example 5.7. Let Γ be a graph consisting of two disjoint cycles of length four, and let $\bar{\Gamma}$ be its complement, which is an $(8, 5)$ -complex. The adjacency graph of Γ , and hence that of $\bar{\Gamma}$, is bipartite, again consisting of two cycles of length four. On the other hand, since Γ is positively decorable but not part of the boundary of a convex polygon, Lemma 4.6 tells us that $\bar{\Gamma}$ is regular but not positively decorable (remark that $\bar{\Gamma}$ cannot be the whole boundary of a simplicial 6-polytope since for that its adjacency graph would need to have degree six at every vertex).

However, the relationship between balancedness and bipartiteness can be made an equivalence under certain additional hypotheses. A pure simplicial complex Γ is called *locally strongly connected* if the adjacency graph of the star of any face is connected. Locally strongly connected complexes are sometimes called *normal*, and they include, for example, all triangulated manifolds, with or without boundary. See, e.g., the paragraph after Theorem A in [16] for more information on them. By results of Joswig [12, Prop. 6] and [12, Cor. 11], a locally strongly connected and simply connected complex Γ on a finite set \mathcal{A} is balanced if and only if its adjacency graph is bipartite; see also [11, Thm. 5]. In particular, we have the following.




Figure 2. The balanced simplicial complex from Example 5.10.

Corollary 5.8. For simply connected triangulated manifolds (in particular, for triangulations of point configurations), one has

$$\text{balanced} \iff \text{positively decorable} \iff \text{bipartite}.$$

We close this section by illustrating two concrete applications of Theorem 5.5.

Corollary 5.9. Assume that a finite full-dimensional point configuration \mathcal{A} in \mathbb{Z}^d admits a regular triangulation, and let Γ be a balanced simplicial subcomplex of this triangulation. Let $\nu : \mathcal{A} \rightarrow \mathbb{R}$ be a function certifying the regularity of the triangulation, and let $\gamma : \text{Vertices}(\Gamma) \rightarrow [d+1]$ be a $(d+1)$ -coloring of Γ . Then, for $t > 0$ sufficiently small, the number of positive solutions of the Viro polynomial system

$$(5.1) \quad \sum_{w \in \text{Vert}(\Gamma)} t^{\nu(w)} e_{\gamma(w)} X^w = 0$$

is not smaller than the number of facets of Γ .

Example 5.10. Let $d = 2$, $\mathcal{A} = \{w_1, \dots, w_7\}$, where $w_1 = (1, -1)$, $w_2 = (-4, -6)$, $w_3 = (-4, 4)$, $w_4 = (6, 0)$, $w_5 = (3, 6)$, $w_6 = (10, 5)$, and $w_7 = (6, -6)$. Choosing heights $\nu(w_1) = \nu(w_2) = \nu(w_3) = 0$, $\nu(w_4) = 3$, $\nu(w_5) = 5$, $\nu(w_6) = 10$, and $\nu(w_7) = 2$ provides a regular triangulation of \mathcal{A} which has the balanced simplicial subcomplex described in Figure 2. By Corollary 5.9, the Viro polynomial system

$$\begin{aligned} X_1 X_2^{-1} - X_1^{-4} X_2^4 + t^5 X_1^3 X_2^6 - t^{10} X_1^{10} X_2^5 - t^2 X_1^6 X_2^{-6} &= 0, \\ X_1^{-4} X_2^{-6} - X_1^{-4} X_2^4 + t^3 X_1^6 - t^{10} X_1^{10} X_2^5 - t^2 X_1^6 X_2^{-6} &= 0 \end{aligned}$$

has at least six solutions in the positive orthant for $t > 0$ sufficiently small.

In particular, we recover the following result, contained implicitly in [20, Lem. 3.9] concerning maximally positive systems.

We use the notation $\text{Vol}(\cdot)$ for the *normalized volume*, that is, $d!$ times the Euclidean volume in \mathbb{R}^d . A triangulation Γ of \mathcal{A} is called *unimodular* if for any facet $\tau \in \Gamma$ we have

$\text{Vol}(\Gamma) = 1$. A polynomial system with support \mathcal{A} is called *maximally positive* if it has $\text{Vol}(Q)$ nondegenerate positive solutions, where Q is the convex hull of \mathcal{A} . By the Kushnirenko theorem [15], if a system is maximally positive, then all its solutions in the complex torus $(\mathbb{C} \setminus \{0\})^d$ lie in the positive orthant $(0, \infty)^d$.

Corollary 5.11 (see [20]). *Assume that Γ is a regular unimodular triangulation of a finite set $\mathcal{A} \subset \mathbb{Z}^d$. Assume, furthermore, that Γ is balanced or, equivalently, that its adjacency graph is bipartite. Let $\nu : \mathcal{A} \rightarrow \mathbb{R}$ be a function certifying the regularity of Γ , and let $\gamma : \text{Vertices}(\Gamma) \rightarrow [d+1]$ be a $(d+1)$ -coloring of Γ . Then, for $t > 0$ sufficiently small, the Viro polynomial system (5.1) is maximally positive.*

Proof. By Corollary 5.9, system (5.1) has at least $\text{Vol}(Q)$ nondegenerate solutions in the positive orthant for $t > 0$ small enough. On the other hand, it has at most $\text{Vol}(Q)$ nondegenerate solutions with nonzero complex coordinates by the Kushnirenko theorem [15]. ■

This result is also a variant of [23, Cor. 2.4], which, with the same hypotheses except that of Γ being balanced, concludes that system (3.1) is “maximally real”: it has $\text{Vol}(Q)$ nondegenerate solutions in $(\mathbb{R} \setminus \{0\})^d$ (and no other solution in $(\mathbb{C} \setminus \{0\})^d$ by the Kushnirenko theorem).

6. A lower bound based on cyclic polytopes. This section is devoted to the construction and analysis of a family of regular and positively decorable complexes obtained as subcomplexes of cyclic polytopes.

Definition 6.1. *Let d and $n > d+1$ be two positive integers and $a_1 < a_2 < \dots < a_n$ be real numbers. The cyclic polytope $C(n, d+1)$ associated to (a_1, \dots, a_n) is the convex hull in \mathbb{R}^{d+1} of the points $(a_i, a_i^2, \dots, a_i^{d+1})$, $i = 1, \dots, n$.*

The cyclic polytope $C(n, d+1)$ is a simplicial $(d+1)$ -polytope whose combinatorial structure does not depend on the choice of the real numbers a_1, \dots, a_n . In particular, let us denote by $\mathbf{C}_{n,d}$ the d -dimensional abstract simplicial complex on the vertex set $[n]$ that forms the boundary of $C(n, d+1)$. One of the reasons why cyclic polytopes are important is that they maximize the number of faces of every dimension among polytopes with a given dimension and number of vertices. We are especially interested in the case of d odd, in which case the complex is as follows.

Proposition 6.2 (see [7]). *If d is odd, the facets in the boundary of the cyclic polytope $C(n, d+1)$ are of the form*

$$\{i_1, i_1 + 1, i_2, i_2 + 1, \dots, i_{\frac{d+1}{2}}, i_{\frac{d+1}{2}} + 1\}$$

with $1 \leq i_1, i_{\frac{d+1}{2}} \leq n$, and $i_{j+1} > i_j + 1$ for all j . (If $i_{\frac{d+1}{2}} = n$, then $i_1 > 1$ is required, and vertex 1 plays the role of $i_{\frac{d+1}{2}} + 1$). The number of them equals

$$\binom{n - (d+1)/2 - 1}{(d+1)/2 - 1} + \binom{n - (d+1)/2}{(d+1)/2}.$$

Unfortunately, not every proper subcomplex of $\mathbf{C}_{n,d}$ can be positively decorated (except in trivial cases) since its adjacency graph is not bipartite.

Example 6.3. The tetrahedra $\{1, 2, 3, 4\}$, $\{1, 2, 4, 5\}$, and $\{2, 3, 4, 5\}$ form a 3-cycle in the adjacency graph of $\mathbf{C}_{6,3}$.

We now introduce the bipartite subcomplexes of $\mathbf{C}_{n,d}$ in which we are interested. For the time being, we assume both $d + 1 = 2k$ and $n = 2m$ to be even. If we represent any facet $\{i_1, i_1 + 1, i_2, i_2 + 1, \dots, i_k, i_k + 1\}$ of $\mathbf{C}_{2m,2k-1}$ by the sequence $\{i_1, \dots, i_k\}$ (where the vertex 1 plays the role of $i_k + 1$ if $i_k = n$, as happened in Proposition 6.2), we have a bijection between facets of $\mathbf{C}_{2m,2k-1}$ and stable sets of size k in a cycle of length $2m$ (recall that a stable set in a graph is a set of vertices, no two of which are adjacent). Consider the $(2k - 1)$ -dimensional subcomplex $\mathbf{S}_{2m,2k-1}$ of $\mathbf{C}_{2m,2k-1}$ whose facets are the $(2k - 1)$ -simplices $\{i_1, \dots, i_k\}$ such that for all $j \in [k - 1]$ either i_j is odd or $i_{j+1} - i_j > 2$, and such that either $i_1 \neq 2$ or $i_k \neq n$. That is, we are allowed to take two consecutive pairs to build a simplex if both their i_j 's are odd but not if they are even. The adjacency graph of the subcomplex $\mathbf{S}_{2m,2k-1}$ is bipartite since the parity of $i_1 + \dots + i_k$ alternates between adjacent simplices.

Example 6.4. For $n = 6$ and $d + 1 = 4$, we have

$$\mathbf{S}_{6,3} = \{\{1, 2, 3, 4\}, \{1, 2, 4, 5\}, \{1, 2, 5, 6\}, \{2, 3, 5, 6\}, \{3, 4, 5, 6\}, \{1, 3, 4, 6\}\}.$$

The tetrahedra are written so as to show that the adjacency graph is a cycle: each is adjacent with the previous and next ones in the list.

In order to find out and analyze the number of facets in the simplicial complexes $\mathbf{S}_{2m,2k-1}$, we introduce the following graphs.

Definition 6.5. The comb graph on $2m$ vertices is the graph consisting of a path with m vertices together with an edge attached to each vertex in the path. The corona graph with $2m$ vertices is the graph consisting of a cycle of length m together with an edge attached to each vertex in the cycle. Figure 3 shows the case $m = 6$ of both.




Figure 3. The comb graph (left) and the corona graph (right) on 12 vertices.

We denote by $D_{h,k}$ (respectively, $F_{h,k}$) the number of matchings of size k in the comb graph (respectively, the corona graph) with $2(h + k)$ vertices. They form sequences A008288 and A102413 in the Online Encyclopedia of Integer Sequences [18]. The following table shows

the first terms.

	$D_{h,k}$							$F_{h,k}$						
$k =$	0	1	2	3	4	5	6	0	1	2	3	4	5	6
$h = 0 :$	1	1	1	1	1	1	1	1	1	1	1	1	1	1
$h = 1 :$	1	3	5	7	9	11	13	1	4	6	8	10	12	14
$h = 2 :$	1	5	13	25	41	61	85	1	6	16	30	48	70	96
$h = 3 :$	1	7	25	63	129	231	377	1	8	30	76	154	272	438
$h = 4 :$	1	9	41	129	321	681	1289	1	10	48	154	384	810	1520
$h = 5 :$	1	11	61	231	681	1683	3653	1	12	70	272	810	2004	4334
$h = 6 :$	1	13	85	377	1289	3653	8989	1	14	96	438	1520	4334	10672

The numbers $D_{h,k}$ are the well-known *Delannoy numbers*, which have been thoroughly studied [1]. Besides matchings in the comb graph, $D_{h,k}$ equals the number of paths from $(0, 0)$ to (h, k) with steps $(1, 0)$, $(0, 1)$, and $(1, 1)$. The equivalence of the two definitions follows from the fact that both satisfy the following recurrence, which can also be taken as a definition of $D_{h,k}$:

$$D_{h,0} = D_{0,k} = 1 \quad \text{and} \quad D_{h,k} = D_{h,k-1} + D_{h-1,k} + D_{h-1,k-1} \quad \forall i, j \geq 1.$$

The Delannoy numbers can also be defined by either of the formulas in (1.2).

Proposition 6.6.

$$|\mathbf{S}_{2(h+k),2k-1}| = F_{h,k} = D_{h,k} + D_{h-1,k-1}.$$

In particular, $D_{h,k} < |\mathbf{S}_{2(h+k),2k-1}| < 2D_{h,k}$.

Proof. To show that $F_{h,k} = D_{h,k} + D_{h-1,k-1}$, observe that the corona graph is obtained from the comb graph by adding an edge between the first and last vertices of the path. We call that edge the reference edge of the corona graph (the edge $1 - 11$ in Figure 3). Matchings in the corona graph that do not use the reference edge are the same as matchings in the comb graph and are counted by $D_{h,k}$. Matchings of size i using the reference edge are the same as matchings of size $i - 1$ in the comb graph obtained from the corona by deleting the two end-points of the reference edge; this graph happens to be a comb graph with $2(h + k - 2)$ edges, so these matchings are counted by $D_{h-1,k-1}$.

To show that $|\mathbf{S}_{2(h+k),2k-1}| = F_{h,k}$, let $m = h + k$. Observe that each simplex in $\mathbf{S}_{2m,2k-1}$ consists of k pairs $(i_j, i_j + 1)$, $j = 1, \dots, k$, with the restriction that when i_j is even, the elements $i_j - 1$ and $i_j + 2$ cannot be used. In the corona graph, pairs with i_j odd correspond to the spikes and pairs with i_j even correspond to the cycle edge between two spikes, which “uses up” the four vertices of two spikes. This correspondence is clearly a bijection.

The last part follows from the previous two since

$$D_{h,k} < D_{h,k} + D_{h-1,k-1} < 2D_{h,k}. \quad \blacksquare$$

Example 6.7. Proposition 6.6 says that

$$|\mathbf{S}_{10,5}| = F_{2,3} = D_{2,3} + D_{1,2} = 25 + 5 = 30.$$

The following is the whole list of 30 simplices in $\mathbf{S}_{10,5}$. Each row is a cyclic orbit, obtained from the first element of the row by even numbers of cyclic shifts. The first two rows, the

next three rows, and the last row, respectively, correspond to matchings using 0, 1, or 2 edges from the cycle in the pentagonal corona, respectively:

$$\begin{aligned} \mathbf{S}_{10,5} = & \{ \{1, 2, 3, 4, 5, 6\}, \{3, 4, 5, 6, 7, 8\}, \{5, 6, 7, 8, 9, 10\}, \{1, 2, 7, 8, 9, 10\}, \{1, 2, 3, 4, 9, 10\}, \\ & \{1, 2, 3, 4, 7, 8\}, \{3, 4, 5, 6, 9, 10\}, \{1, 2, 5, 6, 7, 8\}, \{3, 4, 7, 8, 9, 10\}, \{1, 2, 5, 6, 9, 10\}, \\ & \{1, 2, 3, 4, 6, 7\}, \{3, 4, 5, 6, 8, 9\}, \{1, 5, 6, 7, 8, 10\}, \{2, 3, 7, 8, 9, 10\}, \{1, 2, 4, 5, 9, 10\}, \\ & \{1, 2, 3, 4, 8, 9\}, \{1, 3, 4, 5, 6, 10\}, \{2, 3, 5, 6, 7, 8\}, \{4, 5, 7, 8, 9, 10\}, \{1, 2, 6, 7, 9, 10\}, \\ & \{1, 2, 4, 5, 7, 8\}, \{3, 4, 6, 7, 9, 10\}, \{1, 2, 5, 6, 8, 9\}, \{1, 3, 4, 7, 8, 10\}, \{2, 3, 5, 6, 9, 10\}, \\ & \{1, 2, 4, 5, 8, 9\}, \{1, 3, 4, 6, 7, 10\}, \{2, 3, 5, 6, 8, 9\}, \{1, 4, 5, 7, 8, 10\}, \{2, 3, 6, 7, 9, 10\} \}. \end{aligned}$$

The symmetry $F_{h,k} = F_{k,h}$ (apparent in the table and which follows from the symmetry in the Delannoy numbers) implies that $\mathbf{S}_{2m,2k-1}$ and $\mathbf{S}_{2m,2m-2k-1}$ have the same size. In fact, they turn out to be complementary.

Theorem 6.8. *Let $\mathbf{S}'_{2m,2k-1}$ denote the image of $\mathbf{S}_{2m,2k-1}$ under the following relabeling of vertices: $(1, 2, 3, 4, \dots, 2m-1, 2m) \mapsto (2, 1, 4, 3, \dots, 2m, 2m-1)$. (That is, we swap the labels of i and $i+1$ for every odd i .) Then $\mathbf{S}'_{2m,2k-1}$ is the complement of $\mathbf{S}_{2m,2m-2k-1}$. In particular, $\mathbf{S}_{2m,2k-1}$ is positively decorable for all k and regular for $k \geq 2$.*

Proof. Consider the following obvious involutive bijection ρ between matchings of size k and matchings of size $m-k$ in the corona graph: for a given matching M , let $\rho(M)$ have the same edges of the cycle as M and the complementary set of (available) spikes. Remember that once a matching has decided to use i edges of the cycle, there are $m-2i$ spikes available, of which M uses $k-i$ and $\rho(M)$ uses the other $m-k-i$. The relabeling of the vertices makes that, for each odd i , if the facet of $\mathbf{S}_{2m,2k-1}$ corresponding to M uses the pair of vertices $i+1$ and $i+2$, then in the facet corresponding to $\rho(M)$ we are using the complement set from the four-tuple $\{i, i+1, i+2, i+3\}$ (except they have been relabeled to $i+1$ and $i+2$ again).

Since the complex $\mathbf{S}_{2m,2k-1}$ is a subset of the boundary of the cyclic polytope and a proper subset for $k \geq 2$, it is regular and positively decorable. ■

Corollary 6.9. *For every $h, k \in \mathbb{Z}$ with $h > 0$, $k > 1$, one has*

$$\Xi_{2k,2h} \geq \Xi_{2k-1,2h} \geq F_{h,k} \geq D_{h,k}.$$

Proof. The first inequality follows from Remark 2.2. The middle inequality is a direct consequence of Theorems 6.8 and 4.10 since $\mathbf{S}_{2(k+h),2k-1}$ is regular and positively decorable. The last inequality follows from Proposition 6.6. ■

Remark 6.10. The above result is our tightest bound for $\Xi_{d,k}$ when d is odd and k even. For other parities, we can proceed as follows:

- We define $\mathbf{S}_{2m-1,2k-1}$ to be the deletion of vertex $2m$ in $\mathbf{S}_{2m,2k-1}$. That is, we remove all facets that use vertex $2m$.
- We define $\mathbf{S}_{2m-1,2k-2}$ to be the link of vertex $2m$ in $\mathbf{S}_{2m,2k-1}$. That is, we keep facets that use vertex $2m$ but remove vertex $2m$ in them.

Clearly, $|\mathbf{S}_{2m,2k-1}| = |\mathbf{S}_{2m-1,2k-1}| + |\mathbf{S}_{2m-1,2k-2}|$. Also, since deletion in the complement complex is the complement of the link, we still have that $\mathbf{S}_{2m-1,2k-1}$ and $\mathbf{S}_{2m-1,2m-2k-2}$ are complements to one another. Moreover, since $\mathbf{S}_{2m,2k-1}$ has a dihedral symmetry acting

transitively on vertices and since each facet has a fraction of k/m of the vertices, we have that

$$|\mathbf{S}_{2m-1,2k-1}| = \frac{m-k}{m} |\mathbf{S}_{2m,2k-1}| \quad \text{and} \quad |\mathbf{S}_{2m-1,2k-2}| = \frac{k}{m} |\mathbf{S}_{2m,2k-1}|.$$

This, along with Corollary 6.9, implies

$$\Xi_{2i-1,2j-1} \geq \frac{j}{i+j} F_{i,j} \quad \text{and} \quad \Xi_{2i,2j} \geq \frac{i+1}{i+j+1} F_{i+1,j}.$$

For $\Xi_{2i,2j-1}$, we can say that

$$\Xi_{2i,2j-1} \geq \Xi_{1,1} \cdot \Xi_{2i-1,2(j-1)} \geq 2 F_{i,j-1}.$$

For example, we have that

$$(6.1) \quad \Xi_{d,d} \geq |\mathbf{S}_{2d+1,d}| = \begin{cases} \frac{1}{2} |\mathbf{S}_{2d+2,d}| & \text{if } d \text{ is odd,} \\ \frac{d/2+1}{d+1} |\mathbf{S}_{2d+2,d+1}| & \text{if } d \text{ is even.} \end{cases}$$

The following table shows the lower bounds for $\Xi_{d,d}$ obtained from this formula, which form sequence A110110 in the Online Encyclopedia of Integer Sequences [18].

d	$\frac{1}{2} \mathbf{S}_{2d+2,d} $	$\frac{d/2+1}{d+1} \mathbf{S}_{2d+2,d+1} $	$ \mathbf{S}_{2d+1,d} $
1	$\frac{1}{2} 4 =$		2
2		$\frac{2}{3} 6 =$	4
3	$\frac{1}{2} 16 =$		8
4		$\frac{3}{5} 30 =$	18
5	$\frac{1}{2} 76 =$		38
6		$\frac{4}{7} 154 =$	88
7	$\frac{1}{2} 384 =$		192
8		$\frac{5}{9} 810 =$	450
9	$\frac{1}{2} 2004 =$		1002

7. Comparison of our bounds with previous ones. In order to derive asymptotic lower bounds on $\Xi_{d,k}$, we now look at the asymptotics of Delannoy numbers.

Proposition 7.1. *For every $i, j \in \mathbb{Z}_{>0}$, we have*

$$\lim_{n \rightarrow \infty} (F_{in,jn})^{1/n} = \lim_{n \rightarrow \infty} (D_{in,jn})^{1/n} = \left(\frac{\sqrt{i^2 + j^2} + j}{i} \right)^i \left(\frac{\sqrt{i^2 + j^2} + i}{j} \right)^j.$$

Proof. The first equality follows from Proposition 6.6. For the second one, since we have

$$D_{i,j} = \sum_{\ell=0}^{\min\{i,j\}} 2^\ell \binom{i}{\ell} \binom{j}{\ell},$$

we conclude that

$$\lim_{n \rightarrow \infty} (D_{in,jn})^{1/n} = \lim_{n \rightarrow \infty} \left(2^\ell \binom{in}{\ell} \binom{jn}{\ell} \right)^{1/n},$$

where $\ell = \ell(n) \in [0, \min\{in, jn\}]$ is the integer that maximizes $f(\ell) := 2^\ell \binom{in}{\ell} \binom{jn}{\ell}$. To find ℓ , we observe that

$$\frac{f(\ell)}{f(\ell-1)} = \frac{2(in-\ell)(jn-\ell)}{\ell^2} = \frac{2(i-\alpha)(j-\alpha)}{\alpha^2},$$

where $\alpha := \ell/n$. Since this quotient is a strictly decreasing function of α and since we can think of $\alpha \in [0, \min\{i, j\}]$ as a continuous parameter (because we are interested in the limit $n \rightarrow \infty$), the maximum we are looking for is attained when this quotient equals 1. This happens when

$$(7.1) \quad \alpha^2 = 2(i-\alpha)(j-\alpha),$$

which implies that

$$(7.2) \quad \alpha = i + j - \sqrt{i^2 + j^2}.$$

(We here take a negative sign for the square root since $\alpha = i + j + \sqrt{i^2 + j^2} > \min\{i, j\}$ is not a valid solution.) We then just need to plug $\ell = \alpha n$ in $2^\ell \binom{in}{\ell} \binom{jn}{\ell}$ and use Stirling's approximation:

$$\begin{aligned} \left(2^{\alpha n} \binom{in}{\alpha n} \binom{jn}{\alpha n} \right)^{1/n} &\sim \left(\frac{2^{\alpha n} (in)^{in} (jn)^{jn}}{(\alpha n)^{\alpha n} ((i-\alpha)n)^{(i-\alpha)n} (\alpha n)^{\alpha n} ((j-\alpha)n)^{(j-\alpha)n}} \right)^{1/n} \\ &= \frac{2^{\alpha} i^i j^j}{\alpha^{2\alpha} (i-\alpha)^{i-\alpha} (j-\alpha)^{j-\alpha}} \\ &\stackrel{(*)}{=} \frac{2^{\alpha} i^i j^j}{(2(i-\alpha)(j-\alpha))^{\alpha} (i-\alpha)^{i-\alpha} (j-\alpha)^{j-\alpha}} \\ &= \left(\frac{i}{i-\alpha} \right)^i \left(\frac{j}{j-\alpha} \right)^j \\ &\stackrel{(**)}{=} \left(\frac{i}{\sqrt{i^2 + j^2} - j} \right)^i \left(\frac{j}{\sqrt{i^2 + j^2} - i} \right)^j \\ &= \left(\frac{\sqrt{i^2 + j^2} + j}{i} \right)^i \left(\frac{\sqrt{i^2 + j^2} + i}{j} \right)^j. \end{aligned}$$

In equalities (*) and (**), we have used (7.1) and (7.2), respectively. ■

Theorem 7.2. For every $d, k \in \mathbb{Z}_{>0}$,

$$\lim_{n \rightarrow \infty} (\Xi_{dn, kn})^{1/n} \geq \left(\frac{\sqrt{d^2 + k^2} + k}{d} \right)^{\frac{d}{2}} \left(\frac{\sqrt{d^2 + k^2} + d}{k} \right)^{\frac{k}{2}}.$$

Proof. By Corollary 6.9 and Proposition 7.1,

$$\lim_{n \rightarrow \infty} (\Xi_{dn, kn})^{1/n} \geq \lim_{n \rightarrow \infty} \left(D_{\frac{kn}{2}, \frac{dn}{2}} \right)^{1/n} \geq \left(\frac{\sqrt{d^2 + k^2} + k}{d} \right)^{\frac{d}{2}} \left(\frac{\sqrt{d^2 + k^2} + d}{k} \right)^{\frac{k}{2}}. \quad \blacksquare$$

Recall from Theorem 2.4 that for every pair (d, k) the limit $\lim_{n \rightarrow \infty} \Xi_{dn, kn}^{1/(dn+kn)}$ exists and depends only on the ratio d/k . Moreover, this limit coincides with the value at $\frac{d}{d+k}$ of the function $\alpha \mapsto \xi_{\alpha, 1-\alpha} = \lim_{n \rightarrow \infty} (\Xi_{\lfloor \alpha n \rfloor, \lfloor (1-\alpha)n \rfloor})^{1/n} \in [1, \infty]$ defined over $(0, 1)$ (see section 2). Theorem 7.2 translates to the following.

Corollary 7.3. *For every $\alpha, \beta > 0$,*

$$\xi_{\alpha, \beta} \geq \left(\frac{\sqrt{\alpha^2 + \beta^2} + \beta}{\alpha} \right)^{\frac{\alpha}{2(\alpha+\beta)}} \left(\frac{\sqrt{\alpha^2 + \beta^2} + \alpha}{\beta} \right)^{\frac{\beta}{2(\alpha+\beta)}}.$$

For example, taking $d = k$, the statement above gives

$$\xi_{1/2, 1/2} = \lim_{d \rightarrow \infty} (\Xi_{d, d})^{1/2d} \geq (\sqrt{2} + 1)^{1/2} \approx 1.5538 \dots$$

This bound is worse than the one coming from $\Xi_{2,2} \geq 7$ (see Proposition 2.1), which implies that $(\Xi_{d, d})^{1/2d} \geq 7^{1/4} \approx 1.6266$. But Corollary 7.3 gives meaningful (and new) bounds for a large choice of d/k or, equivalently, of $\alpha \in (0, 1)$. For example, taking $k = 2d$, the statement above gives

$$\xi_{1/3, 2/3} = \lim_{d \rightarrow \infty} (\Xi_{d, 2d})^{1/3d} \geq \left(\frac{\sqrt{22 + 10\sqrt{5}}}{2} \right)^{1/3} \approx 1.4933 \dots$$

and the same bound is obtained for $\xi_{2/3, 1/3} = \lim_{d \rightarrow \infty} (\Xi_{2d, d})^{1/3d}$.

For better comparison, Figure 4 graphs the lower bound for $\xi_{\alpha, 1-\alpha}$ given by Corollary 7.3. The red dots are the lower bounds obtained from the previously known values $\Xi_{2,2} \geq 7$ and $\Xi_{d,1} = \Xi_{1,d} = d + 1$.

Since the function $\alpha \mapsto \xi_{\alpha, 1-\alpha}$ is log-concave (Proposition 2.5), it is a bit more convenient to plot the logarithm of $\xi_{\alpha, 1-\alpha}$; in such a plot, we can take the upper convex envelope of all known lower bounds for ξ and get a new lower bound. This is done in Figure 5, where the black dashed segments show that the use of Corollary 7.3 produces new lower bounds for $\xi_{\alpha, 1-\alpha}$ whenever $\alpha \in (0.2, 0.8)$.

Remark 7.4. Our results are stated asymptotically, but one can also compute explicit examples where the bound of Corollary 6.9 gives systems with more solutions than was previously achievable. For example, for $d = 115$ and $k = 264$, the maximum number of roots obtainable combining the results in Proposition 2.1 is $2.008 \cdot 10^{62}$, while Corollary 6.9 gives $4.073 \cdot 10^{62}$.




Figure 4. The lower bound for $\xi_{\alpha,1-\alpha}$ coming from Theorem 7.2 (blue curve) versus the ones coming from $\Xi_{2,2} \geq 7$ and $\Xi_{d,1} = \Xi_{1,d} = d + 1$ (red dots).




Figure 5. The different lower bounds for $\log \xi_{\alpha,1-\alpha}$, $\alpha \in (0, 1)$. The red line is the best previously known lower bound using Proposition 2.1 and log-concavity. Our lower bound (blue curve) is above the previously known ones for $\alpha \in [0.2434, 0.3659]$. This range can be extended to $\alpha \in [0.2, 0.8]$ using log-concavity (dashed lines).

8. Limitations of the polyhedral method. We finish the paper with an analysis of how far our methods could possibly be taken. For this, let us denote by $R_{d,k}$ the maximum size (i.e., the maximum number of facets) of a regular $(d-1)$ -complex on $d+k$ vertices such that its complement is also regular. Part 2 of Theorem 4.10 says that

$$\Xi_{d,k} \geq R_{d+1,k},$$

and our main result in section 6 was the use of this inequality to provide new lower bounds for $\Xi_{d,k}$. Observe that either $R_{d,k}$ or $R_{d,k} - 1$ equals the maximum size of a regular positively decorable complex (Corollary 4.7).

Remark 8.1. Our shift on parameters for $R_{d,k}$ is chosen to make it symmetric in k and d : $R_{d,k} = R_{k,d}$.

The inequality $\Xi_{d,k} \geq R_{d+1,k}$ is certainly not an equality, as the following table of small values shows:

		$R_{d+1,k}$						$\Xi_{d,k}$			
$d \backslash k$		1	2	3	4	$d \backslash k$		1	2	3	4
0		1	1	1	1	0					
1		1	3	4	5	1		2	3	4	5
2		1	4	7	8	2		3	≥ 7		
3		1	5	8	≥ 16	3		4			

The values of Ξ come from Proposition 2.1 and those of R come from the following:

- $R_{1,k} = R_{k,1} = 1$ is obvious: a regular 0-dimensional complex can only have one point.
- $R_{2,k} = R_{k,2} = k + 1$ since the largest regular 1-complex with $2 + k$ vertices is a path of $k + 1$ edges, and its complement is regular too (Example 4.11).
- $R_{3,k} \leq 2k + 1$ follows from the fact that a triangulated 2-ball with $k + 3$ vertices has at most $2k + 1$ triangles (with equality if and only if its boundary is a 3-cycle). On the other hand, it is easy to construct a balanced 3-polytope with $k + 3$ vertices for every $k \notin \{1, 2, 4\}$: for odd k , consider the bipyramid over a $(k + 1)$ -gon; for even k , glue an octahedron into a facet of the latter. This shows that $R_{3,k} = 2k + 1$ for all such k (but $R_{3,4} = 8$ instead of 9 since no balanced 3-polytope on 7 vertices exists; the best we can do is a double pyramid over a path of length four).
- $R_{4,4} \geq 16$ follows from the complex $\mathbf{S}_{8,3}$ of size $F_{2,2} = 16$.

It is easy to prove analogues of (2.2) and (2.3) for R . Assume for simplicity that both d and k are even and that $d \leq k$. Then, by Proposition 6.6,

$$R_{d,k} \geq |\mathbf{S}_{d+k,d-1}| = F_{d/2,k/2} \geq D_{d/2,k/2} \geq \binom{\frac{d+k}{2}}{\frac{d}{2}},$$

where the last inequality comes from taking the summand $\ell = 0$ in (1.2). For $k = d$, this recovers (2.4) (modulo a sublinear factor) since $\binom{d}{d/2} \in \Theta(2^d/\sqrt{d})$. More generally, using Stirling's approximation, we get

$$R_{d,k} \geq \binom{\frac{d+k}{2}}{\frac{d}{2}} \underset{d,k \rightarrow \infty}{\sim} \left(\frac{d+k}{k}\right)^{k/2} \left(\frac{d+k}{d}\right)^{d/2} \sqrt{\frac{d+k}{\pi k d}}.$$




Figure 6. The lower bound for $\xi_{\alpha,1-\alpha}$ coming from Theorem 7.2 (blue curve) versus the ones coming from $\Xi_{2,2} \geq 7$ and $\Xi_{d,1} = \Xi_{1,d} = d + 1$ (red dots). The green curve, coming from the upper bound theorem for polytopes, is the limit of the lower bounds that could possibly be produced with our method.

For constant d and big k , we can approximate $\left(\frac{d+k}{k}\right)^{k/2} \simeq e^{d/2}$ so that

$$R_{d,k} \geq \frac{e^{d/2}}{\sqrt{\pi d}} \left(\frac{k}{d} + 1\right)^{d/2}.$$

This, except for the constant factor and for the exponent $d/2$ instead of d , is close to (2.2). Doing the same for constant k and big d gives the analogue of (2.3).

Similarly, one has

$$R_{d+d',k+k'} \geq R_{d,k} R_{d',k'}$$

(the analogue of part 1 in Proposition 2.1) since the join of regular complexes is regular and the complement of a join is the join of the complements.

Regarding upper bounds, since the number of facets of a regular complex cannot exceed that of a cyclic polytope, we have that

$$R_{2d,2k} \leq |\mathbf{C}_{2d+2k,2d-1}| = \binom{2k+d}{d} + \binom{2k+d-1}{d-1},$$

so that, by using Stirling's formula, we get

$$\lim_{n \rightarrow \infty} R_{2dn,2kn}^{1/(2dn+2kn)} \leq \left(\frac{d+2k}{2k}\right)^{\frac{k}{d+k}} \left(\frac{d+2k}{d}\right)^{\frac{d}{2(d+k)}}.$$

Figure 6 shows this upper bound (green line) together with the lower bounds from Figure 4 (blue line and red dots). There are many red dots above the green line, meaning that the

upper bound for R is smaller than the lower bound for Ξ . For example, for the case $d = k$, we have that

$$(R_{d,d})^{1/d} \leq 3\sqrt{3}/2 \approx 2.598 < 2.6458 \approx 7^{1/2} \leq (\Xi_{d,d})^{1/d}.$$

Acknowledgments. We are grateful to Alin Bostan, Alicia Dickenstein, Louis Dumont, Éric Schost, Frank Sottile, and Bernd Sturmfels for helpful discussions.

REFERENCES

- [1] C. BANDERIER AND S. SCHWER, *Why Delannoy numbers?*, J. Statist. Plann. Inference, 135 (2005), pp. 40–54.
- [2] D. BERNSTEIN, *The number of roots of a system of equations*, Funct. Anal. Appl., 9 (1975), pp. 183–185.
- [3] F. BIHAN, *Viro method for the construction of real complete intersections*, Adv. Math., 169 (2002), pp. 177–186.
- [4] F. BIHAN, *Polynomial systems supported on circuits and dessins d'enfants*, J. Lond. Math. Soc. (2), 75 (2007), pp. 116–132.
- [5] F. BIHAN, J. M. ROJAS, AND F. SOTTILE, *On the sharpness of fewnomial bounds and the number of components of fewnomial hypersurfaces*, in Algorithms in Algebraic Geometry, Springer, New York, 2008, pp. 15–20.
- [6] F. BIHAN AND F. SOTTILE, *New fewnomial upper bounds from Gale dual polynomial systems*, Mosc. Math. J., 7 (2007), pp. 387–407.
- [7] J. A. DE LOERA, J. RAMBAU, AND F. SANTOS, *Triangulations: Structures for Algorithms and Applications*, Algorithms Comput. Math. 25, Springer-Verlag, Berlin, 2010.
- [8] B. EL HILANY, *Constructing polynomial systems with many positive solutions using tropical geometry*, Rev. Mat. Complut., 31 (2018), pp. 525–544, <https://doi.org/10.1007/s13163-017-0254-1>.
- [9] B. GRÜNBAUM, *Convex Polytopes*, 2nd ed., Grad. Texts in Math. 221, Springer-Verlag, New York, 2003.
- [10] B. GRÜNBAUM AND G. C. SHEPHARD, *Convex polytopes*, Bull. London Math. Soc., 1 (1969), pp. 257–300.
- [11] I. IZMESTIEV, *Color or Cover*, preprint, <https://arxiv.org/abs/1503.00605>, 2015.
- [12] M. JOSWIG, *Projectivities in simplicial complexes and colorings of simple polytopes*, Math. Z., 240 (2002), pp. 243–259.
- [13] M. JOSWIG AND G. ZIEGLER, *Foldable triangulations of lattice polygons*, Amer. Math. Monthly, 121 (2014), pp. 706–710.
- [14] A. KHOVANSKII, *On a class of systems of transcendental equations*, Soviet Math. Dokl., 22 (1980), pp. 762–765.
- [15] A. G. KUSHNIRENKO, *A Newton polyhedron and the number of solutions of a system of k equations in k unknowns*, Uspekhi Mat. Nauk, 30 (1975), pp. 266–267.
- [16] J.-P. LABBÉ, T. MANNEVILLE, AND F. SANTOS, *Hirsch polytopes with exponentially long combinatorial segments*, Math. Program., 165 (2017), pp. 663–688.
- [17] C. R. F. MAUNDER, *Algebraic Topology*, Dover, New York, 1996.
- [18] OEIS FOUNDATION INC., *The On-Line Encyclopedia of Integer Sequences*, <http://oeis.org>, 2017.
- [19] J.-J. RISLER, *Construction d'hypersurfaces réelles [d'après Viro]*, Séminaire Bourbaki, Vol. 1992/93, Astérisque, 216 (1993), Exp. no. 763, 3, pp. 69–86.
- [20] E. SOPRUNOVA AND F. SOTTILE, *Lower bounds for real solutions to sparse polynomial systems*, Adv. Math., 204 (2005), pp. 116–151.
- [21] F. SOTTILE, *Real Solutions to Equations from Geometry*, Univ. Lecture Ser. 57, American Mathematical Society, Providence, RI, 2011.
- [22] R. STANLEY, *Combinatorics and Commutative Algebra*, 2nd ed., Birkhäuser, Boston, 1996.
- [23] B. STURMFELS, *On the number of real roots of a sparse polynomial system*, in Hamiltonian and Gradient Flows, Algorithms and Control, Fields Inst. Commun. 3, American Mathematical Society, Providence, RI, 1994, pp. 137–143.

- [24] B. STURMFELS, *Viro's theorem for complete intersections*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 21 (1994), pp. 377–386.
- [25] O. VIRO, *Gluing of algebraic hypersurfaces, smoothing of singularities and construction of curves*, in Proceedings of the Leningrad International Topological Conference, Nauka, Leningrad, 1983, pp. 149–197.