

Transforming healthcare with chatbots: Uses and applications—A scoping review

DIGITAL HEALTH
Volume 11: 1–22
© The Author(s) 2025
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20552076251319174
journals.sagepub.com/home/dhj



Marina Barreda^{1,2,3} , David Cantarero-Prieto^{1,2,3}, Daniel Coca¹,
Abraham Delgado⁴ , Paloma Lanza-León^{1,2}, Javier Lera^{1,2} ,
Rocío Montalbán⁴ and Flora Pérez⁴

Abstract

Purpose: The COVID-19 pandemic has intensified the demand and use of healthcare resources, prompting the search for efficient solutions under budgetary constraints. In this context, the increasing use of artificial intelligence and telemedicine has emerged as a key strategy to optimize healthcare delivery and resources. Consequently, chatbots have emerged as innovative tools in various healthcare fields, such as mental health and patient monitoring, offering therapeutic conversations and early interventions. This systematic review aims to explore the current state of chatbots in the healthcare sector, meticulously evaluating their effectiveness, practical applications, and potential benefits.

Methods: This systematic review was conducted following PRISMA guidelines, utilizing three databases, including PubMed, Web of Science, and Scopus, to identify relevant studies on the use and cost of chatbots in health over the past 5 years.

Results: Several articles were identified through the database search ($n = 31$). The chatbot interventions were categorized by similar types. The reviewed articles highlight the diverse applications of chatbot interventions in healthcare, including mental health support, medical information, appointment management, health education, lifestyle changes, and COVID-19 management, demonstrating significant potential across these areas.

Conclusion: Furthermore, there are challenges regarding the implementation of chatbots, compatibility with other systems, and ethical considerations that may arise in different healthcare settings. Addressing these issues will be essential to maximize the benefits of chatbots, mitigate risks, and ensure equitable access to these health innovations.

Keywords

Artificial intelligence, chatbots, healthcare, cost, patient engagement

Submission date: 18 September 2024; Acceptance date: 23 January 2025

Introduction

The COVID-19 pandemic has led to an increased use of healthcare resources. In the context of budget constraints and higher demand for resources, as well as social distancing and remote work, this has resulted in greater use of artificial intelligence and telemedicine, aiming for efficiency and efficacy in the delivery of care and promoting health.^{1–5} Efficiency is understood as better use of available resources and benchmarking, as established by public organizations with their well-known spending reviews, to focus resources on the most priority areas or activities.

¹Research Group of Health Economics, IDIVAL, Santander, Cantabria, Spain

²Department of Economics, University of Cantabria, Santander, Cantabria, Spain

³SANFI (Santander Financial Institute), University of Cantabria, Santander, Cantabria, Spain

⁴Health Department, Cantabria Government, Santander, Cantabria, Spain

Corresponding author:

Javier Lera, Research Group of Health Economics, IDIVAL, Santander, Cantabria, Spain; Department of Economics, University of Cantabria, Santander, Cantabria, Spain.

Email: Javier.lera@unican.es



In this way, despite the fact that many countries have a long tradition of evaluating public spending efficiency and have increasingly focused on this issue since the Great Recession fifteen years ago, it is generally observed that many have predominantly relied on linear spending cuts. This approach has persisted even after the lessons learned during the mentioned period, reflecting a global trend toward reducing expenditures without a thorough review of efficiency and effectiveness in resource allocation.

The use of chatbots in healthcare is also being driven by international collaborations and knowledge sharing. Organizations such as the World Health Organization (WHO) have recognized the potential of artificial intelligence to improve public health and have promoted the development of guidelines and standards for the implementation of these technologies. These initiatives aim to ensure that the benefits of healthcare chatbots are accessible globally, promoting equity and quality in healthcare.⁶

In this context, related to the search for efficiency and the growing advancements in artificial intelligence, the idea of applying these technologies to medical uses has emerged. Consequently, the use of chatbots in the healthcare sector has grown significantly, becoming an innovative tool to improve patient care and optimize health processes. Aggarwal et al.⁷ approximated the following definition for “chatbot” in healthcare: “*AI chatbots are conversational agents that mimic human interaction through written, oral, and visual forms of communication with a user.*” This way of defining chatbots was based on the studies of Laranjo et al.⁸ and Oh et al.⁹ This development can help create the potential to improve health outcomes, potentially alleviating some of the burdens on healthcare providers and systems from being the sole voices of public health outreach (Powell et al., 2023).

Chatbots offer various applications in the healthcare sector, from providing information on symptoms and treatments to scheduling appointments and medication reminders but their main focus till now is within mental health, screening, and public health according to Afsahi et al.¹⁰ For example, some studies^{11–13} showed that chatbots can be effective in the field of mental health, providing clinical support and resources to individuals needing therapeutic conversations. This use can be especially beneficial in contexts where access to mental health professionals is limited. Other uses are related to the use of chatbots for changes in patient’s behaviour towards healthier lifestyles⁷ or even chatbots are helpful in the management of chronicity.^{14,15}

Additionally, chatbots have the potential to monitor patient health and provide early interventions. Irfan et al.¹⁶ in their article show that chatbots can monitor health parameters such as blood pressure and glucose levels, sending alerts to medical professionals when abnormal values are detected. This not only improves the quality of care but can also prevent serious complications by allowing for early intervention.

However, despite these benefits, a significant limitation in the literature is the lack of studies estimating the costs associated with the development and implementation of these systems. Evaluating the cost-effectiveness of chatbots is crucial to ensure that their use is not only clinically beneficial but also economically viable. Studies analysing the cost-benefit relationship will help justify the investment in these technologies and promote their adoption in healthcare systems with limited resources.

Chatbots also systematically collect data through specific queries and data retrieval, improving patient engagement and quality of life. Bergmo¹⁷ emphasizes that comprehending the costs and benefits of eHealth interventions is crucial for several reasons: it helps to demonstrate cost-effectiveness, aids in decision-making, and is vital for creating business models and payment systems that can sustain widespread services. Integrating a cost-effectiveness analysis will allow for measuring not only the costs associated with the implementation and use of these platforms but also the economic benefits derived from improved health and reduced cardiovascular events.

Through this analysis, we aim to provide a comprehensive overview of how these interactive systems are being used to facilitate communication, diagnosis, patient follow-up, and medical information management, thereby contributing to more efficient and accessible care.

Methods

Search strategy and eligibility criteria

We have carried out this literature review following the guidelines of Preferred Reporting Items for Literature Review and Meta-Analyses (PRISMA) (Moher et al. 2009). This review is not registered, and a protocol has not been previously published.

We have used three different databases such as PubMed, Web of Science, and Scopus for the search of relevant articles published during the last 5 years to identify the most current studies published regarding the use and cost of chatbots in health. The search strategy is based on mainly three selected keywords: “Chatbots,” “Health” AND “Cost within the fields” “title, abstract and keywords,” depending on the database consulted (Table 1). Nevertheless, although the word “Costs” was a keyword in our search, this survey did not estimate the cost or savings of the chatbot.

Studies are included if they meet the following inclusion criteria: (a) full-text articles; (b) written in English or Spanish; (c) published in peer-reviewed journals; (d) published in the last 5 years; (e) focused on create a chatbot for the use in medicine. Meanwhile, studies are excluded from the further review if (a) results are not reported; (b) are non-English or non-Spanish publications; (c) are review articles, conference presentations, abstracts,

Table 1. Search strategy: PubMed, Web of Science, and Scopus.

#	Search term
<i>PubMed</i>	
#1.	Chatbot* [Title]
#2.	Health [Title/Abstract]
#3.	Cost [Title/Abstract]
<i>Web of Science</i>	
#1.	Chatbot* [Title]
#2.	Health [Title]
#3.	Cost [Title]
#4.	Limit to: journal article; year of publication >= last 5 years; English or Spanish; Web of Science Core Collection; Open Access.
<i>Scopus</i>	
#1.	Chatbot* [Title]
#2.	Health [Title/Abstract/Keywords]
#3.	Cost [Title/Abstract/Keywords]
#4.	Limit to: journal article; year of publication >= last 5 years; English or Spanish; All Open Access.

Source: Authors' elaboration.

editorial letters, or comments; (d) are deemed outside of the scope of the present review, that is, they do not focus on their application in medicine.

Study selection and data extraction

Using the procedure and criteria mentioned above, 231 articles are identified by the search strategy in our databases. Of all, 58 studies are found in PubMed, 123 in Web of Science, and 50 in Scopus. After the exclusion of the duplicates, 169 articles are screened. Titles and abstracts are reviewed, and 86 articles do not fulfill the inclusion criteria, so they are excluded. In total, 83 studies are included for full-text screening and reviewed, of which 52 are excluded, to do this, figures (such as flowcharts) and tables were used to synthesize the data. Therefore, the final number of studies included in this review is 33. Figure 1 summarized the search selection procedure as Table 2 shows the results.

After removing duplicates, two independent reviewers performed the first selection of the screening process. To assess the quality of the articles selected in this systematic

review, the authors applied a structured methodology to ensure the rigor and reliability of the included studies. On the one hand, studies were initially assessed against predefined inclusion criteria, such as the relevance of the approach to the use of chatbots in healthcare (communication, diagnosis, patient monitoring, and medical information management), their methodological design, and the availability of complete data. Articles that did not meet these criteria were excluded. On the other hand, studies that passed the initial review were assessed by at least two independent reviewers to reduce bias in the quality assessment. In case of disagreement, both researchers discussed the article with a third reviewer and a joint decision was reached (include or exclude the article). For each of the studies selected for inclusion, one author designed a data extraction procedure that was approved by the other review authors. The extraction form included the following data: authors and year of publication, country, study objectives, population characteristics and sample size, type of nudge, duration of intervention, setting, study outcome, and outcome.

As shown in Figure 2, the first paper found on the use and cost of chatbots in healthcare was published in 2020, despite this literature review covering the last 5 years. In this document, a total of 31 articles are considered. The temporal evolution has been increasing from 2020 to 2023. Thus, in the first year, there were 2 articles, while the highest number of articles was published in 2023 (14 articles). The annual growth of articles has been as follows. From 2020 to 2021, articles analysing interactive systems to facilitate communication, diagnosis, patient follow-up, and medical information management have increased by 250.00%, from 2 articles to 7. From 2021 to 2022, this percentage has grown to a lesser extent, by 14.29%, from 7 to 8 publications related to our objective. Meanwhile, from 2022 to 2023, the annual growth corresponds to 75.00%, increasing to 14 articles.

Then, Figure 3 plotted the distribution of the selected studies by country. In this context, most studies were conducted in India ($N=10$), followed by China ($N=2$) and USA ($N=2$). The remaining articles considered only one country, such as those listed below: Bahrein, Brazil, Egypt, Italy, Johannesburg, Kazakhstan, Malaysia, Morocco, Pakistan, Philippines, Singapore, and Sri Lanka. Two other articles look at the European continent in general, while three other articles look at several countries at once.

According to the applications of chatbots in health, here were a synthesis of the chatbot interventions described in the reviewed articles. The articles were grouped by similar types of interventions, demonstrating the diverse applications of chatbot interventions in healthcare, ranging from mental health support and medical information to appointment management and health education.

Several articles included mental health support and psychological well-being chatbots. Kaywan et al.²³ assessed




Figure 1. Flow diagram for the search process for identifying and including references for the systematic review.

the feasibility and effectiveness of detecting early and assisting depression through an artificial intelligence chatbot, which showed some potential for assisting with automation and discreet communication. Potts et al.²⁴ evaluated a multilingual mental health and well-being chatbot called ChatPal, which aimed at improving mental well-being of the participants from rural areas. Meanwhile, Sabour et al.²⁷ analyzed effectiveness in reducing symptoms of mental distress through a conversational agent providing cognitive support (Emohaa), concluding significant improvements in symptoms of depression, negative affect, and insomnia. Suharwardy et al.²⁸ evaluated acceptability and efficacy for mood management using a mental health chatbot intervention in postpartum women, showing results such as high user satisfaction but limited impact on depressive symptoms due to baseline conditions. In addition, Upadhyaya and Kaur²⁹ assisted individuals using a

chatbot that employs Cognitive Behavioural Therapy to provide mental health support, help manage emotions, and cope with thoughts and experiences. Zhu et al.¹¹ investigated determinants of user satisfaction, interaction, and clinical support with mental health chatbots, designed for symptom collection, health status monitoring, medication reminders, and patient education on their health conditions and presenting positive influences on user satisfaction and continuance intention. In the case of Hakani et al.,³³ the authors detected signs of anxiety and suggested methods for controlling depression using a digital chatbot, but their results were limited due to lack of data for accurate predictions. Bendig et al.³⁹ improved psychological well-being using agent-based software and they found positive results in psychological well-being. In particular, Gabrielli et al.⁴¹ managed stress and anxiety through a psychoeducational chatbot (Atena) and showed benefits for university

Table 2. Summary of the case studies.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Anjum et al. (2023) ¹⁸	India	To develop a health chatbot powered by AI and natural language processing, which can diagnose conditions and offer relevant information, improving healthcare accessibility while lowering expenses.	Potential patients.	Using a medical Chatbot which offers appropriate guidance on maintaining a healthy lifestyle.		NLP and machine learning are the most effective approaches for building chatbots.	Positive
Bhuvanesh et al. (2023) ¹⁹	India	To create a chatbot that evaluates symptoms and categorizes the potential diseases the patient might be experiencing.	Patients with health issues.	A method based on knowledge graphs was proposed for developing health chatbots that evaluate patient symptoms and determine the disease based on the assessment.		The medical chatbot model based on knowledge graphs achieves 99% accuracy and can offer enhanced diagnoses as more knowledge bases are integrated.	Positive
Booth et al. (2023) ²⁰	Europe	The chatbot's log data is analyzed to provide insights into usage patterns, identify different user types through clustering, and uncover associations	A total of 579 adult users (over 18 years old).	The onset of the COVID-19 outbreak.	A span of 12 weeks	This research has shed light on the demographics of ChatPal chatbot users, usage trends, and the relationships between different app features. These insights can guide future app development by highlighting which features are most frequently utilized.	Positive
Goldnadel et al. (2023) ²¹	USA	Create a digital chatbot capable of providing anonymous interactions with an unlimited number of users around	No personal data was utilized, ensuring that all interactions remained anonymous.	Interactive chatbot for educating the general public about alcohol.	From November 9, 2021, to January 16, 2022.	Preliminary findings suggest a promising potential for leveraging artificial intelligence solutions to improve	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
		the clock on topics related to alcohol, including strategies for reducing risks. It should be available in several languages, free of charge, and compatible with various devices.				health education on the risks associated with alcohol consumption.	
Hasnain (2023) ²²	Pakistan	Offering information on monkeypox (mpox) infections in Pakistan through ChatGPT.	People in general.	Utilize ChatGPT to deliver information on monkeypox infections in Pakistan.		There are potential drawbacks regarding the latest mpox situation in Pakistan and other countries, including ChatGPT's limitations in predicting trends with small datasets and concerns about its performance and reliability.	Negative
Irfan & Zafar (2023) ¹⁶	India	Develop an AI-driven medical chatbot capable of diagnosing illnesses and offering fundamental information about diseases prior to seeking a doctor's consultation.	Individuals experiencing health concerns.	The suggested system is a text-based conversational agent designed to engage users in dialogue about their health concerns. It queries users about their symptoms to assist in diagnosing their condition.		The chatbot is user-friendly and accessible to anyone who can write in their native language, whether on a mobile app or desktop version. It offers personalized diagnoses based on the symptoms provided by users.	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Kaywan et al. (2023) ²³	India	To assess the practicality and efficacy of employing AI-powered chatbots for the early detection of depression. .	50 participants with an average age of 34.7 years.	Determining the levels of depression.		While the chatbot is not designed to replace mental health professionals, it demonstrates potential as a tool for supporting automation and providing discreet communication through validated scoring systems.	Limited
Potts et al. (2023) ²⁴	Northern Ireland, Scotland, the Republic of Ireland, Sweden, and Finland	To assess the impact of a multilingual mental health and well-being chatbot mental well-being.	348 participants from rural areas, with 73% being female, aged between 18 and 73 years (average age: 30).	A pre-and post-intervention study of the ChatPal chatbot.	A duration of 12 weeks	The study evaluated the Warwick-Edinburgh Short Mental Well-Being Scale (SWEMWBS) and observed enhancements in well-being scores. Additionally, the Satisfaction with Life Scale (SWLS) and the World Health Organization Well-Being Index (WHO-5) were assessed, with both showing improvements throughout the study period.	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Prakasam et al. (2023) ²⁵	India	Machine learning algorithms designed to facilitate the process of scheduling, canceling, and rescheduling doctor's appointments based on users' preferred times.	Patients	Development of a chatbot for managing the booking, cancelation, and rescheduling of doctor's appointments.		The WhatsApp chatbot allows patients to automatically book, cancel, and reschedule doctor appointments without the need for human assistance, thereby streamlining continuous communication between patients and doctors.	Positive
Rekik et al. (2023) ²⁶	Egypt	Develop a medical chatbot tailored for the Tunisian dialect. Based on users' symptoms, the chatbot will make health predictions, diagnose conditions using the identified symptoms, and suggest appropriate medical care. Users will be able to interact with the chatbot	356 patients who received the correct diagnosis.	To develop a medical chatbot tailored for the Arabic language, with a specific focus on the Tunisian dialect.		The latter approach achieved the highest performance, with an F1 score of 98.60%. Although the study utilized only 356 data pairs, the results highlighted the effectiveness of our implemented chatbot.	Positive
Sabour et al. (2023) ²⁷	China	To evaluate the effectiveness of the conversational agent in alleviating symptoms of mental distress.	Patients with mental problems	A conversational agent offering cognitive support, known as Emohaa		Participants who utilized the chatbot saw notably greater improvements in symptoms of mental distress, such as depression, negative affect, and insomnia	Positive
Suhaarwardy et al. (2023) ²⁸	California (USA)	To assess the acceptability and initial effectiveness of a mental health chatbot	192 English-speaking postpartum women, aged 18 or older, who	Mental health chatbot intervention versus standard care alone	Assessment of symptoms of depression and	The sample showed no signs of depression at baseline, which	Limited

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Upadhyaya & Kaur (2023) ²⁹	India	Creating a multilingual health awareness chatbot is proposed to assist individuals. This AI-driven chatbot aims to provide answers to questions about diseases, treatments, and preventive measures.	have had a live birth and have access to a smartphone.	The chatbot named SERMO is employed to address the rise in suicide attempts and deaths due to the shortage of mental health professionals. It uses Cognitive Behavioral Therapy (CBT) to assist individuals with mental health conditions in managing their emotions and coping with their thoughts and experiences.	anxiety during hospitalization (baseline) and at 2, 4, and 6 weeks postpartum	constrained the chatbot's ability to address depressive symptoms effectively. However, 91% of users expressed satisfaction with the chatbot, and 80% felt at ease using a mobile app for managing their mood.	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Vera & Palaoag (2023) ³⁰	Philippines	To offer design guidelines for creating a prototype system that collects current information on medicinal plants and their uses, with the aim of distributing this information to underserved regions with limited medical services.	150 participants from remote regions of Catanduanes, who predominantly depend on herbal remedies provided by local clinics.	A prototype of an intelligent interactive system is presented, integrating chatbot technology with artificial intelligence (AI) to address queries about treatment alternatives and the use of various medicinal plants for common health issues, thereby promoting alternative healing methods.		The MedPlantBot system prototype could significantly impact the healthcare industry by promoting sustainability and ethical practices.	Positive
Bhangdia et al. (2021) ³¹	India	To utilize technology to enable more individuals to access support. The proposed approach involves the development of a responsive therapist robot that generates suitable responses based on the user's emotions.	Individuals with mental health conditions.	Developing a reliable and accurate method for detecting emotions.		Developing a multimodal therapy chatbot enables accurate detection of emotions, provision of appropriate responses, and suggestions of activities based on the user's mood. Incorporating facial recognition technology could further improve the effectiveness of the system.	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Binh et al. (2022) ³²	Europe	Designing solutions to improve the lives of individuals who do not need body-specific benefits but could greatly benefit from an active lifestyle, especially those suffering from obesity, is the primary goal.	45 local communities throughout the European Union.	The creation of the platform (STOP KM-EP) will play a crucial role in applying strategies to enhance the quality of life for individuals living with obesity by promoting an active lifestyle.		STOP KM-EP platform is effective and appropriate for gathering data on fitness profiles, fitness information, knowledge resources, logs, and chatbot functions	Positive
Hakani et al. (2022) ³³	India	To combine these advanced technologies into a single system to deliver highly accurate outcomes. This project covers a broad range of areas, focusing on both identifying and addressing depression in individuals.	People with depression.	To develop a digital platform that identifies indicators of anxiety and provides recommendations for managing depression.		The system offers a comprehensive solution for detecting and managing depression, giving it an edge over current approaches. However, there could be cases where the user lacks sufficient tweets to accurately assess their depression score.	Limited
Mehta & Singh (2022) ³⁴	India	To demonstrate that the proposed medical robot can provide greater benefits compared to most chatbots currently used in the healthcare sector.	People in general.	To establish human-computer interaction through the use of NLP technology.		A supportive system has been designed for offices and medical facilities that offers treatment recommendations, identifies prescriptions, and integrates with a medication API.	Positive
Mellah et al. (2022) ³⁵	Morocco	Analysing and predicting whether a patient has COVID-19 using a chatbot.	50 patients.	The chatbot operates in both Arabic and French, providing guidance and actions to safeguard lives and		The system accurately identified high risk in 25 out of 25 COVID-19 patients and low risk in 20 out of 25 individuals	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Natshéh, & Jabed (2022) ³⁶	Bahrain	Design a 24/7 helpline chatbot that offers users crucial support and information to handle the rising demand for COVID-19 care.	People with COVID-19.	curb the virus's transmission based on its analytical findings. Developing an AI-driven chatbot utilizing free and open-source tools and frameworks		without the virus, resulting in a precision rate of 90%.	Positive
Okonkwo et al. (2022) ³⁷	Johannesbourg	Develop a chatbot designed to assess students and verify their COVID-19 vaccination status.	106 university students.	The chatbot (COVID.Bot) is an advanced interactive system capable of engaging with students to ascertain their COVID-19 vaccination status for entry to the institution and participation in its activities.		The results showed that most participants regarded the chatbot as effective (86.8%), well-suited to their devices and lifestyles (80.2%), capable of delivering accurate responses (81.1%), and user-friendly (80.2%). The average reliability score was .87, reflecting satisfactory reliability.	Positive
Peng et al. (2022) ³⁸	Malaysia	Examine the obstacles and enablers affecting the uptake of an AI chatbot for HIV testing and prevention among men who have sex with men in Malaysia, highlighting perceived benefits, challenges, and preferred functionalities.	Five focus group interviews were carried out, each consisting of 5 to 9 participants.	An AI chatbot developed to support HIV testing and prevention.	July- September 2021	Barriers and facilitators for MSM's acceptance of an AI chatbot for HIV testing are identified. Concerns include the chatbot's accuracy, convenience, and social stigma. Facilitators include connections to healthcare and self-testing services. Recommendations include implementing anonymous settings, MSM-friendly platforms, and clear guidance on HIV-related topics.	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Zhu et al. (2022) ¹¹	China	Examine the factors influencing user satisfaction and their intention to continue using mental health chatbots.	371 users.	Mental health chatbot (Xiaolv).	1 week	An analysis of 371 Chinese users revealed that while personalization, enjoyment, learning, and the COVID-19 pandemic have some positive but weak effects on user satisfaction and the intention to keep using the chatbot, satisfaction itself has only a slight positive influence on continued use. Voice interaction did not significantly predict either satisfaction or continuance. This research underscores the relevance of the Theory of Consumption Values for understanding mental health chatbot usage.	Positive
Bendig et al. (2021) ³⁹	Germany, Austria, and Switzerland	Assess the clinical effectiveness and user acceptance of the brief IMI SISU software based on agents, in comparison to a control group that is on a waiting list.	120 Participants of the German language.	Develop a medical chatbot utilizing neural networks to offer diagnostic information and deliver essential details about various diseases.		SISU is regarded as a viable intervention with the potential to enhance participants' psychological well-being. In the feasibility trial, a dropout rate of 39% was observed during the study's course.	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Dhinakaran et al. (2021) ⁴⁰	Singapore	Assess the practicality and appeal of a conversational agent created to promote healthy lifestyle behaviour changes within the general population of Singapore	60 Participants.	To support the design and development of a conversational agent centered on promoting healthy lifestyle changes.	August-September 2019.	Future digital solutions, such as conversational agents targeting healthy lifestyles and diabetes prevention, should tackle the obstacles revealed in this study and inspire individuals to embrace healthier habits.	Positive
Gabrielli et al. (2021) ⁴¹	Italy	Perform an initial evaluation to measure how well Atena, a psychoeducational chatbot, engages users and supports effective stress and anxiety management among university students.	71 university students.	Psychoeducational chatbot designed to assist with managing stress and anxiety in a healthy way.	During the Covid-19 pandemic.	The benefits of employing a digital intervention, such as a chatbot, for promoting healthy coping strategies among university students experiencing high levels of stress are demonstrated.	Positive
Omarov et al. (2021) ⁴²	Kazakhstan	Explore the design and deployment of the mobile chatbot psychologist, emphasizing the AIML-driven conversational engine and the incorporation of cognitive-behavioural therapy (CBT) techniques.	Individuals facing psychological distress	An AI-powered mobile chatbot psychologist is introduced, utilizing Artificial Intelligence Markup Language (AIML) and Cognitive Behavioral Therapy (CBT) to deliver customized psychological support.		The AI-powered mobile chatbot psychologist marks a major advancement in mental health care by leveraging AI and evidence-based therapeutic methods to offer accessible, affordable, and tailored psychological support.	Positive
Larbi et al. (2021) ⁴³	Switzerland/ Norway	To evaluate the usability of the chatbot called MYA.	Thirty volunteers participated, including 15 males aged 18 to 69 years and 15 females aged 18 to 49 years.	To create MYA, a Telegram-based chatbot aimed at encouraging increased physical activity.		Although the results are promising, it is still uncertain whether the chatbot can effectively motivate individuals to	Limited

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Pivithuru et al. (2021) ⁴⁴	Sri Lanka	Develop a versatile, user-friendly technology for an Integrated Electronic Patient Health Record System to replace traditional paper records with a comprehensive electronic system	Doctors	A smartcard-based Integrated e-Patient Card System		boost their physical activity levels.	Positive
						The proposed system, utilizing CNN and GradCAM with a transfer learning approach, attained an accuracy of 95.21%. It is capable of diagnosing lung diseases with over 90% accuracy. The system enables a radiographer to upload an X-ray image, which is then analyzed to provide the disease name, its probability, and its location.	
Daley et al. (2020) ⁴⁵	Brazil	Conduct a preliminary assessment of Vitalik, a mental health chatbot, to evaluate its engagement and effectiveness in reducing anxiety, depression, and stress.	3629 Portuguese-speaking users from Brazil, all over 18 years old with internet access. Of these, 76% were female and 52% were aged 18–24 years.	To create Vitalik, an automated chatbot that provides mental health information through an innovative conversational format.	1 month period	Scores for anxiety, depression, and stress decreased from the initial assessment to the follow-up. Higher user engagement was associated with more significant improvements in anxiety and depression levels after the intervention.	Positive

(continued)

Table 2. Continued.

Article	Country	Objective	Population characteristics	Chatbot intervention	Intervention duration	Study outcome(s)	Results +/-
Srivastava et al. (2020) ⁴⁶	India	Develop a diagnostic chatbot that interacts with patients to address their medical queries and issues, offering a personalized diagnosis based on their symptoms and profile.	Doctors and patients.	Develop a chatbot, named Medibot, to deliver personalized diagnoses.		For extracting diagnosed symptoms, correct symptoms were identified with a recall rate of 65% and a precision rate of 71%.	Positive

Source: Authors' elaboration.
AImL: artificial intelligence markup language; CBL: cognitive behavioral therapy; NLP: natural language processing technology; HIV: human immunodeficiency virus.




Figure 2. Evolution of published papers over time.
Source: Authors' elaboration.




Figure 3. Distribution of individual country articles by country.
Note: Three articles were multi-country (Potts et al. 2023; Bendig et al. 2021; Larbi et al. 2021) and two analyzed Europe (Booth et al. 2023; Binh et al. 2022), and we could not classify them.
Source: Authors' elaboration.

students in stress management. Omarov et al.⁴² provided personalized psychological support using a mobile psychologist chatbot based on artificial intelligence markup language and cognitive behavioral therapy, concluding significant advances in accessible psychological support. For example, Daley et al.⁴⁵ evaluated the engagement and effectiveness of a mental health chatbot (Vitalik), delivering mental health content in a conversational format, showing decreased anxiety, depression, and stress levels.

On one hand, other articles were focused on medical and health information chatbots. Anjum et al.¹⁸ provided accessible healthcare information using artificial intelligence and natural language processing technology through a medical chatbot offering advice on healthy lifestyles, who demonstrated effective results in reducing costs and making healthcare accessible. In this context, Bhuvanesh et al.¹⁹ developed a chatbot for symptom assessment and disease classification, concluding that it presented high accuracy in disease diagnosis. In particular, Hasnain²² provided information about monkeypox infection using

ChatGPT for disease information dissemination, highlighting potential disadvantages in prediction competency. Rekik et al.²⁶ built a medical chatbot for health predictions and medical care based on symptoms for the Tunisian dialect and it also demonstrated high efficiency and accuracy. Vera and Palaoag³⁰ provide information on medicinal plants and their applications using a chatbot for alternative healing methods, which made a significant contribution to healthcare with sustainable practices.

On the other hand, two articles were related to disease diagnosis and communication management chatbots. Irfan and Zafar¹⁶ diagnosed illnesses and provide basic disease information through a text-to-text conversational agent for health problem diagnosis, which was easy to use and offered personalized diagnoses. Prakasam et al.²⁵ analyzed a WhatsApp chatbot that facilitated managing, booking, canceling, and rescheduling doctor appointments and enhanced communication between patients and physicians. Mehta and Singh³⁴ used natural language processing technology, that is, a robot, for human-computer

communication in medical contexts, showing the useful applications in offices and medical centers. In the case of Bhangdia et al.,³¹ they evaluated symptoms and classify possible diseases based on symptoms and emotions, provided appropriate responses, and suggested activities based on the user's mood using a chatbot. Pivithuru et al.⁴⁴ created a multifunctional, user-friendly technology for an Integrated Electronic Patient Health Record System, including, specifically, a chatbot for medical record management and showing 90% accuracy in lung disease diagnosis. Srivastava et al.⁴⁶ built a diagnosis bot (Medibot) for engaging patients in medical queries and providing individualized diagnoses based on symptoms and patient profiles, presenting a recall of 65% and a precision of 71% in symptom identification.

In addition, health education and awareness were contexts covered by some articles considered. Goldnadel et al.²¹ educated on alcohol-related topics with risk reduction strategies through digital conversational agent accessible 24/7, promising scenario for health education on alcohol harms. Peng et al.³⁸ assisted in Human Immunodeficiency Virus (HIV) testing and prevention among marginalized populations using artificial intelligence chatbot for culturally sensitive health tool, and the results showed valuable insights into key features for design and implementation.

Two articles also analyzed chatbots for promoting healthy lifestyle changes. Binh et al.³² improved the lives of people suffering from obesity through an active lifestyle using a specific platform, showing the positive usability and feasibility, suitable for collecting fitness data and logs. Similarly, Dhinakaran et al.⁴⁰ promoted healthy lifestyle changes using a conversational agent, which made useful recommendations for improving future digital interventions. Larbi et al.⁴³ tried to increase physical activity using a Telegram-based chatbot, but their results were inconclusive based on motivation to increase physical activity.

Related to COVID-19 management chatbots, we found some other articles. Booth et al.²⁰ analyzed the chatbot's log data to provide information on usage patterns and different types of users during the beginning of the COVID-19 pandemic, presenting positive insights into user types and usage patterns, helping in further app development. In the case of Mellah et al.,³⁵ they suggested measures against COVID-19 using a bilingual chatbot (Arabic and French) with a 90% accuracy in risk prediction. Natsheh and Javed³⁶ provided 24/7 help to meet the increasing demand of COVID-19 patients, having the capability to meet future healthcare sector needs. Okonkwo et al.³⁷ assessed the vaccination status of students against COVID-19 using an interactive system, with high levels of effectiveness and usability.

Discussion

This article aimed at summarizing the growing adoption and effectiveness of chatbots in health care systems. We

demonstrate that chatbots are revolutionizing healthcare by providing accessible, efficient, and effective solutions across different domains, including mental health and patient management, disease diagnosis, and health information dissemination. This fact indicates a significant shift towards digital health solutions.

The integration of AI and NLP in chatbots enhances their functionality and accuracy. Studies like those by Bhuvanesh et al.¹⁹ and Rekik et al.²⁶ demonstrate high diagnostic accuracy and an improved access to healthcare, suggesting that AI-driven chatbots could become integral components of future healthcare systems. Moreover, chatbots have proven to be effective in engaging patients and providing health information in a user-friendly manner. The ability of chatbots to operate 24/7 and provide immediate responses makes healthcare more accessible, particularly for populations with limited access to traditional healthcare services as Natsheh and Javed³⁶ demonstrated.

Regarding costs, the fact that routine tasks such as appointment scheduling, management of medical records, and processing of administrative data, which reduces the workload of healthcare staff can be automatized and provide preliminary health assessments, chatbots can significantly reduce direct costs. Anjum et al.¹⁸ and Upadhyaya and Kaur²⁹ highlighted the cost-reducing potential of healthcare chatbots, making systems more sustainable and efficient while chatbots improve the quality in healthcare delivery.

On the one hand, studies such as those by Kaywan et al.²³ and Sabour et al.²⁷ provide clear evidence on the potential of AI-driven interventions in addressing mental health problems through accessible and discreet platforms. The evidence suggests that mental health chatbots can serve as valuable tools in extending mental health support, particularly in underserved areas. The success of multilingual and culturally sensitive chatbots, like ChatPal and those targeting rural populations, indicates that such technologies can bridge gaps in mental health care accessibility and equity. Moreover, the diverse applications of chatbots, from cognitive behavioural therapy (CBT) to psychoeducation, illustrate the flexibility and potential of these technologies in delivering personalized mental health care. This adaptability is crucial for developing tailored interventions that can cater to varied demographic and psychological needs.

The broader application of chatbots in health information dissemination, disease diagnosis, and lifestyle promotion demonstrates their potential to contribute significantly to overall health and well-being. Chatbots can offer personalized healthcare solutions by using patient data to provide tailored advice and interventions. This personalized approach is crucial for managing chronic diseases and improving patient outcomes, as demonstrated by Upadhyaya and Kaur²⁹ and Bhangdia et al.³¹

In the case of recent Large Language Models (LLMs), they have significantly surpassed their predecessors in the

ability to understand and generate natural language, enabling much more fluid and accurate interactions with patients.^{47,48} This improves the quality of communication and user satisfaction. Greater accuracy in language analysis and the generation of contextualized responses reduces errors and improves the reliability of diagnosis as well as treatment recommendations.⁴⁹ In addition, LLMs allow chatbots to better adapt to patients' needs, providing more personalized responses based on deeper analysis of available data. This not only increases effectiveness in monitoring and managing medical information but also improves clinical outcomes^{50,51}. Nevertheless, while previous studies showed that chatbots could reduce costs by alleviating the burden on healthcare staff, the new LLMs offer even greater efficiencies, allowing for better scaling without compromising service quality.^{52,53} However, we recognize that the upfront costs of implementing LLMs can be higher due to their complexity and infrastructure requirements. We have also to add to this analysis the fast evolution of chatbots. It presents a challenge when interpreting conclusions from older studies and even possible replication in the near future. Early research in this field often relied on less sophisticated algorithms and limited computational resources, which constrained the functionality, accuracy, and scalability of chatbots. For example, earlier systems lacked the advanced natural language processing and adaptive learning capabilities that are now available in modern large language models (LLMs) that are accessible to the wide public such as GPT-4 and Gemini. These newer models offer substantially improved contextual understanding and personalized interaction, enabling more effective applications in healthcare. As a result, while older studies provide valuable foundational insights, their relevance may be limited in the context of today's technological advancements. The lack of enough data from real-world applications of current-generation chatbots also poses a barrier to totally understand their potential. This discrepancy highlights the need for ongoing and continuously update research to evaluate the impact of brand new chatbots and ensure that conclusions drawn from earlier studies are appropriately contextualized.

As evidenced by studies on COVID-19 management and health education, chatbots can play a critical role in public health responses and preventive care. The studies by Mellah et al.³⁵ and Okonkwo et al.³⁷ highlight how chatbots can be rapidly deployed to support public health efforts and very useful disseminating information, assessing symptoms, and managing patient inquiries such as vaccination status. Chatbots can also play a key role in health education. Goldnadel et al.²¹ and Peng et al.³⁸ showed promising results at this regard in educating the public on critical health issues, promoting preventive measures, and reducing the spread of misinformation.

Despite their benefits, the deployment of chatbots in healthcare comes with challenges such as ensuring data

privacy, addressing ethical concerns, and managing user trust. Some studies like the one from Suharwardy et al.²⁸ reveal some limitations in their study concerning baseline conditions of users. Additionally, Hakani et al.³³ noted the importance of improving the accuracy of their depression predictive model in the case of the lack of extensive data. These issues need to be carefully considered and addressed to fully realize the potential of chatbots in healthcare. Future research should explore the long-term impact of chatbot interventions on patient outcomes, the integration of chatbot technology with electronic health records, and the development of more sophisticated AI models to enhance chatbot capabilities. Additionally, cross-disciplinary collaboration might foster innovative solutions and address the current limitations.

Despite the strengths of this review, some limitations should be noted. First, the review includes studies published in English and Spanish, which may introduce some language bias and limit the generalizability of the findings. Although most of the newest advances in LLM and chatbots may be published in English. Secondly, although we include articles published within the last five years to ensure updated evidence, some of the first studies that could offer additional insights in the evolution of LLM and chatbots may have been not included in our paper. Thirdly, the heterogeneity of the included studies in terms of methodologies, sample sizes, and outcomes makes it challenging to draw definitive conclusions and propose policy recommendations. Furthermore, many studies provided limited details on the cost-effectiveness and real-world implementation challenges of chatbot technologies. Lastly, as chatbot technology continues to evolve rapidly, some findings from included studies may already be outdated, being necessary continuous updates to capture the impact of emerging innovations. Future research should address these limitations by incorporating broader language inclusivity, longitudinal assessments, and real-world data to provide a more comprehensive evaluation of chatbot applications in healthcare.

Conclusions

In summary, chatbots represent a transformative technology in healthcare, offering numerous benefits from increased accessibility and personalized care to cost savings and enhanced patient engagement and empowerment. Despite these benefits, challenges remain in integrating these tools into healthcare systems and assessing their long-term impact. In recent years, the development of LLMs, such as GPT-4 or Gemini among others, has significantly improved the effectiveness and accuracy of chatbots, enabling more fluid and contextualized interactions with patients. Additionally, recent advancements and optimizations in technological infrastructure, along with increasing market competition, have led to a substantial reduction in the costs

associated with using LLMs. This decrease in costs can enhance the cost-effectiveness of chatbots in healthcare settings, enabling their use not only in large hospital systems but also in smaller clinics and rural environments, where access to healthcare is limited. However, the high costs of implementation and operation of these technologies should not be overlooked, as they have been a barrier to their widespread adoption.

To fully leverage the potential of this technology, it will be crucial to continue fostering innovation, supporting robust research, and developing supportive policies that regulate and evaluate the use of chatbots to ensure they meet the required quality and safety standards. Additionally, regulatory frameworks must evolve to ensure that advancements in AI and LLMs are integrated ethically and efficiently into clinical practice, safeguarding patient privacy and data security.

Finally, policymakers should develop frameworks to regulate and evaluate the use of chatbots in healthcare, ensuring they meet quality and safety standards. Support for research, evaluation and development in this field can accelerate the adoption of chatbot technology leading to improved healthcare delivery and outcomes.

Acknowledgements: The authors acknowledge the Government of Cantabria for funding this research.

Contributorship: Conceptualization: DC, AD, and RM conceived the systematic review, defining the study's focus and objectives. Methodology: MB developed the search strategy, inclusion/exclusion criteria, and designed the methodology for study selection and data extraction. Data collection: DC, PL, and JL were responsible for data collection, conducting the search and selection of relevant studies. Data analysis: DC and MB. Writing—original draft preparation: The initial draft of the manuscript was written by MB, PL, and JL. Writing—review & editing: The review and editing of the manuscript were done by DC, AD, RM, and FP. Supervision: DC and MB supervised the entire process, ensuring the consistency and quality of the manuscript.

Declaration of conflicting interests: The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval: Ethical approval to report this research was obtained from «COMITÉ DE ÉTICA DE LA INVESTIGACIÓN CON MEDICAMENTOS DE CANTABRIA» (11/2024–10/05/2024).

Funding: The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by the Government of Cantabria (Grant Number: SUBVTC-2023-0021). This study is also funded by the European Commission in the Horizon H2020 scheme, awarded to the TIMELY project (Grant agreement ID: 101017424).

ORCID iDs: Abraham Delgado  <https://orcid.org/0000-0002-7665-8605>

Javier Lera  <https://orcid.org/0000-0002-0367-265X>

Marina Barreda  <https://orcid.org/0000-0002-5952-6758>

References

1. Mehraeen E, Mehrtak M, SeyedAlinaghi S, et al. Technology in the Era of COVID-19: a systematic review of current evidence. *Infect Disord - Drug Targets [Internet]* 2022 Jun; 22. <https://www.eurekaselect.com/202551/article>.
2. Moutsana Tapolin F, Liaskos J, Zoulias E, et al. A Conversational Web-Based Chatbot to Disseminate COVID-19 Advisory Information. In 2023. Available from: <https://ebooks.iospress.nl/doi/10.3233/SHTI230538>.
3. Mehraeen E, SeyedAlinaghi S, Heydari M, et al. Telemedicine technologies and applications in the era of COVID-19 pandemic: a systematic review. *Health Inform J [Internet]* 2023 Apr 19; 29. <https://journals.sagepub.com/doi/10.1177/14604582231167431>.
4. LeBrón AMW, Schulz AJ, Mentz G, et al. Impact of change over time in self-reported discrimination on blood pressure: implications for inequities in cardiovascular risk for a multi-racial urban community. *Ethn Health [Internet]* 2020 Apr 2; 25: 1. [cited 2022 Jun 20] Available from: [/pmc/articles/PMC6054822/](https://pubmed.ncbi.nlm.nih.gov/346054822/).
5. Almalki M and Azeez F. Health chatbots for fighting COVID-19: a scoping review. *Acta Inform Medica [Internet]* 2020; 28: 241. <https://www.bibliomed.org/?mno=35112>.
6. World Health Organization (WHO). Ethics and governance of artificial intelligence for health [Internet]. 2021. Available from: <https://www.who.int/publications/i/item/9789240029200>.
7. Aggarwal A, Tam CC, Wu D, et al. Artificial intelligence-based chatbots for promoting health behavioral changes: systematic review. *J Med Internet Res [Internet]* 2023 Feb 24; 25: e40789. <https://www.jmir.org/2023/1/e40789>.
8. Laranjo L, Dunn AG, Tong HL, et al. Conversational agents in healthcare: a systematic review. *J Am Med Informatics Assoc [Internet]* 2018 Sep 1; 25: 1248–1258. <https://academic.oup.com/jamia/article/25/9/1248/5052181>.
9. Oh YJ, Zhang J, Fang M-L, et al. A systematic review of artificial intelligence chatbots for promoting physical activity, healthy diet, and weight loss. *Int J Behav Nutr Phys Act [Internet]* 2021 Dec 11; 18: 160. <https://ijbnpa.biomedcentral.com/articles/10.1186/s12966-021-01224-6>.
10. Afsahi AM, Seyed Alinaghi SA, Molla A, et al. Chatbots utility in healthcare industry: an Umbrella review. *Front Heal Informatics [Internet]* 2024 Apr 7; 13: 200. <http://ijmi.ir/index.php/IJMI/article/view/561>.
11. Zhu Y, Wang R and Pu C. “I am chatbot, your virtual mental health adviser.” what drives citizens’ satisfaction and continuance intention toward mental health chatbots during the COVID-19 pandemic? An empirical study in China. *Digit Heal [Internet]* 2022 Jan 30; 8: 20552076221090031. <https://journals.sagepub.com/doi/10.1177/20552076221090031>.
12. Lim SM, Shiau CWC, Cheng LJ, et al. Chatbot-delivered psychotherapy for adults with depressive and anxiety symptoms: a systematic review and meta-regression. *Behav Ther [Internet]* 2022 Mar; 53: 334–347. <https://linkinghub.elsevier.com/retrieve/pii/S0005789421001258>.

13. Ahmed A, Hassan A, Aziz S, et al. Chatbot features for anxiety and depression: a scoping review. *Health Informatics J [Internet]* 2023 Jan 24; 29. <https://journals.sagepub.com/doi/10.1177/14604582221146719>
14. Kurniawan MH, Handiyani H, Nuraini T, et al. A systematic review of artificial intelligence-powered (AI-powered) chatbot intervention for managing chronic illness. *Ann Med [Internet]* 2024 Dec 31; 56. <https://www.tandfonline.com/doi/full/10.1080/07853890.2024.2302980>
15. Bin Sawad A, Narayan B, Alnefaie A, et al. A systematic review on healthcare artificial intelligent conversational agents for chronic conditions. *Sensors [Internet]* 2022 Mar 29; 22: 2625. <https://www.mdpi.com/1424-8220/22/7/2625>
16. Irfan N and Zafar S. A Self Diagnosis Medical Chatbot Using Sklearn. In 2023. p. 197–208. Available from: https://link.springer.com/10.1007/978-981-99-3716-5_18.
17. Bergmo TS. How to measure costs and benefits of eHealth interventions: an overview of methods and frameworks. *J Med Internet Res [Internet]* 2015 Nov 9; 17: e254. <http://www.jmir.org/2015/11/e254/>
18. Anjum K, Sameer M and Kumar S. AI Enabled NLP based Text to Text Medical Chatbot. In: 2023 3rd International Conference on Innovative Practices in Technology and Management (ICIPTM) [Internet], IEEE; 2023. p. 1–5. Available from: <https://ieeexplore.ieee.org/document/10117966/>.
19. Bhuvanesh Shathyan R, Begam MF, Jashwanth K, et al. Knowledge Graph Based Medical Chatbot building. In: 2023 4th IEEE Global Conference for Advancement in Technology (GCAT) [Internet], IEEE; 2023. p. 1–6. Available from: <https://ieeexplore.ieee.org/document/10353415/>.
20. Booth F, Potts C, Bond R, et al. A mental health and well-being chatbot: user event log analysis. *JMIR mHealth UHealth [Internet]* 2023 Jul 6; 11: e43052. <https://mhealth.jmir.org/2023/1/e43052>
21. Monteiro MG, Pantani D, Pinsky I, et al. Using the pan American health organization digital conversational agent to educate the public on alcohol use and health: preliminary analysis. *JMIR Form Res [Internet]* 2023 Apr 6; 7: e43165. <https://formative.jmir.org/2023/1/e43165>
22. Hasnain M. ChatGPT applications and challenges in controlling monkey pox in Pakistan. *Ann Biomed Eng [Internet]* 2023 Sep 6; 51: 1889–1891. <https://link.springer.com/10.1007/s10439-023-03231-z>
23. Kaywan P, Ahmed K, Ibaida A, et al. Early detection of depression using a conversational AI bot: a non-clinical trial. *Saha S, Editor. PLoS One [Internet]* 2023 Feb 3; 18: e0279743. <https://dx.plos.org/10.1371/journal.pone.0279743>
24. Potts C, Lindström F, Bond R, et al. A multilingual digital mental health and well-being chatbot (ChatPal): pre-post multicenter intervention study. *J Med Internet Res [Internet]* 2023 Jul 6; 25: e43051. <https://www.jmir.org/2023/1/e43051>
25. Prakasam S, Balakrishnan N, Kirthickram T R, et al. Design and Development of AI-Powered Healthcare WhatsApp Chatbot. In: 2023 2nd International Conference on Vision Towards Emerging Trends in Communication and Networking Technologies (ViTECoN) [Internet], IEEE; 2023. p. 1–6. Available from: <https://ieeexplore.ieee.org/document/10157423/>.
26. Rekik S, Elamine M and Belguith LH. A Medical Chatbot for Tunisian Dialect using a Rule-Based and Machine Learning Approach. In: 2023 20th ACS/IEEE International Conference on Computer Systems and Applications (AICCSA) [Internet], IEEE; 2023. p. 1–7. Available from: <https://ieeexplore.ieee.org/document/10479249/>.
27. Sabour S, Zhang W, Xiao X, et al. A chatbot for mental health support: exploring the impact of Emohaa on reducing mental distress in China. *Front Digit Heal [Internet]* 2023 May 4; 5. <https://www.frontiersin.org/articles/10.3389/fdgth.2023.1133987/full>.
28. Suharwardy S, Ramachandran M, Leonard SA, et al. Feasibility and impact of a mental health chatbot on post-partum mental health: a randomized controlled trial. *AJOG Glob Reports [Internet]* 2023 Aug; 3: 100165. <https://linkinghub.elsevier.com/retrieve/pii/S2666577823000060>
29. Upadhyaya P and Kaur G. Smart Multi-linguistic Health Awareness System using RASA Model. In: 2023 International Conference on Sustainable Computing and Smart Systems (ICSCSS) [Internet], IEEE; 2023. p. 922–7. Available from: <https://ieeexplore.ieee.org/document/10169275/>.
30. Vera MCS and Palaoag TD. Implementation of a smarter herbal medication delivery system employing an ai-powered chatbot. *Int J Adv Comput Sci Appl [Internet]* 2023; 14. <http://thesai.org/Publications/ViewPaper?Volume=14&Issue=3&Code=IJACSA&SerialNo=58>.
31. Bhangdia Y, Bhansali R, Chaudhari N, et al. Speech Emotion Recognition and Sentiment Analysis based Therapist Bot. In: 2021 Third International Conference on Inventive Research in Computing Applications (ICIRCA) [Internet], IEEE; 2021. p. 96–101. Available from: <https://ieeexplore.ieee.org/document/9544671/>.
32. Vu B, Bruchhaus S, Moorhead A, et al. Towards Continuous Professional Monitoring of Health Status Based on Energetic Balancing. In: 2022 IEEE International Workshop on Sport, Technology and Research (STAR) [Internet], IEEE; 2022. p. 72–7. Available from: <https://ieeexplore.ieee.org/document/9859599/>.
33. Hakani R, Patil S, Patil S, et al. Revivify: A Depression Detection and Control System using Tweets and Automated Chatbot. In: 2022 IEEE World Conference on Applied Intelligence and Computing (AIC) [Internet], IEEE; 2022. p. 796–801. Available from: <https://ieeexplore.ieee.org/document/9848978/>.
34. Mehta D and Singh S. Medical Insurance Virtual Assistant: Vaidya. In: 2022 5th International Conference on Contemporary Computing and Informatics (IC3I) [Internet], IEEE; 2022. p. 584–9. Available from: <https://ieeexplore.ieee.org/document/10072984/>.
35. Mellah Y, Bouchentouf T, Rahmoun N, et al. Bilingual Chatbot For COVID-19 Detection Based on Symptoms Using Rasa NLU. In: 2022 2nd International Conference on Innovative Research in Applied Science, Engineering and Technology (IRASET) [Internet], IEEE; 2022. p. 1–5. Available from: <https://ieeexplore.ieee.org/document/9738103/>.
36. Natsheh E and Javed M. Implementing artificial intelligence-based COVID-19 chatbot in the kingdom of Bahrain. *Int J Intell Syst Appl Eng [Internet]* 2022; 10: 10–17. <https://ijisae.org/index.php/IJISAE/article/view/2131>.
37. Okonkwo CW, Amusa LB and Twinomurinzi H. COVID-Bot, an intelligent system for COVID-19 vaccination screening:

- design and development. *JMIR Form Res [Internet]* 2022 Oct 27; 6: e39157. <https://formative.jmir.org/2022/10/e39157>
38. Peng ML, Wickersham JA, Altice FL, et al. Formative evaluation of the acceptance of HIV prevention artificial intelligence chatbots by men who have sex with men in Malaysia: focus group study. *JMIR Form Res [Internet]* 2022 Oct 6; 6: e42055. <https://formative.jmir.org/2022/10/e42055>
 39. Bendig E, Meißner D, Erb B, et al. Study protocol of a randomised controlled trial on SISU, a software agent providing a brief self-help intervention for adults with low psychological well-being. *BMJ Open [Internet]* 2021 Feb 8; 11: e041573. <https://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-2020-041573>
 40. Dhinakaran DA, Sathish T, Kowatsch T, et al. Public perceptions of diabetes, healthy living, and conversational agents in Singapore: Needs assessment. *JMIR Form Res [Internet]* 2021 Nov 11; 5: e30435. <https://formative.jmir.org/2021/11/e30435>
 41. Gabrielli S, Rizzi S, Bassi G, et al. Engagement and effectiveness of a healthy-coping intervention via chatbot for university students during the COVID-19 pandemic: mixed methods proof-of-concept study. *JMIR mHealth UHealth [Internet]* 2021 May 28; 9: e27965. <https://mhealth.jmir.org/2021/5/e27965>
 42. Omarov B, Zhumanov Z, Gumar A, et al. Artificial intelligence enabled mobile chatbot psychologist using AIML and cognitive behavioral therapy. *Int J Adv Comput Sci Appl [Internet]* 2023; 14. <http://thesai.org/Publications/ViewPaper?Volume=14&Issue=6&Code=IJACSA&SerialNo=16>
 43. Larbi D, Gabarron E and Denecke K. Social Media Chatbot for Increasing Physical Activity: Usability Study. In 2021. Available from: <https://ebooks.iospress.nl/doi/10.3233/SHTI210604>.
 44. Ashan Pivithuru KW, Srinivasan A, Wijeratnam A, et al. E-patient Card: An Integrated Electronic Health Recording System for Patient. In: 2021 3rd International Conference on Advancements in Computing (ICAC) [Internet], IEEE; 2021. p. 395–400. Available from: <https://ieeexplore.ieee.org/document/9671145/>.
 45. Daley K, Hungerbuehler I, Cavanagh K, et al. Preliminary evaluation of the engagement and effectiveness of a mental health chatbot. *Front Digit Heal [Internet]* 2020 Nov 30; 2. <https://www.frontiersin.org/articles/10.3389/fdgth.2020.576361/full>.
 46. Srivastava P and Singh N. Automatized Medical Chatbot (Medibot). In: 2020 International Conference on Power Electronics & IoT Applications in Renewable Energy and its Control (PARC) [Internet], IEEE; 2020. p. 351–4. Available from: <https://ieeexplore.ieee.org/document/9087096/>.
 47. Vera MCS and Palaoag TD. MedPlantBot: AI Chatbot Architectural Design Framework for Responsible Use of Medicinal Plants. In: 2023 IEEE International Conference on ICT in Business Industry & Government (ICTBIG) [Internet], IEEE; 2023. p. 1–8. Available from: <https://ieeexplore.ieee.org/document/10456018/>.
 48. Nazi Z Al and Peng W. Large language models in healthcare and medical domain: a review. *Informatics [Internet]* 2024 Aug 7; 11: 57. <https://www.mdpi.com/2227-9709/11/3/57>
 49. Haltaufderheide J and Ranisch R. The ethics of ChatGPT in medicine and healthcare: a systematic review on large language models (LLMs). *npj Digit Med [Internet]* 2024 Jul 8; 7: 183. <https://www.nature.com/articles/s41746-024-01157-x>
 50. Clark M and Bailey S. Chatbots in health care: connecting patients to information. *Can J Heal Technol [Internet]* 2024 Jan 22; 4. <https://www.canjhealthtechnol.ca/index.php/cjht/article/view/EH0122>
 51. Jiang F, Jiang Y, Zhi H, et al. Artificial intelligence in healthcare: past, present and future. *Stroke Vasc Neurol [Internet]* 2017 Dec; 2: 230–243. <https://svn.bmj.com/lookup/doi/10.1136/svn-2017-000101>
 52. Bohr A and Memarzadeh K. The rise of artificial intelligence in healthcare applications. In: *Artificial intelligence in healthcare [internet]*. Elsevier, 2020, pp.25–60. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780128184387000022>.
 53. Topol E. *Deep medicine: How artificial intelligence can make healthcare human again*. New York: Basic Books, 2019.