

Search for Production of Heavy Particles Decaying to Top Quarks and Invisible Particles in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

T. Aaltonen,²² B. Álvarez González,^{10,x} S. Amerio,^{42a} D. Amidei,³³ A. Anastassov,³⁷ A. Annovi,¹⁸ J. Antos,¹³ G. Apollinari,¹⁶ J. A. Appel,¹⁶ A. Apresyan,⁴⁷ T. Arisawa,⁵⁷ A. Artikov,¹⁴ J. Asaadi,⁵² W. Ashmanskas,¹⁶ B. Auerbach,⁶⁰ A. Aurisano,⁵² F. Azfar,⁴¹ W. Badgett,¹⁶ A. Barbaro-Galtieri,²⁷ V. E. Barnes,⁴⁷ B. A. Barnett,²⁴ P. Barria,^{45b,45a} P. Bartos,¹³ M. Baucé,^{42b,42a} G. Bauer,³¹ F. Bedeschi,^{61a} D. Beecher,²⁹ S. Behari,²⁴ G. Bellettini,^{61b,61a} J. Bellinger,⁵⁹ D. Benjamin,¹⁵ A. Beretvas,¹⁶ A. Bhatti,⁴⁹ M. Binkley,^{16,a} D. Bisello,^{42b,42a} I. Bizjak,^{29,bb} K. R. Bland,⁵ B. Blumenfeld,²⁴ A. Bocci,¹⁵ A. Bodek,⁴⁸ D. Bortoletto,⁴⁷ J. Boudreau,⁴⁶ A. Boveia,¹² B. Brau,^{16,b} L. Brigliadori,^{6b,6a} A. Brisuda,¹³ C. Bromberg,³⁴ E. Brucken,²² M. Bucciantonio,^{45b,45a} J. Budagov,¹⁴ H. S. Budd,⁴⁸ S. Budd,²³ K. Burkett,¹⁶ G. Busetto,^{42b,42a} P. Bussey,²⁰ A. Buzatu,³² C. Calancha,³⁰ S. Camarda,⁴ M. Campanelli,³⁴ M. Campbell,³³ F. Canelli,^{12,16} A. Canepa,⁴⁴ B. Carls,²³ D. Carlsmith,⁵⁹ R. Carosi,^{45a} S. Carrillo,^{17,1} S. Carron,¹⁶ B. Casal,¹⁰ M. Casarsa,¹⁶ A. Castro,^{6b,6a} P. Catastini,¹⁶ D. Cauz,^{53a} V. Cavaliere,^{45b,45a} M. Cavalli-Sforza,⁴ A. Cerri,^{27,g} L. Cerrito,^{29,r} Y. C. Chen,¹ M. Chertok,⁷ G. Chiarelli,^{45a} G. Chlachidze,¹⁶ F. Chlebana,¹⁶ K. Cho,²⁶ D. Chokheli,¹⁴ J. P. Chou,²¹ W. H. Chung,⁵⁹ Y. S. Chung,⁴⁸ C. I. Ciobanu,⁴³ M. A. Ciocci,^{45b,45c} A. Clark,¹⁹ G. Compostella,^{42b,42a} M. E. Convery,¹⁶ J. Conway,⁷ M. Corbo,⁴³ M. Cordelli,¹⁸ C. A. Cox,⁷ D. J. Cox,⁷ F. Crescioli,^{45b,45a} C. Cuenca Almenar,⁶⁰ J. Cuevas,^{10,x} R. Culbertson,¹⁶ D. Dagenhart,¹⁶ N. d'Ascenzo,^{43,v} M. Datta,¹⁶ P. de Barbaro,⁴⁸ S. De Cecco,^{50a} G. De Lorenzo,⁴ M. Dell'Orso,^{45b,45a} C. Deluca,⁴ L. Demortier,⁴⁹ J. Deng,^{15,d} M. Deninno,^{6a} F. Devoto,²² M. d'Errico,^{42b,42a} A. Di Canto,^{45b,45a} B. Di Ruzza,^{45a} J. R. Dittmann,⁵ M. D'Onofrio,²⁸ S. Donati,^{45b,45a} P. Dong,¹⁶ M. Dorigo,^{53a} T. Dorigo,^{42a} K. Ebina,⁵⁷ A. Elagin,⁵² A. Eppig,³³ R. Erbacher,⁷ D. Errede,²³ S. Errede,²³ N. Ershaidat,^{43,aa} R. Eusebi,⁵² H. C. Fang,²⁷ S. Farrington,⁴¹ M. Feindt,²⁵ J. P. Fernandez,³⁰ C. Ferrazza,^{45b,45a} R. Field,¹⁷ G. Flanagan,^{47,t} R. Forrest,⁷ M. J. Frank,⁵ M. Franklin,²¹ J. C. Freeman,¹⁶ Y. Funakoshi,⁵⁷ I. Furic,¹⁷ M. Gallinaro,⁴⁹ J. Galyardt,¹¹ J. E. Garcia,¹⁹ A. F. Garfinkel,⁴⁷ P. Garosi,^{45b,45a} H. Gerberich,²³ E. Gerchtein,¹⁶ S. Giagu,^{50b,50a} V. Giakoumopoulou,³ P. Giannetti,^{45a} K. Gibson,⁴⁶ C. M. Ginsburg,¹⁶ N. Giokaris,³ P. Giromini,¹⁸ M. Giunta,^{45a} G. Giurgiu,²⁴ V. Glagolev,¹⁴ D. Glenzinski,¹⁶ M. Gold,³⁶ D. Goldin,⁵² N. Goldschmidt,¹⁷ A. Golossanov,¹⁶ G. Gomez,¹⁰ G. Gomez-Ceballos,³¹ M. Goncharov,³¹ O. González,³⁰ I. Gorelov,³⁶ A. T. Goshaw,¹⁵ K. Goulianos,⁴⁹ S. Grinstein,⁴ C. Grosso-Pilcher,¹² R. C. Group,⁵⁶ J. Guimaraes da Costa,²¹ Z. Gunay-Unalan,³⁴ C. Haber,²⁷ S. R. Hahn,¹⁶ E. Halkiadakis,⁵¹ A. Hamaguchi,⁴⁰ J. Y. Han,⁴⁸ F. Happacher,¹⁸ K. Hara,⁵⁴ D. Hare,⁵¹ M. Hare,⁵⁵ R. F. Harr,⁵⁸ K. Hatakeyama,⁵ C. Hays,⁴¹ M. Heck,²⁵ J. Heinrich,⁴⁴ M. Herndon,⁵⁹ S. Hewamanage,⁵ D. Hidas,⁵¹ A. Hocker,¹⁶ W. Hopkins,^{16,h} D. Horn,²⁵ S. Hou,¹ R. E. Hughes,³⁸ M. Hurwitz,¹² U. Husemann,⁶⁰ N. Hussain,³² M. Hussein,³⁴ J. Huston,³⁴ G. Introzzi,^{45a} M. Iori,^{50b,50a} A. Ivanov,^{7,p} E. James,¹⁶ D. Jang,¹¹ B. Jayatilaka,¹⁵ E. J. Jeon,²⁶ M. K. Jha,^{6a} S. Jindariani,¹⁶ W. Johnson,⁷ M. Jones,⁴⁷ K. K. Joo,²⁶ S. Y. Jun,¹¹ T. R. Junk,¹⁶ T. Kamon,⁵² P. E. Karchin,⁵⁸ Y. Kato,^{40,o} W. Ketchum,¹² J. Keung,⁴⁴ V. Khotilovich,⁵² B. Kilminster,¹⁶ D. H. Kim,²⁶ H. S. Kim,²⁶ H. W. Kim,²⁶ J. E. Kim,²⁶ M. J. Kim,¹⁸ S. B. Kim,²⁶ S. H. Kim,⁵⁴ Y. K. Kim,¹² N. Kimura,⁵⁷ M. Kirby,¹⁶ S. Klimenko,¹⁷ K. Kondo,⁵⁷ D. J. Kong,²⁶ J. Konigsberg,¹⁷ A. V. Kotwal,¹⁵ M. Krepes,²⁵ J. Kroll,⁴⁴ D. Krop,¹² N. Krumnack,^{5,m} M. Kruse,¹⁵ V. Krutelyov,^{52,e} T. Kuhr,²⁵ M. Kurata,⁵⁴ S. Kwang,¹² A. T. Laasanen,⁴⁷ S. Lami,^{45a} S. Lammel,¹⁶ M. Lancaster,²⁹ R. L. Lander,⁷ K. Lannon,^{38,w} A. Lath,⁵¹ G. Latino,^{45b,45a} T. LeCompte,² E. Lee,⁵² H. S. Lee,¹² J. S. Lee,²⁶ S. W. Lee,^{52,y} S. Leo,^{45b,45a} S. Leone,^{45a} J. D. Lewis,¹⁶ A. Limosani,^{15,s} C.-J. Lin,²⁷ J. Linacre,⁴¹ M. Lindgren,¹⁶ E. Lipeles,⁴⁴ A. Lister,¹⁹ D. O. Litvintsev,¹⁶ C. Liu,⁴⁶ Q. Liu,⁴⁷ T. Liu,¹⁶ S. Lockwitz,⁶⁰ N. S. Lockyer,⁴⁴ A. Loginov,⁶⁰ D. Lucchesi,^{42b,42a} J. Lueck,²⁵ P. Lujan,²⁷ P. Lukens,¹⁶ G. Lungu,⁴⁹ J. Lys,²⁷ R. Lysak,¹³ R. Madrak,¹⁶ K. Maeshima,¹⁶ K. Makhoul,³¹ P. Maksimovic,²⁴ S. Malik,⁴⁹ G. Manca,^{28,c} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁷ C. Marino,²⁵ M. Martínez,⁴ R. Martínez-Ballarín,³⁰ P. Mastrandrea,^{50a} M. Mathis,²⁴ M. E. Mattson,⁵⁸ P. Mazzanti,^{6a} K. S. McFarland,⁴⁸ P. McIntyre,⁵² R. McNulty,^{28,j} A. Mehta,²⁸ P. Mehtala,²² A. Menzione,^{45a} C. Mesropian,⁴⁹ T. Miao,¹⁶ D. Mietlicki,³³ A. Mitra,¹ H. Miyake,⁵⁴ S. Moed,²¹ N. Moggi,^{6a} M. N. Mondragon,^{16,1} C. S. Moon,²⁶ R. Moore,¹⁶ M. J. Morello,¹⁶ J. Morlock,²⁵ P. Movilla Fernandez,¹⁶ A. Mukherjee,¹⁶ Th. Muller,²⁵ P. Murat,¹⁶ M. Mussini,^{6b,6a} J. Nachtman,^{16,n} Y. Nagai,⁵⁴ J. Naganoma,⁵⁷ I. Nakano,³⁹ A. Napier,⁵⁵ J. Nett,⁵² C. Neu,⁵⁶ M. S. Neubauer,²³ J. Nielsen,^{27,f} L. Nodulman,² O. Norniella,²³ E. Nurse,²⁹ L. Oakes,⁴¹ S. H. Oh,¹⁵ Y. D. Oh,²⁶ I. Oksuzian,⁵⁶ T. Okusawa,⁴⁰ R. Orava,²² L. Ortolan,⁴ S. Pagan Griso,^{42b,42a} C. Pagliarone,^{53a} E. Palencia,^{10,g} V. Papadimitriou,¹⁶ A. A. Paramonov,² J. Patrick,¹⁶ G. Pauletta,^{53b,53a} M. Paulini,¹¹ C. Paus,³¹ D. E. Pellett,⁷ A. Penzo,^{53a} T. J. Phillips,¹⁵ G. Piacentino,^{45a} E. Pianori,⁴⁴ J. Pilot,³⁸ K. Pitts,²³ C. Plager,⁹ L. Pondrom,⁵⁹ K. Potamianos,⁴⁷ O. Poukhov,^{14,a} F. Prokoshin,^{14,z} A. Pronko,¹⁶ F. Ptohos,^{18,i} E. Pueschel,¹¹ G. Punzi,^{45b,45a} J. Pursley,⁵⁹ A. Rahaman,⁴⁶ V. Ramakrishnan,⁵⁹ N. Ranjan,⁴⁷ K. Rao,⁸

I. Redondo,³⁰ P. Renton,⁴¹ M. Rescigno,^{50a} F. Rimondi,^{6b,6a} L. Ristori,^{45a,16} A. Robson,²⁰ T. Rodrigo,¹⁰ T. Rodriguez,⁴⁴ E. Rogers,²³ S. Rolli,⁵⁵ R. Roser,¹⁶ M. Rossi,^{53a} F. Rubbo,¹⁶ F. Ruffini,^{45b,45a} A. Ruiz,¹⁰ J. Russ,¹¹ V. Rusu,¹⁶ A. Safonov,⁵² W. K. Sakumoto,⁴⁸ Y. Sakurai,⁵⁷ L. Santi,^{53b,53a} L. Sartori,^{45a} K. Sato,⁵⁴ V. Saveliev,^{43,v} A. Savoy-Navarro,⁴³ P. Schlabach,¹⁶ A. Schmidt,²⁵ E. E. Schmidt,¹⁶ M. P. Schmidt,^{60,a} M. Schmitt,³⁷ T. Schwarz,⁷ L. Scodellaro,¹⁰ A. Scribano,^{45b,45a} F. Scuri,^{45a} A. Sedov,⁴⁷ S. Seidel,³⁶ Y. Seiya,⁴⁰ A. Semenov,¹⁴ F. Sforza,^{45b,45a} A. Sfyrla,²³ S. Z. Shalhout,⁷ T. Shears,²⁸ P. F. Shepard,⁴⁶ M. Shimojima,^{54,u} S. Shiraishi,¹² M. Shochet,¹² I. Shreyber,³⁵ A. Simonenko,¹⁴ P. Sinervo,³² A. Sissakian,^{14,a} K. Sliwa,⁵⁵ J. R. Smith,⁷ F. D. Snider,¹⁶ A. Soha,¹⁶ S. Somalwar,⁵¹ V. Sorin,⁴ P. Squillacioti,¹⁶ M. Stancari,¹⁶ M. Stanitzki,⁶⁰ R. St. Denis,²⁰ B. Stelzer,³² O. Stelzer-Chilton,³² D. Stentz,³⁷ J. Strologas,³⁶ G. L. Strycker,³³ Y. Sudo,⁵⁴ A. Sukhanov,¹⁷ I. Suslov,¹⁴ K. Takemasa,⁵⁴ Y. Takeuchi,⁵⁴ J. Tang,¹² M. Tecchio,³³ P. K. Teng,¹ J. Thom,^{16,h} J. Thome,¹¹ G. A. Thompson,²³ E. Thomson,⁴⁴ P. Tito-Guzmán,³⁰ S. Tkaczyk,¹⁶ D. Toback,⁵² S. Tokar,¹³ K. Tollefson,³⁴ T. Tomura,⁵⁴ D. Tonelli,¹⁶ S. Torre,¹⁸ D. Torretta,¹⁶ P. Totaro,^{42a} M. Trovato,^{45b,45a} Y. Tu,⁴⁴ F. Ukegawa,⁵⁴ S. Uozumi,²⁶ A. Varganov,³³ F. Vázquez,^{17,1} G. Velev,¹⁶ C. Vellidis,³ M. Vidal,³⁰ I. Vila,¹⁰ R. Vilar,¹⁰ J. Vizán,¹⁰ M. Vogel,³⁶ G. Volpi,^{45b,45a} P. Wagner,⁴⁴ R. L. Wagner,¹⁶ T. Wakisaka,⁴⁰ R. Wallny,⁹ S. M. Wang,¹ A. Warburton,³² D. Waters,²⁹ M. Weinberger,⁵² W. C. Wester III,¹⁶ B. Whitehouse,⁵⁵ D. Whiteson,^{44,d} A. B. Wicklund,² E. Wicklund,¹⁶ S. Wilbur,¹² F. Wick,²⁵ H. H. Williams,⁴⁴ J. S. Wilson,³⁸ P. Wilson,¹⁶ B. L. Winer,³⁸ P. Wittich,^{16,h} S. Wolbers,¹⁶ H. Wolfe,³⁸ T. Wright,³³ X. Wu,¹⁹ Z. Wu,⁵ K. Yamamoto,⁴⁰ J. Yamaoka,¹⁵ T. Yang,¹⁶ U. K. Yang,^{12,q} Y. C. Yang,²⁶ W.-M. Yao,²⁷ G. P. Yeh,¹⁶ K. Yi,^{16,n} J. Yoh,¹⁶ K. Yorita,⁵⁷ T. Yoshida,^{40,k} G. B. Yu,¹⁵ I. Yu,²⁶ S. S. Yu,¹⁶ J. C. Yun,¹⁶ A. Zanetti,^{53a} Y. Zeng,¹⁵ and S. Zucchelli^{6b,6a,k}

(CDF Collaboration)

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*²*Argonne National Laboratory, Argonne, Illinois 60439, USA*³*University of Athens, 157 71 Athens, Greece*⁴*Institut de Física d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*⁵*Baylor University, Waco, Texas 76798, USA*^{6a}*Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy*^{6b}*University of Bologna, I-40127 Bologna, Italy*⁷*University of California, Davis, Davis, California 95616, USA*⁸*University of California, Irvine, Irvine, California 92697, USA*⁹*University of California, Los Angeles, Los Angeles, California 90024, USA*¹⁰*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*¹¹*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*¹²*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*¹³*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*¹⁴*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*¹⁵*Duke University, Durham, North Carolina 27708, USA*¹⁶*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*¹⁷*University of Florida, Gainesville, Florida 32611, USA*¹⁸*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*¹⁹*University of Geneva, CH-1211 Geneva 4, Switzerland*²⁰*Glasgow University, Glasgow G12 8QQ, United Kingdom*²¹*Harvard University, Cambridge, Massachusetts 02138, USA*²²*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*²³*University of Illinois, Urbana, Illinois 61801, USA*²⁴*The Johns Hopkins University, Baltimore, Maryland 21218, USA*²⁵*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*²⁶*Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea*²⁷*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*²⁸*University of Liverpool, Liverpool L69 7ZE, United Kingdom*²⁹*University College London, London WC1E 6BT, United Kingdom*³⁰*Centro de Investigaciones Energéticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*³¹*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

- ³²*Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3*
- ³³*University of Michigan, Ann Arbor, Michigan 48109, USA*
- ³⁴*Michigan State University, East Lansing, Michigan 48824, USA*
- ³⁵*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- ³⁶*University of New Mexico, Albuquerque, New Mexico 87131, USA*
- ³⁷*Northwestern University, Evanston, Illinois 60208, USA*
- ³⁸*The Ohio State University, Columbus, Ohio 43210, USA*
- ³⁹*Okayama University, Okayama 700-8530, Japan*
- ⁴⁰*Osaka City University, Osaka 588, Japan*
- ⁴¹*University of Oxford, Oxford OX1 3RH, United Kingdom*
- ^{42a}*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- ^{42b}*University of Bologna, I-40127 Bologna, Italy*
- ⁴³*LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*
- ⁴⁴*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ^{45a}*Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy*
- ^{45b}*University of Pisa, I-56127 Pisa, Italy*
- ^{45c}*University of Siena, I-56127 Pisa, Italy*
- ^{45d}*Scuola Normale Superiore, I-56127 Pisa, Italy*
- ⁴⁶*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- ⁴⁷*Purdue University, West Lafayette, Indiana 47907, USA*
- ⁴⁸*University of Rochester, Rochester, New York 14627, USA*
- ⁴⁹*The Rockefeller University, New York, New York 10065, USA*
- ^{50a}*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, I-00185 Roma, Italy*
- ^{50b}*Sapienza Università di Roma, I-00185 Roma, Italy*
- ⁵¹*Rutgers University, Piscataway, New Jersey 08855, USA*
- ⁵²*Texas A&M University, College Station, Texas 77843, USA*
- ^{53a}*Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, I-33100 Udine, Italy*
- ^{53b}*University of Trieste/Udine, I-33100 Udine, Italy*
- ⁵⁴*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁵⁵*Tufts University, Medford, Massachusetts 02155, USA*
- ⁵⁶*University of Virginia, Charlottesville, Virginia 22906, USA*
- ⁵⁷*Waseda University, Tokyo 169, Japan*
- ⁵⁸*Wayne State University, Detroit, Michigan 48201, USA*
- ⁵⁹*University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁶⁰*Yale University, New Haven, Connecticut 06520, USA*
- ^{61a}*Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy*
- ^{61b}*University of Pisa, I-56127 Pisa, Italy*

(Received 12 March 2011; published 12 May 2011)

We present a search for a new particle T' decaying to top quark via $T' \rightarrow t + X$, where X is an invisible particle. In a data sample with 4.8 fb^{-1} of integrated luminosity collected by the CDF II detector at Fermilab in $p\bar{p}$ collisions with $\sqrt{s} = 1.96 \text{ TeV}$, we search for pair production of T' in the lepton + jets channel, $p\bar{p} \rightarrow t\bar{t} + X + X \rightarrow \ell\nu bqq'b + X + X$. We interpret our results primarily in terms of a model where T' are exotic fourth generation quarks and X are dark matter particles. Current direct and indirect bounds on such exotic quarks restrict their masses to be between 300 and 600 GeV/c^2 , the dark matter particle mass being anywhere below $m_{T'}$. The data are consistent with standard model expectations, and we set 95% confidence level limits on the generic production of $T'\bar{T}' \rightarrow t\bar{t} + X + X$. For the dark matter model we exclude T' at 95% confidence level up to $m_{T'} = 360 \text{ GeV}/c^2$ for $m_X \leq 100 \text{ GeV}/c^2$.

DOI: 10.1103/PhysRevLett.106.191801

PACS numbers: 14.65.Jk, 12.60.-i, 13.85.Rm, 14.80.-j

Despite an intensive program of research [1], the precise nature of dark matter remains elusive, though it is clear that it must be long-lived on cosmological time scales. Such a long lifetime could be due to a conserved charge under an unbroken symmetry. However, none of the unbroken symmetries of the standard model (SM) suffice to provide such

a charge, so it follows that dark matter must be charged under a new, unbroken symmetry. The prospects of creating dark matter at particle colliders are excellent, but only if the dark matter particles X couple to standard model particles directly or indirectly. One potential mechanism is via a connector particle Y , which carries SM charges so

that it can be produced at particle colliders as well as carrying the new dark charge, so that it can decay to the dark matter particle, $Y \rightarrow f + X$, where f is a SM particle. One compelling recent model [2] uses an exotic fourth generation up-type quark T' as the connector particle, which decays to a top quark and dark matter, $T' \rightarrow t + X$. Current direct and indirect bounds on such exotic quarks restrict their masses to be between 300 and 600 GeV [2].

The pair production of such exotic quarks and their subsequent decay to top quarks and dark matter has a collider signal comprising of top-quark pairs ($t\bar{t}$) and missing transverse momentum (E_i) due to the invisible dark matter particles. These types of signals, in general, are of great interest as they appear in numerous new physics scenarios including many dark matter motivated models, little Higgs models with T -parity conservation [3] and models in which baryon and lepton numbers are gauge symmetries [4]. Supersymmetry, which includes a natural dark matter candidate and provides a framework for unification of the forces, also predicts a $t\bar{t} + E_i$ signal from the decay of a supersymmetric top \tilde{t} quark to a top quark and the lightest supersymmetric particle [5], $\tilde{t} \rightarrow t + \chi^0$. There are currently no experimental bounds on a new heavy particle Y decaying via $Y \rightarrow t + X$.

This Letter reports a search for such a generic signal $t\bar{t} + E_i$ via the pair production of a heavy new particle T' with prompt decay $T' \rightarrow t + X$. We consider the mode $p\bar{p} \rightarrow t\bar{t} + X + X \rightarrow WbWb + X + X$ in which one W decays leptonically (including τ decays to e or μ) and one decays hadronically to qq' , this decay mode allows for large branching ratios while suppressing SM backgrounds. Such a signal is similar to top-quark pair production and decay, but with additional missing transverse energy due to the invisible particles.

Events were recorded by CDF II [6], a general purpose detector designed to study collisions at the Fermilab Tevatron $p\bar{p}$ collider at $\sqrt{s} = 1.96$ TeV. A charged-particle tracking system immersed in a 1.4 T magnetic field consists of a silicon microstrip tracker and a drift chamber. Electromagnetic and hadronic calorimeters surrounding the tracking system measure particle energies and drift chambers located outside the calorimeters detect muons. We use a data sample corresponding to an integrated luminosity of 4.8 ± 0.3 fb $^{-1}$.

The data acquisition system is triggered by e or μ candidates [7] with transverse momentum p_T [8] greater than 18 GeV/ c . Electrons and muons are reconstructed offline and are selected if they have a pseudorapidity η [8] magnitude less than 1.1, $p_T \geq 20$ GeV/ c and satisfy the standard identification and isolation requirements [7]. Jets are reconstructed in the calorimeter using the JETCLU [9] algorithm with a clustering radius of 0.4 in azimuth-pseudorapidity space and corrected using standard techniques [10]. Jets are selected if they have $p_T \geq 15$ GeV/ c and $|\eta| < 2.4$.

Missing transverse momentum [11] is reconstructed using fully corrected calorimeter and muon information [7].

Production of T' pairs and their subsequent decays to top-quark pairs and two dark matter particles would appear as events with a charged lepton and missing transverse momentum from one leptonically decaying W and the two dark matter particles, and four jets from the two b quarks and the hadronic decay of the second W boson. We select events with at least one electron or muon, at least four jets, and large missing transverse momentum. The missing transverse energy in a signal event depends on the masses $m_{T'}$ and m_X , for each pair of signal masses we optimize for the minimum amount of missing transverse energy required (ranging from 100 GeV/ c to 160 GeV/ c).

We model the production and decay of T' pairs with MADGRAPH [12]. Additional radiation, hadronization and showering are described by PYTHIA [13]. The detector response for all simulated samples is modeled by the official CDF detector simulation.

The dominant SM background is top-quark pair production. We model this background using PYTHIA $t\bar{t}$ production with $m_t = 172.5$ GeV/ c^2 . We normalize the $t\bar{t}$ background to the NLO cross section [14], and confirm that it is well modeled by examining $t\bar{t}$ -dominated regions in the data.

The second dominant SM background process is the associated production of W boson and jets. Samples of simulated W + jets events with light- and heavy-flavor jets are generated using the ALPGEN [15] program, interfaced with the parton-shower model from PYTHIA. The W + jets samples are normalized to the measured W cross section, with an additional multiplicative factor for the relative contribution of heavy- and light-flavor jets, the standard technique in measuring the top-quark pair-production cross section [16]. Multijet background, in which a jet is misreconstructed as a lepton, is modeled using a jet-triggered sample normalized in a background-dominated region at low missing transverse momentum. The remaining backgrounds, single top and diboson production, are modeled using PYTHIA and normalized to next-to-leading order cross sections [17].

We differentiate the signal events from these backgrounds by comparing the reconstructed transverse mass of the leptonically decaying W candidate,

$$m_T^W \equiv m_T(p_T^\ell, \not{p}_T) = \sqrt{2|p_T^\ell| |\not{p}_T| (1 - \cos[\Delta\phi(p_T^\ell, \not{p}_T)])},$$

where p_T^ℓ is the transverse momentum of the lepton and \not{p}_T is the missing transverse momentum. In background events, the \not{p}_T comes primarily from the neutrino in $W \rightarrow \ell\nu$ decay, and m_T^W will show a strong peak at the W -boson mass. The signal event, $T' \rightarrow t + X$, has additional missing transverse momentum due to the invisible particles X and thus does not reconstruct the W mass in m_T^W . Figure 1 shows the m_T^W distributions of the backgrounds versus the signals.




FIG. 1. Reconstructed transverse mass of the W , m_T^W , for the standard model backgrounds, the observed data, and for three choices of $(m_{T'}, m_X)$.

We consider several sources of systematic uncertainty on both the background rates and distributions, as well as on the expectations for the signal. Each affects the expected sensitivity to new physics expressed as an expected cross-section upper limit in the no-signal assumption. The dominant systematic uncertainties are the jet energy scale [10], contributions from additional interactions, and descriptions of initial and final state radiation [18]. In each case, we treat the unknown underlying quantity as a nuisance parameter and measure the distortion of the m_T^W spectrum for positive and negative fluctuations. As mentioned before we optimize the minimum missing transverse energy required for each signal point, Table I compares the number of events expected with uncertainties for

TABLE I. Number of events, for example, signal points compared to backgrounds and data for two E_T cuts after initial selection is made.

Cut:	$E_T \geq 100 \text{ GeV}/c$	$E_T \geq 150 \text{ GeV}/c$
$T'T' \rightarrow ttXX$ [GeV/c ²]		
$m_{T'}, m_X = 300, 90$	$22.9^{+5.8}_{-4.7}$	$4.1^{+2.4}_{-2.1}$
$m_{T'}, m_X = 310, 80$	$22.6^{+4.9}_{-5.1}$	$6.4^{+2.3}_{-2.6}$
$m_{T'}, m_X = 330, 70$	$17.6^{+3.7}_{-3.6}$	$7.3^{+2.5}_{-2.4}$
$m_{T'}, m_X = 350, 1$	$13.1^{+2.7}_{-2.8}$	$6.7^{+2.0}_{-1.9}$
$t\bar{t}$	189^{+54}_{-50}	$26.3^{+11.6}_{-9.8}$
$W + \text{jets}$	105^{+31}_{-14}	$16.6^{+4.5}_{-2.1}$
Single top	1.86 ± 0.2	0.18 ± 0.02
Diboson	9.69 ± 0.1	1.53 ± 0.1
$Z + \text{jets}$	4.00 ± 0.4	0.46 ± 0.05
QCD	0.04 ± 0.01	0.04 ± 0.01
Total Background	310^{+80}_{-64}	45^{+14}_{-11}
Data	309	42

backgrounds and signals to data for two example missing transverse energy cuts.

We validate our modeling of the SM backgrounds in two background-dominated control regions. We validate our modeling of the large m_T^W region in events with high missing transverse energy and exactly three jets, and validate our modeling of four-jet events in events with small missing transverse energy ($< 100 \text{ GeV}/c$). Figure 2 shows good agreement of our background modeling with data in the control regions.

There is no evidence for the presence of $T' \rightarrow t + X$ events in the data. We calculate 95% C.L. upper limits on the $T' \rightarrow t + X$ cross section, by performing a binned




FIG. 2 (color online). Reconstructed transverse mass of the W , m_T^W , in signal-depleted control regions. Left, events with at least four jets and small missing transverse momentum ($< 100 \text{ GeV}/c$). Right, events with exactly three jets and large missing transverse momentum ($> 100 \text{ GeV}/c$).

TABLE II. Expected 95% C.L. upper limit on $T'\bar{T}'$ production cross section, σ_{exp} , the range of expected limits which includes 68% of pseudoexperiments, and the observed limit, σ_{obs} , for representative signal points in $(m_{T'}, m_X)$.

$m_{T'}, m_X$ (GeV/ c^2)	σ_{exp} [pb]	+34%	-34%	σ_{obs} [pb]
200, 1	1.31	1.86	0.83	1.21
220, 40	1.40	2.17	0.93	1.20
260, 1	0.23	0.40	0.14	0.20
280, 1	0.16	0.27	0.09	0.15
280, 20	0.18	0.29	0.11	0.17
280, 40	0.17	0.27	0.11	0.12
$m_{T'}, m_X$ (GeV/ c^2)	σ_{exp} [pb]	+34%	-34%	σ_{obs} [pb]
300, 100	0.34	0.51	0.24	0.39
310, 90	0.19	0.29	0.11	0.21
320, 80	0.15	0.24	0.08	0.12
350, 50	0.07	0.10	0.04	0.02
360, 110	0.09	0.19	0.05	0.09
370, 1	0.06	0.10	0.04	0.05

maximum-likelihood fit in the $m_{T'}^W$ variable, allowing for systematic and statistical fluctuations via template morphing [19] which performs an interpolation in each bin as a function of the nuisance parameters. We use the likelihood-ratio ordering prescription [20] to construct classical confidence intervals in the theoretical cross section by generating ensembles of simulated experiments that describe expected fluctuations of statistical and systematic uncertainties on both signal and backgrounds. The observed limits are consistent with expectation in the background-only hypothesis, for a few example signal mass points we tabulate the expected and observed limits (see Table II). We convert the observed upper limits on the pair-production cross sections to an exclusion curve in mass parameter space for the dark matter model involving fourth generation quarks; see Fig. 3.

In conclusion, we have searched for new physics particles T' decaying to top quarks with invisible particles X with a detector signature of $t\bar{t} + E_\tau$. We calculate upper




FIG. 3. Observed versus expected exclusion in $(m_{T'}, m_X)$ along with the cross-section upper limits.

limits on the cross section of such events and exclude any dark matter model involving exotic fourth generation quark up to $m_{T'} = 360$ GeV/ c^2 . Our cross-section limits on the generic decay, $T' \rightarrow t + X$, may be applied to the many other models that predict the production of a heavy particle T' decaying to top quarks and invisible particles X , such as the supersymmetric process $\tilde{t} \rightarrow t + \chi^0$. A similar search performed at the LHC, given its higher energy regime, would be able to provide limits on such a supersymmetric decay.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

^aDeceased.

^bWith visitors from University of Massachusetts Amherst, Amherst, MA 01003, USA.

^cWith visitors from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy.

^dWith visitors from University of California Irvine, Irvine, CA 92697, USA.

^eWith visitors from University of California Santa Barbara, Santa Barbara, CA 93106, USA.

^fWith visitors from University of California Santa Cruz, Santa Cruz, CA 95064, USA.

^gWith visitors from CERN, CH-1211 Geneva, Switzerland.

^hWith visitors from Cornell University, Ithaca, NY 14853, USA.

ⁱWith visitors from University of Cyprus, Nicosia CY-1678, Cyprus.

^jWith visitors from University College Dublin, Dublin 4, Ireland.

^kWith visitors from University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017.

^lWith visitors from Universidad Iberoamericana, Mexico D.F., Mexico.

^mWith visitors from Iowa State University, Ames, IA 50011, USA.

ⁿWith visitors from University of Iowa, Iowa City, IA 52242, USA.

- ^oWith visitors from Kinki University, Higashi-Osaka City, Japan 577-8502.
- ^pWith visitors from Kansas State University, Manhattan, KS 66506, USA.
- ^qWith visitors from University of Manchester, Manchester M13 9PL, England.
- ^rWith visitors from Queen Mary, University of London, London, E1 4NS, England.
- ^sWith visitors from University of Melbourne, Victoria 3010, Australia.
- ^tWith visitors from Muons, Inc., Batavia, IL 60510, USA.
- ^uWith visitors from Nagasaki Institute of Applied Science, Nagasaki, Japan.
- ^vWith visitors from National Research Nuclear University, Moscow, Russia.
- ^wWith visitors from University of Notre Dame, Notre Dame, IN 46556, USA.
- ^xWith visitors from Universidad de Oviedo, E-33007 Oviedo, Spain.
- ^yWith visitors from Texas Tech University, Lubbock, TX 79609, USA.
- ^zWith visitors from Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile.
- ^{aa}With visitors from Yarmouk University, Irbid 211-63, Jordan.
- ^{bb}On leave from J. Stefan Institute, Ljubljana, Slovenia.
- [1] J. Feng, *Annu. Rev. Astron. Astrophys.* **48**, 495, (2010).
- [2] J. Feng *et al.*, *Phys. Rev. D* **81**, 114027 (2010).
- [3] H. C. Cheng and I. Low, *J. High Energy Phys.* **08** (2004) 061.
- [4] P. Fileviez Perez and M. B. Wise, *Phys. Rev. D* **82**, 011901 (2010).
- [5] M. Drees, R. Godbole, and P. Roy (World Scientific, Hackensack, 2004), p. 555.
- [6] D. Acosta *et al.* (The CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005).
- [7] A. Abulencia *et al.* (The CDF Collaboration), *Phys. Rev. Lett.* **97**, 082004 (2006). D. Acosta *et al.* (The CDF Collaboration), *Phys. Rev. Lett.* **94**, 091803 (2005).
- [8] CDF uses a cylindrical coordinate system with the z axis along the proton beam axis. Pseudorapidity is $\eta \equiv -\ln(\tan(\theta/2))$, where θ is the polar angle relative to the proton beam direction, and ϕ is the azimuthal angle while $p_T = |p| \sin\theta$, $E_T = E \sin\theta$.
- [9] F. Abe *et al.* (The CDF Collaboration), *Phys. Rev. D* **45**, 1448 (1992).
- [10] A. Bhatti *et al.*, *Nucl. Instrum. Methods* **566**, 375 (2006).
- [11] Missing transverse momentum, E_T , is defined as the magnitude of the vector $-\sum_i E_T^i \vec{n}_i$ where E_T^i are the magnitudes of transverse energy contained in each calorimeter tower i , and \vec{n}_i is the unit vector from the interaction vertex to the tower in the transverse (x, y) plane.
- [12] J. Alwall *et al.*, *J. High Energy Phys.* **09** (2007) 028, version 4.4.24.
- [13] T. Sjostrand *et al.*, *Comput. Phys. Commun.* **238**, 135 (2001), version 6.422.
- [14] R. Bonciani, S. Catani, M.L. Mangano, and P. Nason, *Nucl. Phys.* **B529**, 424 (1998).
- [15] M. Mangano *et al.*, *J. High Energy Phys.* **07** (2003) 001.
- [16] A. Abulencia *et al.* (The CDF Collaboration), *Phys. Rev. D* **74**, 072006 (2006).
- [17] J.M. Campbell and R. K. Ellis, *Phys. Rev. D* **60**, 113006 (1999).
- [18] A. Abulencia *et al.* (The CDF Collaboration), *Phys. Rev. D* **73**, 32003 (2006); J. Pumplin *et al.* (The CTEQ Collaboration), *J. High Energy Phys.* **07** (2002) 012; A.D. Martin *et al.* (The MRST Collaboration), *Phys. Lett. B* **356**, 89 (1995).
- [19] A. Read, *Nucl. Instrum. Methods Phys. Res., Sect. A* **425**, 357 (1999).
- [20] G.J. Feldman and R.D. Cousins, *Phys. Rev. D* **57**, 3873 (1998); M. Cacciari, S. Frixione, M.L. Mangano, P. Nason, and G. Ridolfi, *J. High Energy Phys.* **04** (2004) 068.